



The size-weight illusion induced through human echolocation

Journal:	<i>Psychological Science</i>
Manuscript ID:	PSCI-14-1032.R1
Manuscript Type:	Research report
Date Submitted by the Author:	n/a
Complete List of Authors:	Buckingham, Gavin; Heriot-Watt University, Psychology Milne, Jennifer; University of Western Ontario, Brain and Mind Institute Byrne, Caitlin; University of Western Ontario, Brain and Mind Institute Goodale, Melvyn; University of Western Ontario, The Brain and Mind Institute
Keywords:	Perception, Perceptual Motor Coordination

SCHOLARONE™
Manuscripts

Only

1
2
3 The size-weight illusion induced through human echolocation
4
5
6

7 G. Buckingham^{1,2}
8

9 J. L. Milne²
10

11 C. M. Byrne²
12

13 M.A. Goodale²
14
15
16

17 1. Department of Psychology, School of Life Sciences, Heriot-Watt University, Edinburgh, UK
18

19 2. The Brain and Mind Institute, Department of Psychology, Western University, Ontario, Canada
20
21
22

23 Corresponding author:
24

25 Gavin Buckingham
26

27 David Brewster Building
28

29 School of Life Sciences
30

31 Heriot-Watt University
32

33 Edinburgh, UK
34

35 EH14 4AS
36

37 Phone: +44 131 451 3643
38

39 Email: g.buckingham@hw.ac.uk
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3 Certain blind individuals have learned to interpret the echoes of self-generated sounds to perceive
4 the structure of objects in their environment. The current work sought to examine how far the
5 influence of this unique form of visual substitution extends. In order to determine whether
6 echolocation-induced representations of object size could influence weight perception, a small
7 group of echolocation experts made tongue-clicks or finger-snaps toward cubes of varying sizes and
8 weights before lifting them. The echolocators experienced a robust size-weight illusion, providing
9 the first demonstration of a sensory substitution technique where the substituted sense influences
10 the conscious perception of an intact sense.
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

For Review Only

Introduction

Echolocation, the process of using the echoes from self-generated acoustic signals in order to navigate, is seen in a number of non-primate species. This skill is one which can be learned - a number of blind humans have taught themselves to echolocate to replace vision (for a recent review see Kolarik, Cirstea, Pardhan, & Moore, 2014). Typically, these individuals use their echolocation ability to process the spatial layout of their environment in order to facilitate successful locomotion (Thaler, 2013). Echolocation is not, however, merely a functional tool to facilitate navigation. Recent work has demonstrated that human echolocators can decode echoes to identify a variety of object properties such as size, shape, and distance (Milne, Anello, Goodale, & Thaler, in press; Milne, Goodale, & Thaler, 2014; Teng, Puri, & Whitney, 2012). This unique self-generated form of sensory substitution appears to lead to cortical reorganization, with blind echolocation experts recruiting regions of primary visual cortex when making echo-based judgements about object identity (Thaler, Arnott, & Goodale, 2011). Furthermore, recent research has shown that echolocation appears to have vision-like properties such as size constancy – the integration of size and distance cues to maintain an accurate percept of object size in spite of variations in the echo equivalent of retinal size (i.e., stimulus strength and aperture, Milne et al., in press). The current work examines whether this unique form of sensory substitution can influence our perceptions of other senses, specifically in the context of heaviness perception and weight illusions.

It is well known that one's perception of object mass can be influenced by an object's appearance. For example, in the classic 'size-weight illusion' (SWI - Charpentier, 1891; Ross, 1969), small objects feel heavier than equally-weighted large objects. This powerful illusion does not require a haptic sense of object size, and can be induced entirely with visual cues (e.g., when objects are lifted with identically-sized handles or string-and-pulley systems - see Buckingham & Goodale, 2010b; Ellis & Lederman, 1993; Masin & Crestoni, 1988). Initial explanations for the SWI proposed that the effect stemmed from efferent/afferent mismatches (Davis & Roberts, 1976), although recent work examining fingertip force control when lifting illusion-inducing objects has rendered such sensorimotor explanations untenable (Flanagan & Beltzner, 2000, although see Dijk, 2014).

The current view is that the SWI is a consequence of a lifter's cognitive expectations of heaviness, based on their implicit understanding of the relationship between object volume and object mass (for review, see Buckingham, in press). Continuous experience of an object's volume is also not necessary to experience the SWI – a short visual preview of object size is sufficient to induce a robust weight illusion when lifting without visual feedback (Buckingham & Goodale, 2010a; Buckingham, Ranger, & Goodale, 2011). The goal of the current work was to determine whether the percept of object size extracted during human echolocation is sufficiently powerful and reliable to influence an echolocator's experience of object mass and induce a SWI. To this end, we examined perceptions of heaviness in a small group of human echolocators as they lifted sets of objects with different sizes, but identical weights.

Materials and methods

Three blind echolocators were recruited from Ontario, Canada to take part in an object lifting task. The blind echolocation group was made up of one 22 year old female and two males aged 49 and 57. All of these echolocators use echolocation techniques throughout their daily lives for navigation during locomotion. The female echocator lost her sight in adolescence and uses a finger-snap as the sound source for creating echoes; the two males were blinded early in life and use a tongue-click. Their perception of real and illusory weight differences was compared to a group of three non-echolocating blind individuals (two males, one female, mean age = 36.3 years, st. dev = 7.8) and four sighted controls (one male, three females, mean age= 20.5 years, st. dev = 1.1).

Testing took place at the University of Western Ontario in a small (width: 275 cm, length: 300 cm, height: 244 cm) room with walls which were covered with 3.8 cm thick convoluted foam sheets, to dampen echoes. All blind participants wore a blindfold throughout the task, whereas the sighted participants were permitted visual feedback during the entire task except for the periods between trials when the objects were changed over. All participants gave informed consent, and all testing procedures were approved by the university ethics board.

Two sets of these cubes were created - one set was adjusted to weigh 1.5kg (light objects) and another set was adjusted to weigh 2kg (heavy objects). Each set was made up of three cubes – small (15 x 15 x 15 cm), medium (35 x 35 x 35 cm), and large (55 x 55 x 55 cm). The cubes, which were made from 50 mm thick white foam core board, had wooden panels fixed onto the inner panel of their top surfaces to increase their weight and facilitate the attachment of a hook (Figure 1). In order to minimise distracting echoes from the room, only one cube was presented to participants at a time, with the remaining five cubes hidden behind a felt curtain in the corner of the room.




Figure 1. The light set of small, medium, and large cubes lifted by participants in this study. Participants also interacted with a heavier set of identical-looking cubes.

1
2
3 In order to determine if echolocation could induce a SWI, it was essential to ensure that participants
4 received no cues to object size other than those derived through echoes. To this end, we developed
5 a modified version of the task used by Masin and Crestoni (1988), where objects were lifted via a
6 cord and pulley system. All the cubes had a small brass loop screwed into the centre of their top
7 surface, through which a cord was threaded. This cord, which was 329 cm long, was threaded
8 through a small pulley in the ceiling 94 cm directly above the cube, and attached to the wall of the
9 room 87 cm from the ground. The cubes were placed in the centre of the room atop a free-standing
10 pedestal at head height (see Milne et al., 2014; Thaler et al., 2011 for descriptions of similar
11 experimental setups). The height of the pedestal was adjusted for each cube such that the top
12 surface of each object was always 150 cm from the floor and the cable connected to the box via the
13 pulley always maintained the same tension. Participants lifted the objects and made their weight
14 judgements by moving to the edge of the room where the cable was fixed, placing their hands on
15 the cable and pushing it downwards in order to lift the cube off of its pedestal. This procedure
16 eliminated any differences between the cubes' haptically-perceived size and rotational torques,
17 ensuring that subjects' impressions of object volume would stem entirely from echolocation- or
18 vision-derived cues.
19
20
21
22

23
24 Prior to commencing the experimental trials, participants lifted the medium-sized cube of the heavy
25 set of stimuli five times in a row, which they were told weighed 50 out of 100 and on which they
26 were to base their subsequent perceptual judgements. On each trial, participants in all groups were
27 allowed unlimited time and whatever method they wished (without touching the object) to try to
28 determine the size of the cube. Typically, the individuals in the echolocation group made tongue-
29 clicks (the two early-blind males) or finger-snaps (the late-blind female) toward the cube, identifying
30 its edges for 5-15 seconds. The sighted controls looked briefly at the cubes, and the blind non-
31 echolocators used tongue-clicks and hand claps. Once they had finished 'viewing' the cube, subjects
32 retreated to the edge of the room where the wire was affixed to the wall and lifted the object with
33 the cable and pulley system by pushing down on the wire. Once they had lifted the cube,
34 participants gave a numerical rating of how heavy the object felt, relative to the medium-sized
35 object of the heavy set of cubes. A video showing the trial procedure and experimental setup can be
36 found in the supplementary materials. Participants lifted each of the six cubes six times apiece, for a
37 total of 36 lifts. The procedure took between 30 and 60 minutes to complete, depending on the
38 strategy adopted by the participant.
39
40
41
42

43 To examine how each group experienced object weight as a function of size, the ratings given to the
44 small, medium, and large objects were collapsed across the 1.5kg and 2kg sets of objects. To
45 combine the ratings given to the heavy and light objects, the ratings given on each trial were
46 transformed into a proportion of the highest value given by each subject in each set. ANOVA and
47 post hoc analyses were run with IBM SPSS Statistics (Version 21), and effect size calculations were
48 performed in Microsoft Excel with Lakens' (2013) open science spreadsheet.
49
50
51
52
53
54
55
56
57
58
59
60

Results

The combined normalized ratings of heaviness (Figure 2) were first examined in a mixed design 3 (object size) \times 3 (group) ANOVA. This omnibus analysis revealed a significant main effect of object size ($F(2,14) = 18.39, p < .001, \eta_p^2 = 0.72$) as well as a significant interaction between object size and group ($F(4,14) = 9.57, p < .005, \eta_p^2 = 0.73$).




Figure 2. The average normalized ratings of heaviness (averaged across object weight) given by each group for the small, medium, and large cubes. Error bars indicate standard error of the mean.

In order to examine the presence or absence of a perceptual illusion, the magnitude of the SWI was calculated for each individual by subtracting the rating they gave to the large cube from the rating they gave to the small cube (Figure 3A). This value, representing the magnitude of the illusory weight difference, was then compared to zero with one-tailed, one-sample t-tests. Both the echolocation ($t(2) = 3.4, p < .05$) and the sighted ($t(3) = 8.8, p < .005$) groups showed a significant SWI, whereas the non-echolocating blind control group showed no such effect ($t(2) = .66, p = .57$). In order to explicitly examine how the SWI varied across the groups, we performed a one way ANOVA, followed by LSD post hoc comparisons. The ANOVA indicated that significant differences existed between the groups ($F(2,7) = 19.45, p < .005, \eta_p^2 = 0.85$). Post hoc analyses confirmed the sighted group showed a larger illusion than the blind control group (mean difference = 33.6, $p < .001$) and the echolocation group (mean difference = 18, $p < .05$). Crucially, however, the echolocation group also showed a larger illusion than the blind control group (mean difference = 15.6, $p < .05$). By contrast, a one way ANOVA comparing the average difference in raw ratings, collapsed across size, given to the heavy (2kg) and light (1.5kg) sets of cubes found no indication that the groups differed in how they perceived real differences in object mass ($F(2,7) = 4.09, p = .07, \eta_p^2 = 0.54$, Figure 3B).




Figure 3. (A) The average rating given to the large objects subtracted from the average rating given to the small objects (collapsed across both object weights) and (B) the average rating given to the light set of objects subtracted from the average rating given to the heavy set of objects (collapsed across all three object sizes). Error bars indicate standard error of the mean.

Discussion

The current work investigated the extent of the perceptual capabilities of human echolocators in the context of the size-weight illusion (SWI). A small sample of blind individuals who regularly use echolocation for navigation made self-generated tongue-clicks or finger-snaps toward a variety of cubes which were balanced at head height atop a free-standing pedestal. Once they had echolocated toward a cube, they reported how heavy it felt after lifting it off the pedestal with a rope and pulley system. The echolocators' perceptions of object weight was compared to a small group of sighted individuals, who lifted the objects with full vision, as well as a small group of non-echolocating blind individuals who received no cues to object size. When judging how heavy the differently-sized cubes felt, the non-echolocating blind individuals reported (unsurprisingly) that the identically-weighted objects felt as if they weighed the same amount as one another. By contrast, the echolocating blind individuals experienced a SWI, reporting that the smallest cubes felt significantly heavier than the largest cubes. Interestingly, the illusion experienced by the echolocators was significantly smaller than that experienced by the sighted controls, who experienced a more robust SWI (presumably due to the higher resolution of vision as compared to echolocation). By contrast, no statistical differences were observed between how each of the groups experienced a 500g weight variation. The numerical trend for the sighted participants to report that weight difference as smaller than the non-sighted groups is consistent with earlier reports that blind individuals show a greater sensitivity for weight discrimination than sighted counterparts (Grouios, Alevraidou, & Koidou, 2001)

It is well-established that blind individuals can experience a robust SWI when given haptic feedback of an object's volume (Ellis & Lederman, 1993). However, no experiment to date has reported blind individuals experiencing this effect through any other modality. The current findings give new insights into the perceptual abilities of blind humans who have taught themselves to echolocate. It has been well-reported that blind echolocators have a wide repertoire of unique perceptual skills which allows them to successfully interact with the world around them (Kolarik et al., 2014). This technique does not merely solve the same spatial mapping problems that visual feedback normally deals with, but shows a range of vision-like properties (Milne et al., in press; Thaler et al., 2011).

1
2
3 Given that other substitutive techniques have been shown to have an automatic influence on
4 sensorimotor control (e.g., Danilov, Tyler, Skinner, Hogle, & Bach-y-Rita, 2007), it is likely that
5 echolocation experts would also tend to use this unique perceptual ability to apply initial grip and
6 load forces to objects based upon their size and material properties (Buckingham, Cant, & Goodale,
7 2009; Flanagan & Beltzner, 2000; Gordon, Forssberg, Johansson, & Westling, 1991). However, the
8 current work has shown that the echolocation-derived representations of object size influence the
9 conscious perception of the ostensibly unrelated variable of object weight. This is, to our knowledge,
10 the first example of a substituted sense influencing the conscious perception of an intact sense
11 (touch), speaking to this technique's efficacy as a full (i.e., not merely functional) form of sensory
12 substitution.
13
14
15

16 To conclude, our findings highlight the degree of confidence that echolocators place in their echo-
17 induced judgements of object properties, indicating that human echolocation has the potential to
18 replace not only the functional aspects of vision, but also individuals' conscious perception of visual
19 aspects of the world.
20
21
22
23

24 Author contributions

25
26 G. Buckingham, J. Milne, and M. Goodale developed the study concept. Testing and data collection
27 were undertaken by C. Byrne, and supervised by J. Milne and M. Goodale. Analysis and
28 interpretation was undertaken by G. Buckingham, who also drafted the manuscript. All authors
29 approved the final version of the manuscript for submission.
30
31
32
33

34 Funding

35
36 The first author was supported with an NSERC Banting Postdoctoral Fellowship.
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3 References
4

- 5 Buckingham, G. (in press). Getting a grip on heaviness perception: a review of weight illusions and
6
7 their probable causes. *Experimental Brain Research*, 1–7. doi:10.1007/s00221-014-3926-9
8
- 9 Buckingham, G., Cant, J. S., & Goodale, M. A. (2009). Living in A Material World: How Visual Cues to
10
11 Material Properties Affect the Way That We Lift Objects and Perceive Their Weight. *Journal*
12
13 *of Neurophysiology*, 102(6), 3111–3118. doi:10.1152/jn.00515.2009
14
- 15 Buckingham, G., & Goodale, M. A. (2010a). Lifting without Seeing: The Role of Vision in Perceiving
16
17 and Acting upon the Size Weight Illusion. *PLoS ONE*, 5(3), e9709.
18
19 doi:10.1371/journal.pone.0009709
20
- 21 Buckingham, G., & Goodale, M. A. (2010b). The influence of competing perceptual and motor priors
22
23 in the context of the size–weight illusion. *Experimental Brain Research*, 205(2), 283–288.
24
25 doi:10.1007/s00221-010-2353-9
26
- 27 Buckingham, G., Ranger, N. S., & Goodale, M. A. (2011). The Role of Vision in Detecting and
28
29 Correcting Fingertip Force Errors During Object Lifting. *Journal of Vision*, 11(1).
30
31 doi:10.1167/11.1.4
32
- 33 Charpentier, A. (1891). Analyse expérimentale quelques éléments de la sensation de poids. *Archives*
34
35 *de Physiologie Normales et Pathologiques*, 3, 122–135.
36
37
- 38 Danilov, Y. P., Tyler, M. E., Skinner, K. L., Hogle, R. A., & Bach-y-Rita, P. (2007). Efficacy of
39
40 electrotactile vestibular substitution in patients with peripheral and central vestibular loss.
41
42 *Journal of Vestibular Research: Equilibrium & Orientation*, 17(2-3), 119–130.
43
44
- 45 Davis, C. ., & Roberts, W. (1976). Lifting movements in the size-weight illusion. *Perception &*
46
47 *Psychophysics*, 20, 33–36.
48
49
- 50 Dijker, A. J. M. (2014). The role of expectancies in the size-weight illusion: A review of theoretical
51
52 and empirical arguments and a new explanation. *Psychonomic Bulletin & Review*, 1–11.
53
54
55 doi:10.3758/s13423-014-0634-1
56
57
58
59
60

- 1
2
3 Ellis, R. R., & Lederman, S. J. (1993). The role of haptic versus visual volume cues in the size-weight
4
5 illusion. *Perception & Psychophysics*, *53*(3), 315–324.
6
7 Flanagan, J. R., & Beltzner, M. A. (2000). Independence of perceptual and sensorimotor predictions
8
9 in the size-weight illusion. *Nature Neuroscience*, *3*(7), 737–741. doi:10.1038/76701
10
11 Gordon, A. M., Forssberg, H., Johansson, R. S., & Westling, G. (1991). Visual size cues in the
12
13 programming of manipulative forces during precision grip. *Experimental Brain Research*,
14
15 *83*(3), 477–482.
16
17 Grouios, G., Alevraidou, A., & Koidou, I. (2001). Weight-Discrimination Sensitivity in Congenitally
18
19 Blind and Sighted Adults. *Journal of Visual Impairment & Blindness*, *95*(1), 30.
20
21 Kolarik, A. J., Cirstea, S., Pardhan, S., & Moore, B. C. J. (2014). A summary of research investigating
22
23 echolocation abilities of blind and sighted humans. *Hearing Research*, *310*, 60–68.
24
25 doi:10.1016/j.heares.2014.01.010
26
27
28
29 Lakens, D. (2013). Calculating and reporting effect sizes to facilitate cumulative science: a practical
30
31 primer for t-tests and ANOVAs. *Frontiers in Psychology*, *4*, 863.
32
33 doi:10.3389/fpsyg.2013.00863
34
35 Masin, S. C., & Crestoni, L. (1988). Experimental demonstration of the sensory basis of the size-
36
37 weight illusion. *Perception & Psychophysics*, *44*(4), 309–312.
38
39 Milne, J. L., Anello, M., Goodale, M. A., & Thaler, L. (in press). A blind human expert echolocator
40
41 shows size constancy for objects perceived by echoes. *Neurocase*, *0*(0), 1–6.
42
43 doi:10.1080/13554794.2014.922994
44
45
46 Milne, J. L., Goodale, M. A., & Thaler, L. (2014). The role of head movements in the discrimination of
47
48 2-D shape by blind echolocation experts. *Attention, Perception, & Psychophysics*, 1–10.
49
50 doi:10.3758/s13414-014-0695-2
51
52
53 Ross, H. E. (1969). When is a weight not illusory? *The Quarterly Journal of Experimental Psychology*,
54
55 *21*(4), 346–355. doi:10.1080/14640746908400230
56
57
58
59
60

1
2
3 Teng, S., Puri, A., & Whitney, D. (2012). Ultrafine spatial acuity of blind expert human echolocators.

4
5 *Experimental Brain Research*, 216(4), 483–488. doi:10.1007/s00221-011-2951-1

6
7 Thaler, L. (2013). Echolocation may have real-life advantages for blind people: an analysis of survey

8
9 data. *Integrative Physiology*, 4, 98. doi:10.3389/fphys.2013.00098

10
11 Thaler, L., Arnott, S. R., & Goodale, M. A. (2011). Neural correlates of natural human echolocation in
12 early and late blind echolocation experts. *PLoS One*, 6(5), e20162.

13
14
15 doi:10.1371/journal.pone.0020162

16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

For Review Only




Figure 1. The light set of small, medium, and large cubes lifted by participants in this study. Participants also interacted with a heavier set of identical-looking cubes.

94x51mm (300 x 300 DPI)

Review Only

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60




Figure 2. The average normalized ratings of heaviness (averaged across object weight) given by each group for the small, medium, and large cubes. Error bars indicate standard error of the mean.
291x100mm (96 x 96 DPI)

Review Only




Figure 3. (A) The average rating given to the large objects subtracted from the average rating given to the small objects (collapsed across both object weights) and (B) the average rating given to the light set of objects subtracted from the average rating given to the heavy set of objects (collapsed across all three object sizes). Error bars indicate standard error of the mean
370x118mm (96 x 96 DPI)

Review Only

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60