

# Draft Genome Sequences of *Achromobacter piechaudii* GCS2, *Agrobacterium* sp. Strain SUL3, *Microbacterium* sp. Strain GCS4, *Shinella* sp. Strain GWS1, and *Shinella* sp. Strain SUS2 Isolated from Consortium with the Hydrocarbon-Producing Alga *Botryococcus braunii*

Katy J. Jones, Karen Moore,  Christine Sambles, John Love,  David J. Studholme, Stephen J. Aves

Department of Biosciences, College of Life and Environmental Sciences, University of Exeter, Exeter, United Kingdom

**A variety of bacteria associate with the hydrocarbon-producing microalga *Botryococcus braunii*, some of which may influence its growth. We report here the genome sequences for *Achromobacter piechaudii* GCS2, *Agrobacterium* sp. strain SUL3, *Microbacterium* sp. strain GCS4, and *Shinella* sp. strains GWS1 and SUS2, isolated from a laboratory culture of *B. braunii*, race B, strain Guadeloupe.**

Received 6 November 2015 Accepted 9 November 2015 Published 14 January 2016

**Citation** Jones KJ, Moore K, Sambles C, Love J, Studholme DJ, Aves SJ. 2016. Draft genome sequences of *Achromobacter piechaudii* GCS2, *Agrobacterium* sp. strain SUL3, *Microbacterium* sp. strain GCS4, *Shinella* sp. strain GWS1, and *Shinella* sp. strain SUS2 isolated from consortium with the hydrocarbon-producing alga *Botryococcus braunii*. *Genome Announc* 4(1):e01527-15. doi:10.1128/genomeA.01527-15.

**Copyright** © 2016 Jones et al. This is an open-access article distributed under the terms of the [Creative Commons Attribution 3.0 Unported license](http://creativecommons.org/licenses/by/3.0/).

Address correspondence to Stephen J. Aves, [sj.aves@exeter.ac.uk](mailto:sj.aves@exeter.ac.uk).

*Botryococcus braunii* is a green microalga that is able to produce and accumulate high levels of hydrocarbons (up to 60% of its dry weight, depending on the strain), which can be converted into liquid fuels (1, 2). Strains of *B. braunii* are typically not axenic but consist of a single algal strain accompanied by a variety of microorganisms associated as both a biofilm and planktonic population in the water column, which may greatly influence *B. braunii* growth and hydrocarbon production (3–6).

Five bacterial strains (GCS2, SUL3, GCS4, GWS1, and SUS2) were isolated from a long-term laboratory culture of *B. braunii* strain Guadeloupe (race B) (7), and the colonies were grown on LB plates. Genomic DNA fragmented by sonication was concentrated and purified using a QIAquick column (Qiagen), and genomic libraries were prepared using the NEBNext DNA Library Prep Master Mix Set for Illumina (New England BioLabs). One hundred fifty-base pair paired-end sequencing, with a custom bar code, was carried out on an Illumina MiSeq, and *de novo* assembly of the sequence data was carried out using SPAdes version 3.5.0 (8). The number of contigs, coverage,  $N_{50}$ , and total sequence length were, respectively, 20, 41×, 626,645, and 6,180,134 bp for *Achromobacter piechaudii* GCS2; 72, 52×, 358,737, and 6,013,863 bp for *Agrobacterium* sp. strain SUL3; 7, 73×, 944,354, and 3,652,908 bp for *Microbacterium* sp. strain GCS4; 141, 44×, 176,944, and 6,989,725 bp for *Shinella* sp. strain GWS1; and 112, 32×, 264,748, and 7,005,767 bp for *Shinella* sp. strain SUS2. The genomes were annotated using the NCBI Prokaryotic Genome Annotation Pipeline (PGAP), which predicted 5,433, 5,585, 3,325, 6,636, and 6,361 coding sequences (CDSs), respectively, for the five strains. The G+C contents varied between 59.2% (*Agrobacterium* sp. SUL3) and 69.5% (*Microbacterium* sp. GCS4). The 16S rRNA gene sequences were extracted using RNAmmer (9).

Phylogenetic analysis of 16S rRNA gene sequence, *atpD*, *recA*, *rpoB*, and *tyrB* suggested that GCS2 belongs to the species *A. piechaudii*; its genome shares 98.4% average nucleotide identity (ANI) with that of *A. piechaudii* strain ATCC 43553. Phylogenetic analysis of 16S rRNA gene sequence, *gyrB*, *ppk*, *recA*, and *rpoB* suggested that GCS4 belongs to the genus *Microbacterium*. Phylogenetic analysis of 16S rRNA gene sequence, *atpD*, and *recA* suggested that GWS1 and SUS2 both belong to the genus *Shinella*, and SUL3 belongs to the genus *Agrobacterium*.

None of these five bacteria were previously reported in association with *B. braunii*, except for *Shinella* sp. GWS1 and SUS2, which, on the basis of partial 16S rRNA gene sequence, are indistinguishable from a biofilm isolate found to enhance the growth of a laboratory-maintained monoculture of *B. braunii* (4) (reported as *Rhizobium* sp. M14). Many species of algae are auxotrophic for vitamin B<sub>1</sub> (thiamine), vitamin B<sub>7</sub> (biotin), and vitamin B<sub>12</sub> (cobalamin) (10). We identified genes encoding pathways for thiamine synthesis in all five genomes; cobalamin is likely to be synthesized by *Agrobacterium* sp. SUL3 and *Shinella* spp. GWS1 and SUS2, while biotin is likely to be synthesized by *A. piechaudii* GCS2 and *Agrobacterium* sp. SUL3. No evidence of nitrogen fixation genes was found in any of the bacterial genomes, based on tBLASTn and BLASTn searches.

**Nucleotide sequence accession numbers.** These genome sequences have been deposited at GenBank under the following accession numbers: *A. piechaudii* GCS2, [LGYD000000000](https://www.ncbi.nlm.nih.gov/nuccore/LGYD000000000); *Agrobacterium* sp. SUL3, [LGZB000000000](https://www.ncbi.nlm.nih.gov/nuccore/LGZB000000000); *Microbacterium* sp. GCS4, [LGYE000000000](https://www.ncbi.nlm.nih.gov/nuccore/LGYE000000000); *Shinella* sp. GWS1, [LGYF000000000](https://www.ncbi.nlm.nih.gov/nuccore/LGYF000000000); and *Shinella* sp. SUS2, [LGYG000000000](https://www.ncbi.nlm.nih.gov/nuccore/LGYG000000000). The versions described in this paper are the first versions: [LGYD010000000](https://www.ncbi.nlm.nih.gov/nuccore/LGYD010000000), [LGZB010000000](https://www.ncbi.nlm.nih.gov/nuccore/LGZB010000000), [LGYE010000000](https://www.ncbi.nlm.nih.gov/nuccore/LGYE010000000), [LGYF010000000](https://www.ncbi.nlm.nih.gov/nuccore/LGYF010000000), and [LGYG010000000](https://www.ncbi.nlm.nih.gov/nuccore/LGYG010000000).

## ACKNOWLEDGMENTS

K.J.J. was funded by a Ph.D. studentship from the Biotechnology and Biological Sciences Research Council (BBSRC) LoLa award BB/K003240/2. The Exeter Sequencing Service was supported by the Wellcome Trust Institutional Strategic Support Fund (WT097835MF), a Wellcome Trust Multi User Equipment Award (WT101650MA), and a BBSRC LoLa award (BB/K003240/2).

We gratefully acknowledge the Exeter Sequencing Service and computational core facilities at the University of Exeter.

## FUNDING INFORMATION

The Wellcome Trust provided funding to Karen Moore under grant numbers WT097835MF and WT101650MA. The Biotechnology and Biological Sciences Research Council (BBSRC) provided funding to Katy J. Jones and Karen Moore under grant number BB/K003240/2.

The funders had no role in study design, data collection and interpretation, or the decision to submit the work for publication.

## REFERENCES

1. Banerjee A, Sharma R, Chisti Y, Banerjee UC. 2002. *Botryococcus braunii*: a renewable source of hydrocarbons and other chemicals. *Crit Rev Biotechnol* 22:245–279. <http://dx.doi.org/10.1080/0738850290789513>.
2. Metzger P, Largeau C. 2005. *Botryococcus braunii*: a rich source for hydrocarbons and related ether lipids. *Appl Microbiol Biotechnol* 66: 486–496. <http://dx.doi.org/10.1007/s00253-004-1779-z>.
3. Chirac C, Casadevall E, Largeau C, Metzger P. 1985. Bacterial influence upon growth and hydrocarbon production of the green alga *Botryococcus braunii*. *J Phycol* 21:380–387.
4. Rivas MO, Vargas P, Riquelme CE. 2010. Interactions of *Botryococcus braunii* cultures with bacterial biofilms. *Microb Ecol* 60:628–635. <http://dx.doi.org/10.1007/s00248-010-9686-6>.
5. Tanabe Y, Kato S, Matsuura H, Watanabe MM. 2012. A *Botryococcus* strain with bacterial ectosymbionts grows fast and produces high amount of hydrocarbons. *Procedia Environ Sci* 15:22–26. <http://dx.doi.org/10.1016/j.proenv.2012.05.005>.
6. Tanabe Y, Okazaki Y, Yoshida M, Matsuura H, Kai A, Shiratori T, Ishida K, Nakano S, Watanabe MM. 2015. A novel alphaproteobacterial ectosymbiont promotes the growth of the hydrocarbon-rich green alga *Botryococcus braunii*. *Sci Rep* 5:10467. <http://dx.doi.org/10.1038/srep10467>.
7. Metzger P, Berkaloff C, Casadevall E, Coute A. 1985. Alkadiene- and botryococcene-producing races of wild strains of *Botryococcus braunii*. *Phytochemistry* 24:2305–2312. [http://dx.doi.org/10.1016/S0031-9422\(00\)83032-0](http://dx.doi.org/10.1016/S0031-9422(00)83032-0).
8. Bankevich A, Nurk S, Antipov D, Gurevich AA, Dvorkin M, Kulikov AS, Lesin VM, Nikolenko SI, Pham S, Pribelski AD, Pyshtkin AV, Sirotkin AV, Vyahhi N, Tesler G, Alekseyev MA, Pevzner PA. 2012. SPAdes: a new genome assembly algorithm and its applications to single-cell sequencing. *J Comput Biol* 19:455–477. <http://dx.doi.org/10.1089/cmb.2012.0021>.
9. Lagesen K, Hallin P, Rødland EA, Staerfeldt HH, Rognes T, Ussery DW. 2007. RNAmmer: consistent and rapid annotation of ribosomal RNA genes. *Nucleic Acids Res* 35:3100–3108. <http://dx.doi.org/10.1093/nar/gkm160>.
10. Croft MT, Warren MJ, Smith AG. 2006. Algae need their vitamins. *Eukaryot Cell* 5:1175–1183. <http://dx.doi.org/10.1128/EC.00097-06>.