

Technical University of Denmark



Multivariate decoding of mutually interacting brains reveals complementary neural mechanisms in leaders and followers

Konvalinka, Ivana

Publication date:
2014

Document Version
Publisher's PDF, also known as Version of record

[Link back to DTU Orbit](#)

Citation (APA):
Konvalinka, I. (2014). Multivariate decoding of mutually interacting brains reveals complementary neural mechanisms in leaders and followers. Abstract from FENS Brain Conference, The Social Brain, 2014, Copenhagen, Denmark.

DTU Library

Technical Information Center of Denmark

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Multivariate decoding of mutually interacting brains reveals complementary neural mechanisms in leaders and followers

Neural mechanisms underlying real-time social interaction remain largely unknown, as it has been difficult to implement paradigms in a methodologically feasible and conceptually relevant way. Researchers have increasingly begun to implement two-person paradigms by simultaneously recording brain activity from both people. Most of these studies have investigated symmetrical, and synchronized, neural mechanisms of two-people in a variety of interactive tasks. However, given that people often take on complementary, rather than symmetrical, roles during interaction, there is a real need to develop methods for quantifying complementary two-brain mechanisms. To investigate this, we implemented a multivariate decoding method to reveal complementary patterns of individual, rather than coupled, brain mechanisms in a dyad. We employed a synchronized finger-tapping task while measuring dual-EEG from pairs of participants who either tapped together in a bidirectional interactive task, or followed a computer metronome. Interactive versus non-interactive trials were classified using a multivariate analysis on both brains. This analysis revealed asymmetric patterns of frontal alpha-suppression in each pair, during task anticipation and execution, such that only one member showed the frontal component. Behavioural data analysis showed that this distinction coincided with the leader-follower relationship in 8/9 pairs, with the leaders characterized with a stronger frontal alpha-suppression during the interactive condition. This suggests that leaders invest more resources in planning and control, and shows that leader-follower relationships can be predicted from EEG recordings. Moreover, it provides a novel method for quantifying complementary neural mechanism during real-time social interaction.