Technical University of Denmark



Programmed cell death: The life ambition of the barley aleurone layer

Mark, Christina; Zor, Kinga; Heiskanen, Arto; Dufva, Martin; Emnéus, Jenny; Finnie, Christine

Publication date: 2014

Document Version Peer reviewed version

Link back to DTU Orbit

Citation (APA):

Mark, C., Zor, K., Heiskanen, A., Dufva, M., Emnéus, J., & Finnie, C. (2014). Programmed cell death: The life ambition of the barley aleurone layer. Abstract from 1st INPPO World Congress on Plant Proteomics, Hamburg, Germany.

DTU Library Technical Information Center of Denmark

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

• Users may download and print one copy of any publication from the public portal for the purpose of private study or research.

- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Programmed cell death: The life ambition of the barley aleurone layer

<u>Christina Mark¹</u>, Kinga Zór², Arto Heiskanen², Martin Dufva³, Jenny Emnéus² and Christine Finnie¹

¹Agricultural and Environmental Proteomics, Department of Systems Biology, Technical University of Denmark, ²Bioanalytics, Department of Micro- and Nanotechnology, Technical University of Denmark, ³Fluidic Array Systems and Technology, Department of Micro- and Nanotechnology, Technical University of Denmark

We have developed a 24-well multiplate tissue culture system with electrochemical and optical detection techniques for cultivation of immobilised barley aleurone layers. We have applied the system for the purpose of studying the underlying mechanisms of programmed cell death (PCD) in plants.

We have optimised an electrochemical, intracellular, whole-cell redox activity assay [1] that probes the NAD(P):NAD(P)H ratio via a double-mediator system. Experiments show that redox activity changes depend on phytohormone activation or inactivation of aleurone layer metabolism and subsequent PCD.

We have successfully applied a fluorescent double-probe system [2] to detect PCD to ensure that our redox activity data match with known responses of barley aleurone layers to phytohormones.

We have also used the system for transformation of barley aleurone cells with α -amylase-GFP constructs for the purpose of studying the timing of α -amylase production in relation to PCD. These studies will be combined with activity assays and quantitative proteomics studies of α -amylase and other target enzymes.

- [1] Heiskanen et al., 2009, Anal Biochem, 384, 11-19
- [2] Fath et al., 2001, Plant Physiol, 126, 156-166

Keywords:

Barley aleurone layer, redox activity, programmed cell death, plant tissue culture

Remark / Message to the Program Committee and Chairs:

In case this abstract should be chosen for oral presentation: Quantitative proteomics studies are planned for August/September and can therefore not be guaranteed performed and analysed for the Congress. Despite this, I would be happy to give a presentation of the work already performed.