

Bioenergy production from agri-industrial biomass residues a consequential LCA

Tonini, Davide; Hamelin, Lorie; Astrup, Thomas Fruergaard

Publication date:
2014

Document Version
Publisher's PDF, also known as Version of record

[Link back to DTU Orbit](#)

Citation (APA):

Tonini, D., Hamelin, L., & Astrup, T. F. (2014). Bioenergy production from agri-industrial biomass residues: a consequential LCA. Poster session presented at 9th International Conference on Life Cycle Assessment in the Agri-food Sector, San Francisco, United States.

DTU Library Technical Information Center of Denmark

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.


Bioenergy production from agri-industrial biomass residues: a consequential LCA



Davide Tonini, Lorie Hamelin and Thomas Astrup
dait@env.dtu.dk


Intro

Biofuels from residues (of industrial and agricultural production) promise sustainable bioenergy and greenhouse gases mitigation. However, many studies tend to forget that these biomasses are today used for specific purposes (e.g., feeding). Thus, their use for energy may trigger an increase in the international demand of feed products that may finally induce an expansion of cropland into other ecosystems (and/or an intensification). Failing to account for these consequences may lead to results that misrepresent the actual environmental impacts.





Materials and method

Tools	Consequential LCA with biochemical model (integrated)
Functional Unit	1 metric tonne of biomass (dry basis)
Geographic & time scope	Europe EU27 (time scope: 2015-2030)
Assessment method	ILCD-recommended: global warming (GW), acidification (AC), aquatic eutrophication (AE)
Biomasses investigated	Whey, brewer's grain, wheat straw, nature grass, beet molasses, beet top, beet pulp, potato pulp
Scenarios	I) Bioethanol production; molasses-for-biogas (CHP) II) Bioethanol production; molasses-for-feed III) Biogas-for-transport (PSA upgrading) IV) Biogas-for-CHP All scenarios involve use of residual solids for CHP



Marginal regions
Responding to an increased demand by increasing production and export (future)

Maize 
Expansion: 0.57 kg CO₂-eq. kg⁻¹
Intensification: 0.18 kg CO₂-eq. kg⁻¹

Soybean 
Expansion: 2.9 kg CO₂-eq. kg⁻¹
Intensification: 0.01 kg CO₂-eq. kg⁻¹

Share of intensification and expansion for each region (future)

Increased fertilizers use due to intensification (future)


Carbon losses from ecosystems conversion

iLUC model

Deterministic approach based on simplified modeling of future agricultural markets

Conclusion

General

Residues with high nutritional value should be preferably used for feed 

iLUC is the most important contributor to the induced impacts 

Best biomasses

Straw and grass promise the highest environmental savings (no competition with feed involved) 

Best scenario overall

Production of biogas (for CHP) because of higher efficiencies

Best scenario for transport fuel

Bioethanol appears better than biogas (considering PSA upgrading)

