Technical University of Denmark



Magnetic High-Field Phases of Magnetoelectric LiNiPO4

Fogh, Ellen; Toft-Petersen, R.; Nojiri, H.; Granroth, G.; Kihara, T.; Christensen, Niels Bech

Publication date: 2014

Document Version Publisher's PDF, also known as Version of record

Link back to DTU Orbit

Citation (APA):

Fogh, E., Toft-Petersen, R., Nojiri, H., Granroth, G., Kihara, T., & Christensen, N. B. (2014). Magnetic High-Field Phases of Magnetoelectric LiNiPO4. Abstract from Neutron Scattering in Magnetic Fields Above 15 Tesla, Berlin, Germany.

DTU Library Technical Information Center of Denmark

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

• Users may download and print one copy of any publication from the public portal for the purpose of private study or research.

- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Magnetic High-Field Phases of Magnetoelectric LiNiPO₄

E. Fogh ¹, *R.* Toft-Petersen ², *H.* Nojiri ³, *G.* Granroth ⁴, *T.* Kihara ³ and *N. B.* Christensen ¹

¹ Department of Physics, Technical University of Denmark, Kongens Lyngby, Denmark
 ² Helmholtz Zentrum Berlin f
ür Materialen und Energie, Berlin-Wannsee, Germany
 ³ Institute for Material Research, Tohoku University, Sendai, Japan
 ⁴ Neutron Scattering Sciences Division, Oak Ridge National Laboratory, Oak Ridge,

Tennesee

In this work the magnetic high-field phases and magnetoelectric properties of LiNiPO_4 are studied up to 30T.

LiNiPO₄ is a member of the family of lithium orthophosphates which are of orthorhombic structure and exhibit the magnetoelectric effect [1]. LiNiPO₄ is a quasi-2D system with long range antiferromagnetic order below T_N =21.8T in zero-field. The magnetic phase diagram of the material has previously been studied up to 17.3T [2,3] and pulsed-field magnetization measurements up to 23T show several additional phase transitions [4].

Here LiNiPO₄ is investigated for fields up to 30T by pulsed high magnetic field time-of-flight Laue neutron diffraction and by magnetization and electric polarization measurements. Two high-field phases were found in agreement with the previous observations. The ordering vectors were determined and possible spin structures were suggested. The polarization measurements show a reentrance of the magnetoelectric effect for μ_0 H≈19−21T, which points towards a commensurate ferrimagnetic state. However, the behavior of the effect is not fully understood. The effect vanishes above 21T where either a linearly polarized spin structure or a spin spiral is present.

The experimental technique using pulsed high magnetic fields in connection with neutrons is relatively new [5]. Nevertheless, the results displayed here show that the technique presents a future possibility to study high-field phases of certain magnetic materials.

References

[1] M. Mercier, J. Gareyt, and E. F. Bertaut, C. R. Acad. Sci. Ser. B 264, 979, 1967
[2] T. B. S. Jensen, N. B. Christensen, M. Kenzelmann, H. M. Rønnow, C. Niedermayer, N. H. Andersen, K. Lefmann, J. Schefer, M. Zimmermann, J. Li, J. L. Zarestky, and D. Vaknin, Phys. Rev. B, 79(092412), 2009
[3] R. Toft-Petersen, J. Jensen, T. B. S. Jensen, N. H. Andersen, N. B. Christensen, C. Niedermayer, M. Kenzelmann, M. Skoulatos, M. D. Le, K. Lefmann, S. R. Hansen, J. Li, J. L. Zarestky, and D. Vaknin, Phys. Rev. B, 84(054408), 2011
[4] V.M. Khrustalyov, V.N. Savitsky, and N.F. Kharchenko, Czechoslovak Journal of Physics, 54, 2004

[5] S. Yoshii, K. Ohoyama, K. Kurosawa, H. Nojiri, M. Matsuda, P. Frings, F. Duc, B. Vignolle, G. L. J. A. Rikken, L.-P. Regnault, S. Michimura, and F. Iga, Phys. Rev. Letters, 103(077203), 2009