

Technical University of Denmark



NuSTAR Reveals Relativistic Reflection but no Ultra-fast Outflow in the Quasar PG1211+143

Zoghbi, A.; Miller, J. M.; Walton, D. J.; Harrison, F. A.; Fabian, A. C.; Reynolds, C. S.; Boggs, S. E.; Christensen, Finn Erland; Craig, W.; Hailey, C. J.; Stern, D.; Zhang, W. W.

Published in:

The Astrophysical Journal Letters

Link to article, DOI:

[10.1088/2041-8205/799/2/L24](https://doi.org/10.1088/2041-8205/799/2/L24)

Publication date:

2015

Document Version

Peer reviewed version

[Link back to DTU Orbit](#)

Citation (APA):

Zoghbi, A., Miller, J. M., Walton, D. J., Harrison, F. A., Fabian, A. C., Reynolds, C. S., ... Zhang, W. W. (2015). NuSTAR Reveals Relativistic Reflection but no Ultra-fast Outflow in the Quasar PG1211+143. *The Astrophysical Journal Letters*, 799(2), [L24]. DOI: 10.1088/2041-8205/799/2/L24

DTU Library

Technical Information Center of Denmark

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

NuSTAR REVEALS RELATIVISTIC REFLECTION BUT NO ULTRA-FAST OUTFLOW IN THE QUASAR PG 1211+143

A. ZOGHBI¹, J. M. MILLER¹, D. J. WALTON^{2,3}, F. A. HARRISON³, A. C. FABIAN⁴, C. S. REYNOLDS⁵, S. E. BOGGS⁶, F. E. CHRISTENSEN⁷, W. CRAIG⁶, C. J. HAILEY⁸, D. STERN², W. W. ZHANG⁹

Draft version January 9, 2015

ABSTRACT

We report on four epochs of observations of the quasar PG 1211+143 using *NuSTAR*. The net exposure time is 300 ks. Prior work on this source found suggestive evidence of an “ultra-fast outflow” (or, UFO) in the Fe K band, with a velocity of approximately $0.1c$. The putative flow would carry away a high mass flux and kinetic power, with broad implications for feedback and black hole-galaxy co-evolution. *NuSTAR* detects PG 1211+143 out to 30 keV, meaning that the continuum is well-defined both through and above the Fe K band. A characteristic relativistic disk reflection spectrum is clearly revealed, via a broad Fe K emission line and Compton back-scattering curvature. The data offer only weak constraints on the spin of the black hole. A careful search for UFO’s show no significant absorption feature above 90% confidence. The limits are particularly tight when relativistic reflection is included. We discuss the statistics and the implications of these results in terms of connections between accretion onto quasars, Seyferts, and stellar-mass black holes, and feedback into their host environments.

Subject headings: galaxies: active – accretion disks – black hole physics – X-rays: binaries – galaxies: individual (PG 1211+143)

1. INTRODUCTION

The observation of a relation between the masses of supermassive black holes at the centers of galaxies and the stellar velocity dispersion ($M - \sigma$ relation; Ferrarese and Merritt 2000; Gültekin et al. 2009) suggests a direct link between black holes and their host galaxies. Energy and momentum driven out from the central regions push gas and dust away, halting star formation and stopping AGN fueling (Silk and Rees 1998; Di Matteo et al. 2005; Churazov et al. 2005; see Fabian 2012 for a review). The action of AGN feedback could be achieved through the powerful radio jets in the kinetic mode (e.g. McNamara and Nulsen 2007). In the radiative mode, accretion disks drive powerful winds that could contribute significantly to the energy budget of the BH-galaxy system.

Observing the properties of such a wind is of great importance, particularly in X-rays where most of the radiation from the expelled material is produced. Although warm absorber winds are common in the X-ray spectra of AGN (Reynolds 1997; Crenshaw et al. 2003; Blustin

et al. 2005), with outflow velocities of $\sim 1000 \text{ km s}^{-1}$ and column densities of $\log(N_{\text{h}}) \sim 20 - 23 \text{ cm}^{-2}$, they are weak, providing only $\sim 0.01\%$ of the AGN bolometric luminosity (Blustin et al. 2005). The more powerful winds seen in several objects with outflow velocities of $\sim 0.1c$ and column densities of $\log(N_{\text{h}}) = 24 \text{ cm}^{-2}$ could carry power that is a few percent of the bolometric luminosity (Pounds et al. 2003; Pounds and Reeves 2009; Reeves et al. 2003; Cappi et al. 2009; Chartas et al. 2009; Tombesi et al. 2010, 2012; Gofford et al. 2013).

These ultra-fast outflows (UFO) seem to be present in at least 35% of observed AGN in X-rays (Tombesi et al. 2010). However, this number could be an overestimate when alternative modeling and more conservative statistical analyses are considered (Kaspi and Behar 2006; Vaughan and Uttley 2008). Establishing how common these outflows are, their physical and geometrical properties is therefore crucial to understanding their contribution to the energy and momentum budget of black holes and their hosts. In this letter, we present analysis of the *NuSTAR* (Harrison et al. 2013) observation of the quasar PG 1211+143. *NuSTAR* band (3 – 79 keV) with the unprecedented sensitivity at hard ($> 10 \text{ keV}$) X-rays, fixes the continuum and thus allows a meaningful search for blue-shifted absorption below 10 keV.

PG 1211+143 ($z = 0.0809$) is the archetypal case for the ultra-fast outflows in active galaxies. The first observation with *XMM-Newton* in 2001 showed evidence for highly blue-shifted absorption lines that are reminiscent of mildly relativistic disk winds ($\sim 0.1c$; Pounds et al. 2003). The same dataset was analyzed by Kaspi and Behar (2006) who find a best fit outflow velocity of 3000 km s^{-1} instead of the high $24,000 \text{ km s}^{-1}$. A *Chandra* LETG observation showed two redshifted (instead of blueshifted!) absorption lines at 4.56 and 5.33 keV in the source frame (Reeves et al. 2005), which, when identified

abzoghbi@umich.edu

¹Department of Astronomy, University of Michigan, 1085 South University Avenue, Ann Arbor, MI 48109, USA

²Jet Propulsion Laboratory, California Institute of Technology, Pasadena, CA 91109, USA

³Space Radiation Laboratory, California Institute of Technology, Pasadena, CA 91125, USA

⁴Institute of Astronomy, University of Cambridge, Madingley Road, Cambridge CB3 0HA, UK

⁵Department of Astronomy, University of Maryland, College Park, MD 20742-2421, USA

⁶Space Science Laboratory, University of California, Berkeley, California 94720, USA

⁷DTU Space. National Space Institute, Technical University of Denmark, Elektrovej 327, 2800 Lyngby, Denmark

⁸Columbia Astrophysics Laboratory, Columbia University, New York, New York 10027, USA

⁹NASA Goddard Space Flight Center, Greenbelt, Maryland 20771, USA




Figure 1. *Left:* 3–30 keV light curves from all the four *NuSTAR* observations of PG 1211+143. The net exposure is shown in each panel. **Right:** The spectrum from the combined observations (combining FPMA and FPMB from all observations) unfolded to a constant to remove the effective area of the detector, and plotted as $EF(E)$ using `plot_eufspec` in XSPEC.

as the H-like $K\alpha$ lines corresponds to inflowing velocities of $(0.2 - 0.4)c$. Later *XMM-Newton* observations in 2004 and 2007 showed weaker lines but seem to be consistent with the original observations (Pounds and Reeves 2009), or possibly with no absorption lines at all (Tombesi et al. 2010)

2. OBSERVATIONS & DATA REDUCTION

NuSTAR observed PG 1211+143 in four exposures between February and July 2014 (The exact dates are: 18 February 2014, 08 and 09 April 2014 and 07 July 2014). The four observations had net exposures 111, 48, 64 and 74 ks, totaling to nearly 300 ks. The data were reduced using HEASOFT v6.16 with the latest calibration (version 20141020). We used the scripts `nupipeline` and `nuproducts` to extract the spectral products. Source and background spectra were extracted from regions on (with a radius of 2 arcmin) and off source respectively and grouped so that there are least 100 source counts per bin. The spectra were analyzed using XSPEC v12.8.2. Spectral analysis was performed on individual and combined spectra as discussed in section 3. Spectra from the two focal point modules A and B (FPMA and FPMB) and from different epochs were combined using `addspec` tool in HEASOFT. The response files were combined using `addrmf` with the proper weighting.

The resulting 3–30 keV light curves from the four exposures are shown in Fig. 1-left. The average 3–10 keV flux (from a power-law fit) is 2.7×10^{-12} ergs cm^{-2} s^{-1} which is about the same as the first *XMM-Newton* observation of 2001, which had a 3–10 keV flux of 2.5×10^{-12} ergs cm^{-2} s^{-1} . The source showed some flux variability between observations. No strong spectral changes are seen apart from a normalization change in the main power-law continuum (see section 3).

3. SPECTRAL ANALYSIS

One of the goals of the *NuSTAR* observation was to search for absorption lines from high velocity outflows. The spectrum from the new datasets is shown in Fig. 1-right. It has a clear iron K emission line and an excess above 10 keV that is most likely due to the Compton reflection hump. To be systematic in the search, we consider several baseline models, including simple fits to the

3–10 keV band so we can directly compare with the baseline model used in Tombesi et al. (2010).

3.1. Searching for Emission and Absorption Features

In the following discussions, we search for absorption (and emission) features by adding a narrow¹⁰ Gaussian line and doing a systematic scan of the residuals for each baseline model. We use a grid of energies between 3.5 and 9.5 keV in 100 eV steps, and normalization values between -1×10^{-5} and 1×10^5 (to account for both emission and absorption). We use a Monte Carlo method to obtain significance estimates. As pointed out in Prottasov et al. (2002), a simple F -test done by comparing the improvement in χ^2 after adding a Gaussian line is *not* appropriate for obtaining meaningful statistical confidence intervals. The baseline model is not known a priori, and neither is the reference (null) distribution of the F -statistic.

This inapplicability of the F -test is not related to the quality of data, but rather it is fundamental to any case of an added spectral component. Using Monte Carlo methods provides a way of *estimating* the unknown reference distribution and then using it to assess the significance of deviations from the null model. The following steps are used to obtain a reference distribution that takes into account both the uncertainty in the baseline model parameters. These are similar to those in Porquet et al. (2004) and Tombesi et al. (2010) but additionally take into account the baseline model uncertainties:

- A) Start with a **baseline** model with its best fit χ_0^2 and covariance matrix that encodes the uncertainty of best value parameters.
- B) Draw N random sets of parameters using the covariance matrix ($N = 1000$ in our case). This can be achieved for instance by running a Markov Chain Monte Carlo (MCMC) parameter search and taking parameters directly from the resulting chains (after they have converged).
- C) For each parameter set, fake spectra using the observed response and background files. This produces N simulated spectra that are drawn from the baseline model

¹⁰ If a broad component is present in the residuals, several neighboring narrow lines will provide a χ^2 improvements.




Figure 2. A line search significance plot for five baseline models. The regions with darkest colors indicate that the null model (baseline without an additional emission or absorption feature) can be rejected with that significance. The color scale corresponds to $\log(1 - \text{significance})$ and the corresponding significance values are shown in the color bar. The energy axis corresponds to the source rest frame energy ($z = 0.0809$).

taking into account its uncertainty.

D) Add a narrow Gaussian line and scan in energy and normalization and record the maximum improvement in $\Delta\chi^2$. This gives N values of $\Delta\chi^2$ that are used to construct a reference distribution to which observed $\Delta\chi^2$ values are compared. The significance of an observed $\Delta\chi^2_{obs}$ is $1 - N_{(\Delta\chi^2 > \Delta\chi^2_{obs})}/N$, where $N_{(\Delta\chi^2 > \Delta\chi^2_{obs})}$ is the number of simulated $\Delta\chi^2$ values greater than $\Delta\chi^2_{obs}$.

Several baseline models are used to assess their effect on estimating absorption features:

1. **powerlaw + zgauss.** This consists of a power-law plus a broad Gaussian fitted to the 3–10 keV spectrum similar to those used in Tombesi et al. (2010) and Gofford et al. (2013).
2. **powerlaw+reflionx** (Ross and Fabian 2005).
3. **powerlaw+xillver** (García et al. 2014).
4. **relxill** (Dauser et al. 2013).

For models in 2,3 and 4, the whole *NuSTAR* band where the source is significantly detected above background is used. In all models, Galactic absorption is fixed at $2.7 \times 10^{20} \text{ cm}^{-2}$. For the relativistic reflection models, we assume a power-law emissivity and we fit for its index (q) along with the black hole spin, photon index, ionization parameter, iron abundance, inclination, high energy cutoff and reflection fraction. The details of the model parameters are shown in Table 1. For each baseline model, we repeat steps A-D. The results are shown in Fig. 2.

3.2. Line Search Results

It is clear from Fig. 2 that there is *no* significant absorption feature that is persistent with all baseline models. The phenomenological description of the first model gives an excellent description to the data in the 3 – 10 keV band with no significant residuals. In the **reflionx** fit, an absorption feature at 7.1 keV is apparent along with the emission feature at 5.8 keV. The same feature at 5.8 keV persists for the **xillver** fit. Both the **reflionx** and **xillver** models account for the Fe $K\alpha$ emission

$E_{z\text{gauss}}$	powerlaw+zgauss (3–10 keV)
$\sigma_{z\text{gauss}}$	$6.28 \pm 0.07 \text{ keV}$
$A_{\text{Fe}} \text{ (solar)}$	powerlaw+reflionx (3–50 keV)
Γ	0.7 ± 0.1
$\log(\xi)$	2.45 ± 0.06
$A_{\text{Fe}} \text{ (solar)}$	powerlaw+xillver (3–50 keV)
Γ	0.5 ± 0.1
$\log(\xi)$	2.50 ± 0.06
q	relxill (3–50 keV)
a	$2.2 \pm .5$
$incl.$	-0.13 ± 0.8
Γ	28 ± 7
$\log(\xi)$	2.51 ± 0.2
$A_{\text{Fe}} \text{ (solar)}$	1.3 ± 0.9
E_{cut}	0.7 ± 0.1
Ref. fract.	$124 \text{ (keV) (lower limit)}$
	$2.9^{+1.6}_{-0.4}$

Table 1

Fit parameters for the baseline models discussed in section 3.1.

at 6.4 keV. The residuals at ~ 5.8 indicate that additional emission is required red-ward of the line, suggesting line broadening. If broadening is included in the fit (**relxill**, **relxill+xillver**), the feature disappears.

The absorption in the **reflionx** fit at 7.1 keV disappears too when relativistic blurring is included. The last model (**relxill+xillver**) is included for completeness. It models reflection from both the inner (with relativistic broadening) and outer regions. Although, the additional component improves the χ^2 slightly compared to **relxill** alone, its requirement is not very significant. The conclusion from Fig. 2 is that absorption features depend on the chosen continuum, and when the continuum is pinned down with *NuSTAR* coverage at high energies and the inclusion of relativistic reflection, no significant absorption is seen.

The analysis summarized in Fig. 2 was done using a combination of all the spectra (FPMA and FPMB from four observations) to obtain the best signal. We tested for any possible flux dependency by grouping observa-

tions 1 and 4 together and 2 and 3 together (see light curve in Fig. 1-left), and the results do not change; no additional absorption lines are required by the data.

3.3. Comparison with *XMM-Newton*

The first *XMM-Newton* observation in 2001 showed an absorption line at ~ 7.1 keV in the observer’s frame plus possible additional lines at lower energies (Pounds et al. 2003). We reanalyzed this *XMM-Newton* dataset using the same method discussed in section 3.1 and find the line at ~ 7.5 keV to be significant at the $\sim 99\%$ level with an equivalent width of 104 ± 52 eV. The flux level of the first *XMM-Newton* observation and the *NuSTAR* observation discussed here are comparable; therefore, we also assess whether an absorption line similar to that seen in *XMM-Newton* would have been detectable in the current *NuSTAR* data. We fitted the *XMM-Newton* PN spectrum with a model consisting of a power-law, relativistic reflection and an absorption line. We then used this model to fake *NuSTAR* data, and followed the procedure of section 3.1 to search for absorption line. The line is detected significantly in this simulated data showing that an absorption line would have been easily ($> 8\sigma$ significance) detected if it was present in the data. Adding a Gaussian line to our best fit model (`relxill`) with the energy and the width of absorption line seen in the *XMM-Newton* spectrum provides a fit improvement of $\Delta\chi^2 = 3$ for one degree of freedom. The equivalent width of this line is 22 ± 22 eV.

Subsequent to the first observation, *XMM-Newton* observed PG 1211+143 several times, and when we analyze these observations self-consistently, none of them show any significant absorption features when using a baseline model consisting of a power-law and a relativistic line.

3.4. Relativistic Reflection Parameters

The best fit model to the *NuSTAR* data, shown in Fig. 3, consists of an absorbed power-law and relativistic reflection, both modeled with `relxill`. Only data below 10 keV are shown to highlight the differences among models. As was clear from Fig. 2, the data show an excess at ~ 5 keV when fitted with simple reflection models. The addition of relativistic blurring (`relxill`) accounts for the excess. The data, however, do not allow very strong constraints on the spin of the black hole. The parameters of the best fit model are shown in Table 1.

If we fit for the inner radius of emission instead of the black hole spin, we obtain the marginal probability contours shown in the bottom panel of Fig. 3. This figure is obtained by running a Monte Carlo Markov Chain¹¹ on the best fit `relxill` to obtain posterior probability distributions of the fit parameters and then marginalize over the parameters that are not of interest. The inner radius of emission and emissivity index are degenerate in the fit, but it is clear that although the extra broadening of the Fe $K\alpha$ line is required, it is not very large. The inner radius of emission ranges from ~ 4 to 40 gravitational radii ($r_g = GM/c^2$) depending on the disk emissivity. The solution with a steep emissivity and large radii might be unphysical, and although this data do not allow us to




Figure 3. *Top:* The spectrum of PG 1211+143 along with the best fit baseline models discussed in section 3.1 plotted in the 3–10 keV only where the models differ. The x-axis represents the rest-frame energy of the source. *Bottom:* Marginal probability density contours in the `relxill` model for the inner radius of emission and the emissivity index. The contours are for the 68, 90 and 99% confidence levels.

firmly locate the X-ray emission, the detection of reverberation time delays in this objects in the *XMM-Newton* data (de Marco et al. 2011) would favor the solution with a small radius for the emission.

4. DISCUSSION AND CONCLUSIONS

We have analyzed a deep 300 ks observation of the quasar PG 1211+143, obtained with *NuSTAR* in four time intervals spanning six months. When fit with a simple power-law model, the total spectrum shows clear signatures of disk reflection (Fig. 1-right). Fits with the newest, most sophisticated reflection models currently available provide significantly improved fits. The residuals to fits with simple reflection show an excess around ~ 5 keV indicating that relativistic blurring is required, clearly signaling that X-ray emission probes the very innermost portion of the accretion flow in PG 1211+143. The data do not permit a strong constraint on the spin of the black hole, but when reverberation delays seen previously in *XMM-Newton* data are considered, a solution with a small inner radius ($\sim 4r_g$, corresponding to a black hole spin of ~ 0.5) is supported by the data. We thoroughly examined the *NuSTAR* data both in segments and in total, and the spectra show no compelling evi-

¹¹ We use `xspec_emcee` available from: https://github.com/jeremysanders/xspec_emcee

dence of a steady or variable ultra-fast outflow seen previously in this source. The limits are particularly tight once disk reflection has been modeled.

NuSTAR clearly has the sensitivity and spectral resolution required to detect extraordinary outflows, when they are present. Observations of PDS 456, for instance, reveal a likely P-Cygni profile in that quasar, signaling a massive and powerful outflow from that black hole into its host galaxy (Nardini et al. 2014, submitted). *NuSTAR* spectra from observations of NGC 1365 were able to disentangle variable absorption, outflows, and disk reflection in that source, leading to a robust measure of the black hole spin (Risaliti et al. 2013; Walton et al. 2014). At the opposite end of the mass scale, *NuSTAR* observations of the black hole candidate 4U 1630–472 revealed strong absorption that may be a UFO-like outflow, in addition to disk reflection (King et al. 2014). The true extent of these ultra-fast outflows and their persistence is unclear.

The non-detection of ultra-fast outflows in the prototypical object PG 1211+143 in the current observation (and also previous observations subsequent to the first *XMM-Newton* detection) raises several questions on how common these might be in this and other sources. Systematic searches suggest that $\sim 40\%$ of objects observed with *XMM-Newton* and *Suzaku* show them (Tombesi et al. 2010; Gofford et al. 2013). This number could be much lower if a careful statistical search is performed that takes into account physical baseline models along with their observational uncertainties. This comes in the form of including the best fit uncertainties in simulating spectra during the Monte Carlo tests and also allowing for that uncertainty when fitting for narrow features at arbitrary energies in the spectrum.

The true extent of UFO's could be affected by the significance of any single detection, which we have discussed, and also by their variability. A non-detection could simply be due to the fact they are transient in nature as suggested by the the simulations (Proga and Kallman 2004). The case of PG 1211+143 shows that the UFO is seen in one out of eleven observations (*XMM-Newton*, *Suzaku*, *Chandra* and *NuSTAR*) between 2001 and 2014. With the caveat that we still have small numbers, if one assumes the observations are done randomly as far as the AGN is concerned, it appears that a variability argument would also suggest that the contribution of UFO's is an order of magnitude lower than previously inferred.

ACKNOWLEDGMENT

This work made use of data from the *NuSTAR* mission, a project led by the California Institute of Technology, managed by the Jet Propulsion Laboratory, and funded by the National Aeronautics and Space Administration.

REFERENCES

- Blustin, A. J., Page, M. J., Fuerst, S. V., Branduardi-Raymont, G., and Ashton, C. E.: 2005, *A&A* **431**, 111
- Cappi, M., Tombesi, F., Bianchi, S., et al.: 2009, *A&A* **504**, 401
- Chartas, G., Saez, C., Brandt, W. N., Giustini, M., and Garmire, G. P.: 2009, *ApJ* **706**, 644
- Churazov, E., Sazonov, S., Sunyaev, R., et al.: 2005, *MNRAS* **363**, L91
- Crenshaw, D. M., Kraemer, S. B., and George, I. M.: 2003, *ARA&A* **41**, 117
- Dauser, T., Garcia, J., Wilms, J., et al.: 2013, *MNRAS* **430**, 1694
- de Marco, B., Ponti, G., Uttley, P., et al.: 2011, *MNRAS* **417**, L98
- Di Matteo, T., Springel, V., and Hernquist, L.: 2005, *Nature* **433**, 604
- Fabian, A. C.: 2012, *ARA&A* **50**, 455
- Ferrarese, L. and Merritt, D.: 2000, *ApJ* **539**, L9
- García, J., Dauser, T., Lohfink, A., et al.: 2014, *ApJ* **782**, 76
- Gofford, J., Reeves, J. N., Tombesi, F., et al.: 2013, *MNRAS* **430**, 60
- Gültekin, K., Richstone, D. O., Gebhardt, K., et al.: 2009, *ApJ* **698**, 198
- Harrison, F. A., Craig, W. W., Christensen, F. E., et al. *ApJ* **770**, 103
- Kaspi, S. and Behar, E.: 2006, *ApJ* **636**, 674
- King, A. L., Walton, D. J., Miller, J. M., et al.: 2014, *ApJ* **784**, L2
- McNamara, B. R. and Nulsen, P. E. J.: 2007, *ARA&A* **45**, 117
- Porquet, D., Reeves, J. N., Uttley, P., and Turner, T. J.: 2004, *A&A* **427**, 101
- Pounds, K. A. and Reeves, J. N.: 2009, *MNRAS* **397**, 249
- Pounds, K. A., Reeves, J. N., King, A. R., et al.: 2003, *MNRAS* **345**, 705
- Proga, D. and Kallman, T. R.: 2004, *ApJ* **616**, 688
- Protassov, R., van Dyk, D. A., Connors, A., Kashyap, V. L., and Siemiginowska, A.: 2002, *ApJ* **571**, 545
- Reeves, J. N., O'Brien, P. T., and Ward, M. J.: 2003, *ApJ* **593**, L65
- Reeves, J. N., Pounds, K., Uttley, P., et al.: 2005, *ApJ* **633**, L81
- Reynolds, C. S.: 1997, *MNRAS* **286**, 513
- Risaliti, G., Harrison, F. A., Madsen, K. K., et al.: 2013, *Nature* **494**, 449
- Ross, R. R. and Fabian, A. C.: 2005, *MNRAS* **358**, 211
- Silk, J. and Rees, M. J.: 1998, *A&A* **331**, L1
- Tombesi, F., Cappi, M., Reeves, J. N., and Braito, V.: 2012, *MNRAS* **422**, L1
- Tombesi, F., Cappi, M., Reeves, J. N., et al.: 2010, *A&A* **521**, A57
- Vaughan, S. and Uttley, P.: 2008, *MNRAS* **390**, 421
- Walton, D. J., Risaliti, G., Harrison, F. A., et al.: 2014, *ApJ* **788**, 76