

Ecotoxicity Testing of Nanoparticles for Remediation of Contaminated Soil and Groundwater

Hjorth, Rune; Coutris, Claire ; Nguyen, Nhung ; Sevcu, Alena ; Baun, Anders; Joner, Erik

Publication date:
2015

[Link back to DTU Orbit](#)

Citation (APA):

Hjorth, R., Coutris, C., Nguyen, N., Sevcu, A., Baun, A., & Joner, E. (2015). Ecotoxicity Testing of Nanoparticles for Remediation of Contaminated Soil and Groundwater. Poster session presented at 13th International UFZ-Deltares Conference on Sustainable Use and Management of Soil, Sediment and Water Resources, Copenhagen, Denmark.

DTU Library

Technical Information Center of Denmark

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

NanoRem is a four year, €14 million research project funded through the European Commission FP7.

Background and Objectives

As innovation in remediation technology brings new solutions to old problems, regulatory bodies will require hazard and safety data as new technologies are brought from lab-scale to real-world application.

Going beyond zero-valent iron

A range of nanomaterials are developed in NanoRem in order to extend the spectrum of treatable soil and groundwater contaminants from halogenated organics to non-halogenated substances and non-reducible metals.

Contributing to hazard assessment of nanomaterials

These nanomaterials are meant to help solve the environmental problem of contaminated soil and groundwater, not to create additional ones. Therefore, it is important that their intrinsic toxicity is assessed, as required under REACH.

Ecotoxicity tests

We have tested a range of particles developed for remediation purposes, using standard ecotoxicity tests to assess their potential hazard towards groups of organisms used for hazard identification for classification and labeling purposes.

- Bacteria *Vibrio fischeri*, 15 min luminescence, ISO 11348-3
- Algae *Pseudokirchneriella subcapitata*, 48h growth, OECD 201
- Crustacean *Daphnia magna*, 48h immobilization, OECD 202
- Earthworm *Eisenia fetida*, 48h mortality, OECD 207
- Plant *Raphanus sativus*, *L. multiflorum*, 6d root length, OECD 208

Besides the standardized protocols, we also tested the particles effect on:

- Bacteria *Escherichia coli*, 6h growth and 24h cell viability
- Algae *Chlamydomonas sp.*, 48h photosynthesis efficiency
- Oligochaete *Lumbriculus variegatus*, 96h mortality

The tested particles were dispersed as recommended by the producers and the suspensions were characterized by DLS, NTA, ORP, and pH.



Nanomaterials

Carbo-Iron

- Composite of activated carbon and zero-valent iron
- Adsorption and reduction of halogenated contaminants

Photo: Wagner/Leipzig

Fe Zeolites

- Nanoporous aluminosilicate loaded with iron Fe³⁺ catalyst
- Oxidation of small molecules, e.g. BTEX, MTBE, dichloroethane, and chloroform

Photo: E. Ormen/Bioforsk

Fe Oxides

- Pristine iron oxides, mostly goethite Fe³⁺O(OH) stabilized with humic acids
- Oxidation (catalytic effect on bioremediation) of biodegradable contaminants, such as BTEX

Photo: M. Klever/NMBU

Milled zero-valent iron

- Mechanically ground zero-valent iron
- Reduction of halogenated contaminants (same spectrum as for nZVI produced by thermal reduction)

Photo: CTM

Results and Discussion

	Algal growth OECD 201	Algal photosynthesis efficiency	Root elongation OECD 208	Earthworm survival OECD 207	Lumbriculus survival	Bacterial luminescence ISO 11348-3	E. coli viability and growth	Daphnia immobilization OECD 202
Carbo-Iron	EC ₅₀ > 100 mg/L	EC ₅₀ > 100 mg/L	EC ₅₀ > 100 mg/L	EC ₅₀ > 100 mg/L	EC ₅₀ > 100 mg/L	EC ₅₀ > 100 mg/L	EC ₅₀ > 100 mg/L	EC ₅₀ > 100 mg/L
Fe Oxides	EC ₅₀ > 100 mg/L	EC ₅₀ > 100 mg/L	EC ₅₀ > 100 mg/L	EC ₅₀ > 100 mg/L	EC ₅₀ > 100 mg/L	EC ₅₀ > 100 mg/L	EC ₅₀ > 100 mg/L	EC ₅₀ > 100 mg/L
Fe Zeolites	EC ₅₀ > 100 mg/L	EC ₅₀ > 100 mg/L	EC ₅₀ > 100 mg/L	EC ₅₀ > 100 mg/L	EC ₅₀ > 100 mg/L	EC ₅₀ > 100 mg/L	EC ₅₀ > 100 mg/L	EC ₅₀ > 100 mg/L
Milled ZVI	ND	ND	ND	EC ₅₀ < 100 mg/L	EC ₅₀ < 100 mg/L	EC ₅₀ < 100 mg/L	EC ₅₀ < 100 mg/L	ND

Standard ecotoxicity testing of nanoparticles has in general proven technically difficult and it may be questioned whether proper hazard identification of engineered nanoparticles needed for environmental risk assessment is currently feasible. Aggregation, agglomeration, sedimentation, shading, and other physical effects are known to confound the measuring principles behind the tests and these inferences were also observed for the tested particles.

This was pronounced for tests on algae, bacteria, and crustaceans and require inclusion of additional controls to ensure a correct data interpretation. E.g. a reduction in luminescence is seen as a sign of toxicity in the *V. fischeri* assay, but the turbidity of the tested particles absorb the light emitted by the bacteria before detection and is thus not a toxic effect.

Conclusion

The low toxicities found in the standard organisms do not lead to any hazard classification according to EU regulation for any of the tested particles and the results indicate that the particles, except the milled ZVI particles, can be considered non-toxic.

Nanomaterials with the lowest toxicity profile should be preferred over other materials with similar field scale efficacy and reactivity towards the target contaminants.

¹Department of Environmental Engineering, Technical University of Denmark
Miljøvej, Building 113, 2800 Kgs. Lyngby, Denmark
Correspondence to ruhj@env.dtu.dk

²Department for Soil Quality and Climate, Bioforsk
Høgskoleveien 7, 1430 Ås, Norway

³Institute for Nanomaterials, Advanced Technologies and Innovations, Technical University of Liberec
Studentská 2, 46117 Liberec, Czech Republic

