

Technical University of Denmark



Coping with anthelmintic resistance in ruminants: the potential use of chicory (*Cichorium intybus*) as an antiparasitic forage in cattle

Pena-Espinoza, Miguel Angel; Boas, Ulrik; Thamsborg, Stig M.; Enemark, Heidi

Publication date:
2015

Document Version
Publisher's PDF, also known as Version of record

[Link back to DTU Orbit](#)

Citation (APA):

Pena-Espinoza, M. A., Boas, U., Thamsborg, S. M., & Enemark, H. (2015). Coping with anthelmintic resistance in ruminants: the potential use of chicory (*Cichorium intybus*) as an antiparasitic forage in cattle. Abstract from Joint Spring Symposium, Frederiksberg, Denmark.

DTU Library

Technical Information Center of Denmark

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Coping with anthelmintic resistance in ruminants: the potential use of chicory (*Cichorium intybus*) as an antiparasitic forage in cattle

Miguel Peña-Espinoza (1-2), Ulrik Boas (1), Stig M. Thamsborg (2) and Heidi L. Enemark (1)

(1) National Veterinary Institute, Technical University of Denmark, Denmark; (2) Department of Veterinary Disease Biology, University of Copenhagen, Denmark

Studies were performed to test the anthelmintic activity of chicory against cattle nematodes and to investigate the role of sesquiterpene lactones (SL) as active compounds. In study 1, 2-4 months-old calves were allocated into a chicory (CHI, n=9) or control (CON, n=6) group. CHI and CON were stabled and fed with chicory silage (cv. Spadona) or hay, resp., ad libitum. After 2 weeks all calves were infected with 10,000 *Ostertagia ostertagi* and 65,000 *Cooperia oncophora* third-stage (L3) larvae. In study 2, 4-6 months-old calves were allocated into a chicory (cv. Spadona, CHI, n=10) or ryegrass/white clover (CON, n=10) pasture. After 1 week all calves were infected with 20,000 *O. ostertagi* L3 larvae. Fecal egg counts were calculated as number of eggs per g of dried feces (FECDM). At day 56 (study 1) and day 36 (study 2) post-infection calves were killed for worm recovery. FECDM and worm counts were analysed by ANOVA. In study 3, SL extracts were purified from leaves of chicory cv. Spadona and cv. Puna II. *O. ostertagi* adults were incubated at decreasing concentrations of SL extracts and worm motility was evaluated after 6, 24 and 48 h of incubation (37°C). SL profile in the extracts was analysed by liquid chromatography (LC). In study 1 mean FECDM were not significantly different between groups. *O. ostertagi* mean worm counts were 1599 and 3752 in CHI and CON groups, respectively ($P < 0.05$). *C. oncophora* burdens were not statistically different between groups. In study 2 FECDM was decreased in CHI by 48-65% as compared to CON ($P < 0.05$). Worm counting of study 2 is undergoing. In study 3 Spadona-SLs showed higher potency and exerted faster worm paralysis than Puna II-SLs. LC analyses revealed a different composition of SL between cultivars. In conclusion, chicory demonstrated a marked in vivo anthelmintic effect against *O. ostertagi*, but not on *C. oncophora*. Different anthelmintic potency of chicory SL can guide the identification and selection of antiparasitic cultivars.