brought to you by I CORE

## Technical University of Denmark



## Comparison of Different Strategies for Selection/Adaptation of Mixed Microbial Cultures Able to Ferment Crude Glycerol Derived from Second-Generation Biodiesel

Varrone, Cristiano; Heggeset, T. M. B.; Le, S. B.; Haugen, T.; Markussen, S.; Skiadas, Ioannis V; Gavala, Hariklia N.

Published in:

BioMed Research International

Link to article, DOI: 10.1155/2015/932934 10.1155/2017/7602495

Publication date:

2015

Document Version Publisher's PDF, also known as Version of record

Link back to DTU Orbit

Citation (APA):

Varrone, C., Heggeset, T. M. B., Le, S. B., Haugen, T., Markussen, S., V. Skiadas, I., & Gavala, H. N. (2015). Comparison of Different Strategies for Selection/Adaptation of Mixed Microbial Cultures Able to Ferment Crude Glycerol Derived from Second-Generation Biodiesel. BioMed Research International, 2015, [932934]. DOI: 10.1155/2015/932934, 10.1155/2017/7602495

## DTU Library

Technical Information Center of Denmark

#### General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- · You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

Hindawi Publishing Corporation BioMed Research International Volume 2017, Article ID 7602495, 1 page https://doi.org/10.1155/2017/7602495



## Corrigendum

# Corrigendum to "Comparison of Different Strategies for Selection/Adaptation of Mixed Microbial Cultures Able to Ferment Crude Glycerol Derived from Second-Generation Biodiesel"

C. Varrone,<sup>1,2</sup> T. M. B. Heggeset,<sup>3</sup> S. B. Le,<sup>3</sup> T. Haugen,<sup>3</sup> S. Markussen,<sup>3</sup> I. V. Skiadas,<sup>1,2</sup> and H. N. Gavala<sup>1,2</sup>

Correspondence should be addressed to C. Varrone; cristiano.varrone@gmail.com

Received 9 November 2016; Accepted 22 January 2017; Published 6 February 2017

Copyright © 2017 C. Varrone et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

In the article titled "Comparison of Different Strategies for Selection/Adaptation of Mixed Microbial Cultures Able to Ferment Crude Glycerol Derived from Second-Generation Biodiesel" [1], there is a systematic error in the yields reported (in g/g) in section "3.1.1. Activated Sludge." It is reported again in section "4. Conclusions." These yields need to be divided by a factor 10, due to a conversion mistake. This does not alter or affect the discussions and conclusions that have been reached in this paper, since the scope was to make a comparison among the different mixed cultures obtained.

#### References


[1] C. Varrone, T. M. B. Heggeset, S. B. Le et al., "Comparison of different strategies for selection/adaptation of mixed microbial cultures able to ferment crude glycerol derived from second-generation biodiesel," *BioMed Research International*, vol. 2015, Article ID 932934, 14 pages, 2015.


<sup>&</sup>lt;sup>1</sup>Department of Chemistry and Biosciences, Aalborg University, 2350 Copenhagen, Denmark


<sup>&</sup>lt;sup>2</sup>Department of Chemical and Biochemical Engineering, Technical University of Denmark, 2800 Lyngby, Denmark

<sup>&</sup>lt;sup>3</sup>Biotechnology and Nanomedicine, SINTEF Materials and Chemistry, 7465 Trondheim, Norway






















Submit your manuscripts at https://www.hindawi.com











Enzyme

Reséarch





