



## Modeling derivative properties and binary mixtures with CO<sub>2</sub> using the CPA and the quadrupolar CPA equations of state

**Bjørner, Martin Gamel; Kontogeorgis, Georgios**

*Published in:*  
Fluid Phase Equilibria

*Link to article, DOI:*  
[10.1016/j.fluid.2016.04.019](https://doi.org/10.1016/j.fluid.2016.04.019)  
[10.1016/j.fluid.2015.08.011](https://doi.org/10.1016/j.fluid.2015.08.011)

*Publication date:*  
2016

*Document Version*  
Peer reviewed version

[Link back to DTU Orbit](#)

### *Citation (APA):*

Bjørner, M. G., & Kontogeorgis, G. (2016). Modeling derivative properties and binary mixtures with CO<sub>2</sub> using the CPA and the quadrupolar CPA equations of state. *Fluid Phase Equilibria*, 408, 151-169. DOI: 10.1016/j.fluid.2016.04.019, 10.1016/j.fluid.2015.08.011

## DTU Library

Technical Information Center of Denmark

---

### General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

# Modeling derivative properties and binary mixtures with CO<sub>2</sub> using the CPA and the quadrupolar CPA equations of state

M.G. Bjørner<sup>a,\*</sup>, G.M. Kontogeorgis<sup>a</sup>

<sup>a</sup>*Center for Energy Resources Engineering (CERE), Department of Chemical and Biochemical Engineering, Technical University of Denmark, Building 229, DK-2800 Kgs. Lyngby, Denmark.*

---

## Abstract

The cubic plus association (CPA) equation of state (EoS) is extended to include quadrupolar interactions. The quadrupolar term is based on a modification of the perturbation terms by Larsen et al. (*Mol. Phys.* **1977**, *33*, 4, 987) for a hard sphere fluid with a symmetric point quadrupole moment. The new quadrupolar CPA (qCPA) can be used without introducing any additional pure compound parameters. Alternatively a single additional adjustable parameter can be employed.

To evaluate qCPA several pure compound properties are predicted. The model is furthermore evaluated for its ability to predict and correlate binary vapor-liquid equilibria (VLE) and liquid-liquid equilibria (LLE) of mixtures containing CO<sub>2</sub> and hydrocarbons, water, alcohols, or selected quadrupolar compounds.

The results indicate that most pure compound property predictions are satisfactory but similar to other CPA approaches. When binary mixtures are considered, qCPA appear to offer a systematic improvement as compared to the cases where quadrupolar interactions are ignored. This improvement is particularly pronounced when mixtures of CO<sub>2</sub> and hydrocarbons are considered, where the model is almost fully predictive. Using the same modeling approach

---

\*Corresponding author

*Email address:* mgabj@kt.dtu.dk (M.G. Bjørner)

qCPA can accurately correlate both the phase behaviour of CO<sub>2</sub> + hydrocarbon mixtures as well as mixtures of CO<sub>2</sub> + a self-associating compound.

*Keywords:* equation of state, CPA, quadrupole, CO<sub>2</sub>, acid gas, association

---

## 1. Introduction

In recent years CO<sub>2</sub> has received a significant amount of negative attention due to its status as a greenhouse gas and the fact that the amount of CO<sub>2</sub> in the atmosphere continues to rise. This is believed to be largely due to the combustion of fossil fuels. Technologies are thus needed, which can limit the emission of CO<sub>2</sub> to the atmosphere. One such potential technology is carbon capture and storage. During this process, transport of CO<sub>2</sub> rich mixtures is an important step which requires accurate knowledge of the phase behaviour, as well as other thermodynamic properties, of mixtures containing hydrocarbons, water and other fluids such as alcohols [1].

Accurate equilibrium predictions of CO<sub>2</sub> containing mixtures, however, are challenging with traditional equations of state. The reason for this may be that CO<sub>2</sub> has a large quadrupole moment. The large quadrupole moment of CO<sub>2</sub>, as well as size asymmetry, are for instance believed to be the reasons for the liquid liquid equilibrium (LLE) between CO<sub>2</sub> and heavy hydrocarbons, and the reason for the low temperature azeotrope formed between mixtures of CO<sub>2</sub> and lighter hydrocarbons. Cubic equations of state such as the Soave-Redlich-Kwong (SRK) equation of state (EoS), however, treat CO<sub>2</sub> as an inert compound. Even in a modern EoS such as the Statistical Association Fluid Theory (SAFT) only dispersive forces are usually considered for CO<sub>2</sub>. The continued use of these models may be attributed to the fact that several mixtures, such as CO<sub>2</sub> + hydrocarbons, are described quite well, when a single binary interaction parameter ( $k_{ij}$ ), is used.

To deal with polar and quadrupolar interactions a number of multipolar terms have been suggested in the literature. These terms are mainly based on a third order perturbation theory developed from statistical mechanics by Stell

and co-workers [2–5]. The perturbation theory was originally developed for pure fluids using the Stockmayer potential or the conceptually simpler hard sphere model with a central point dipole or quadrupole. Using the former potential,  
30 Gubbins and Twu [6, 7] developed directly applicable expressions for polar and quadrupolar fluid mixtures.

During the last decade a number of quadrupolar terms have been included in the SAFT framework. Inspired by Stell and co-workers [2–5] Gross [8] developed a new quadrupolar expression using the two center Lennard-Jones pair poten-  
35 tial as the reference fluid. The resulting quadrupolar contribution was added to the Perturbed-Chain SAFT (PC-SAFT) to give the Perturbed-Chain Polar SAFT (PCP-SAFT). This EoS can be used without any additional adjustable parameters.

Karakatsani et al. [9, 10] and Karakatsani and Economou [11] introduced  
40 two quadrupolar terms to the PC-SAFT framework. Both terms are based on the perturbation terms from by Larsen et al. [5], which use the pure hard sphere fluid as the reference fluid. The authors suggest two quadrupolar terms; an expression which employs the full correlation integrals from Larsen et al. [5], this term does not use any additional adjustable parameters, and a simpler  
45 version where the correlation integrals are truncated at the zeroth order term.

Later, NguyenHuynh et al. [12] extended a group contribution SAFT EoS to quadrupolar (and polar) fluid mixtures. The quadrupolar term used by the authors is based on the theory of Gubbins and Twu [6, 7]. The term is extended to chain molecules using a procedure suggested by Jog et al. [13] and Jog and  
50 Chapman [14].

It seems that improved predictions and correlations (smaller  $k_{ij}$ ) are typically obtained for binary vapor liquid equilibrium (VLE) when a quadrupolar term is coupled to SAFT or PC-SAFT. The quadrupolar models, however, have some limitations; The models are only applicable to molecules with a highly  
55 symmetric quadrupole moment, so that the quadrupole moment reduces to a single scalar value. It has furthermore been shown that false liquid-liquid splits may be predicted by the models [15]. Mixtures of several quadrupolar molecules

are challenging, and the results are often better if only one component is assumed to have a quadrupole moment [8].

60 Another more pragmatic approach to account for the quadrupole moment of  $\text{CO}_2$  is to consider  $\text{CO}_2$  to be a self-associating compound. Such procedures often work well resulting in improved predictions and correlations with small interaction parameters [16–19]. Unfortunately the improvement is obtained at the cost of additional pure component parameters and, in some cases, an extra  
65 adjustable parameter for the binary mixtures.

In this work, inspired by the recent advances within the SAFT-family, and in an effort to obtain a physically more correct and predictive model, a quadrupolar term is proposed and combined with the well-known cubic plus association (CPA) EoS (Kontogeorgis et al. [20]). The term is based on the explicit ex-  
70 pressions developed by Larsen et al. [5] for a hard sphere fluid with a point quadrupole. To simplify the expressions we truncate the correlation integrals at the zeroth order term, similar to the approach used in the truncated Perturbed-Chain Polar SAFT (tPC-PSAFT) [10, 11].

To evaluate the new quadrupolar CPA (qCPA) the pure fluid properties  
75 of  $\text{CO}_2$  are predicted both in the critical region and in the compressed liquid region. The model is furthermore applied for the prediction and correlation of binary mixtures with  $\text{CO}_2$  and n-alkanes, water, alcohols and a few quadrupolar compounds. The model is compared with two other CPA approaches namely non-associating (n.a.) CPA, where  $\text{CO}_2$  is assumed to be an inert compound,  
80 and the case when  $\text{CO}_2$  is assumed to be self-associating following the 4C association scheme (notation from Huang and Radosz [21]). Tsivintzelis et al. [18] have published similar results with the two CPA approaches for most of the binary mixtures investigated in this work.

## 2. Models

85 The CPA EoS (Kontogeorgis et al. [20]) is an engineering EoS which has been extensively studied in the literature (e.g. [22–24]). It combines the SRK

EoS with the association term from Wertheim’s theory, which is also employed in SAFT [25–27]. The SRK term accounts for the physical interactions between molecules, while the association term takes hydrogen bonding interactions into  
 90 account.

Polar and quadrupolar interactions are not explicitly taken into account in the original formulation. However, to obtain a physically more consistent model the effect of the quadrupole is explicitly included in this work by introducing a simplified quadrupolar term to CPA.

In the literature CPA is typically presented as a pressure explicit EoS, however, it is often much more convenient to express an EoS in terms of the residual Helmholtz energy,  $(A^r(T, V, \mathbf{n}))$ , since all other residual properties can be obtained as partial derivatives of the state variables  $T$ ,  $V$  and  $\mathbf{n}$ . The reduced residual Helmholtz free energy for qCPA can be expressed as the addition of the physical SRK term, Wertheim’s association term and the quadrupolar term:

$$\frac{A_{CPA}^r(T, V, \mathbf{n})}{RT} = \frac{A_{SRK}^r(T, V, \mathbf{n})}{RT} + \frac{A_{Assoc}^r(T, V, \mathbf{n})}{RT} + \frac{A_{Quad}^r(T, V, \mathbf{n})}{RT} \quad (1)$$

95 where  $R$  is the ideal gas constant. In the absence of association or quadrupolar interactions equation (1) reduces to the SRK EoS.

### 2.1. The SRK contribution

Consider a mixture of total composition  $n$ , with total volume  $V$  and temperature  $T$ . The reduced residual Helmholtz energy of the SRK EoS for  $n$  moles of a mixture is expressed as [28]:

$$\frac{A_{SRK}^r(T, V, \mathbf{n})}{RT} = -n \ln \left( 1 - \frac{B}{V} \right) - \frac{D(T)}{RTB} \ln \left( 1 + \frac{B}{V} \right) \quad (2)$$

If the conventional van der Waals one-fluid mixing rules are employed then  $D(T)$  and  $B$  are given by

$$D(T) = \sum_i n_i \sum_j n_j a_{ij}(T) \quad (3a)$$

$$B = \sum_i n_i b_{ii} \quad (3b)$$

where  $b_{ii}$  is the pure compound co-volume parameter for component  $i$ , also denoted  $b_0$ . The cross energetic parameter,  $a_{ij}$ , is calculated using the classical geometric mean rule

$$a_{ij} = \sqrt{a_{ii}(T)a_{jj}(T)}(1 - k_{ij}) \quad (4)$$

where  $k_{ij}$  is a binary interaction parameter, typically correlated to experimental data. The temperature dependent energetic parameter,  $a_{ii}(T)$ , is given as

$$a_{ii}(T) = a_0 \left(1 + c_1 \left(1 - \sqrt{T_r}\right)\right)^2 \quad (5)$$

where  $a_0$  and  $c_1$  are pure compound parameters and  $T_r$  is the reduced temperature ( $= T/T_c$ ).

## 100 2.2. Association contribution

The contribution to the reduced residual Helmholtz free energy from the association term is given by [25–28]:

$$\frac{A_{Assoc}^r(T, V, \mathbf{n})}{RT} = \sum_i n_i \sum_{A_i} (\ln X_{A_i} - \frac{1}{2}X_{A_i} + \frac{1}{2}) \quad (6)$$

where  $A_i$  indicates bonding sites on molecule  $i$  and  $X_{A_i}$  denotes the fraction of A-sites on molecule  $i$  *not* bonded to another association site. These fractions are found by solving the system of non-linear equations given by

$$X_{A_i} = \frac{1}{1 + \frac{1}{V} \sum_j n_j \sum_{B_j} X_{B_j} \Delta^{A_i B_j}} \quad (7)$$

where the association strength,  $\Delta^{A_i B_j}$ , between site A on molecule  $i$  and site B on molecule  $j$  depends on both  $T$ ,  $V$  and  $n$  according to

$$\Delta^{A_i B_j} = g(n, V) \left( \exp\left(\frac{\varepsilon^{A_i B_j}}{RT}\right) - 1 \right) b_{ij} \beta^{A_i B_j} \quad (8)$$

where  $\varepsilon^{A_i B_j}$  and  $\beta^{A_i B_j}$  are the association energy and volume between site A of molecule  $i$  and site B of molecule  $j$  respectively.  $b_{ij}$  is the cross-covolume calculated from the classical arithmetic mean rule

$$b_{ij} = \frac{b_{ii} + b_{jj}}{2} \quad (9)$$

and  $g$  is the Radial distribution function (RDF). Originally the RDF was approximated with the expression from the Carnahan-Starling (CS) EoS. Use of the CS RDF in CPA is an approximation, however, since CPA uses the van der Waals repulsive term rather than the CS hard-sphere term employed in SAFT. Kontogeorgis et al. [29] proposed a simpler expression for  $g$

$$g(n, V) = \frac{1}{1 - 1.9\eta} \quad (10)$$

Which essentially is the RDF of CPA.  $\eta(= B/4V)$  is the reduced fluid density. No mixing rules are required in the association term, but combining rules are required for  $\epsilon^{A_i B_j}$  and  $\beta^{A_i B_j}$  if more than one associating compound is present. One of the most common combining rules is the so-called CR-1, given as;

$$\epsilon^{A_i B_j} = \frac{1}{2} (\epsilon^{A_i B_i} + \epsilon^{A_j B_j}) \quad (11a)$$

$$\beta^{A_i B_j} = \sqrt{\beta^{A_i B_i} \beta^{A_j B_j}} \quad (11b)$$

The association term is dependent on the association scheme (see Eq. (7)), i.e. the number and type of association sites for the associating compound. To denote the different association schemes the notation of Huang and Radosz [21] is adopted.

### 105 2.3. Quadrupole contribution

The expression for the quadrupolar term is an adaptation of the third order perturbation theory developed by Stell and co-workers [2–5]. It uses the same base model as Karakatsani et al. [10, 11] employed in the development of their polar PC-SAFT equations of state. The reduced residual Helmholtz free energy for the expression is calculated from a Padé approximation as

$$\frac{A_{Quad}^r(T, V, \mathbf{n})}{RT} = \frac{A_{2,Quad}^r(T, V, \mathbf{n})/RT}{1 - A_{3,Quad}^r(T, V, \mathbf{n})/A_{2,Quad}^r(T, V, \mathbf{n})} \quad (12)$$

Where  $A_2^r$  and  $A_3^r$  indicate the second- and third-order perturbation terms respectively. The first-order term is effectively zero [5]. The third-order term is the sum of both a two-body and a three-body contribution, since it has been



shown that for multi-polar interactions three-body contributions cannot be ignored [4]. That is;

$$\frac{A_{3,Quad}^r(T, V, \mathbf{n})}{RT} = \frac{A_{3,2,Quad}^r(T, V, \mathbf{n})}{RT} + \frac{A_{3,3,Quad}^r(T, V, \mathbf{n})}{RT} \quad (13)$$

Larsen et al. [5] developed explicit expressions for pure symmetric hard spheres with a point quadrupole at their centre. In this work we extend these expressions to mixtures and relate them to the model parameters of CPA. The expressions are

$$\frac{A_{2,Quad}^r}{RT} = -\frac{7}{10} \frac{N_A}{V (k_b T)^2} \sum_i^{nc} n_i \sum_j^{nc} n_j \frac{Q_{ij}^4}{\sigma_{ij}^7} I_{10}^{HS} \quad (14)$$

$$\frac{A_{3,2,Quad}^r}{RT} = \frac{36}{245} \frac{N_A}{V k_b^2 T^3} \sum_i^{nc} n_i \sum_j^{nc} n_j \frac{Q_{ij}^6}{\sigma_{ij}^{12}} I_{15}^{HS} \quad (15)$$

$$\frac{A_{3,3,Quad}^r}{RT} = -\frac{1}{6400} \frac{N_A^2}{V^2 (k_b T)^3} \sum_i^{nc} n_i \sum_j^{nc} n_j \sum_k^{nc} n_k \frac{Q_{ijk}^6}{\sigma_{ij}^3 \sigma_{ik}^3 \sigma_{jk}^3} I_{TQ}^{HS} \quad (16)$$

where  $N_A$  is the Avogadro constant,  $k_b$  is Boltzmann's constant,  $Q$  is the quadrupolar moment,  $\sigma$  is the hard sphere diameter and  $I_n$  and  $I_{TQ}$  are correlation integrals with the hard sphere model as the reference fluid.

In the work of Larsen et al. [5] the correlation integrals in equations (14)-(16) were approximated by analytical density polynomials of fifth and third order respectively. To simplify the model, and in particular the volume derivatives, we assume that these correlation functions can be truncated already at the zeroth order term. The zeroth order coefficients are given analytically as [5]:

$$I_n^{HS} \approx J_{0,n} = \frac{4\pi}{n-3} \quad (17a)$$

$$I_{TQ}^{HS} \approx J_{0,TQ} = 54\pi^2 \quad (17b)$$

Thus, rather than being a function of the volume, the correlation functions are simply three constants. This approach has been employed with a measure of success in tPC-PSAFT [9–11], although an additional adjustable parameter was needed to retain a performance comparable to the non-truncated version.

No mixing rules are required in the quadrupolar term (Eq. (14)-(16)). Combining rules are, however, required if more than one quadrupolar compound

is present. If geometric-mean combining rules are employed the form of the quadrupolar term becomes similar to the quadrupolar expressions proposed by Gubbins and Twu [6, 7] as well as that suggested by Gross [8]. Thus the cross quadrupolar moment for two- and three-body contributions are calculated as

$$Q_{ij} = \sqrt{Q_{ii}Q_{jj}} \quad (18)$$

$$Q_{ijk} = \sqrt[3]{Q_{ii}Q_{jj}Q_{kk}} \quad (19)$$

Finally, it would be advantageous to relate the molecular diameter of the hard-sphere, to a co-volume parameter similar to that used in CPA. When de Villiers et al. [30] extended CPA with the dipolar theories of Gross [31] and Jog and Chapman [14] they assumed that the original definition of the co-volume in terms of the molecular hard-sphere diameter could be employed ( $b = (2/3) N_A \pi \sigma^3$ ). The quadrupolar term, however, is based on a hard sphere reference fluid, whereas CPA is based on the van der Waals repulsive term. Wong and Prausnitz [32] showed that the difference between the CS EoS and the van der Waals repulsive term can be reduced if  $b^{eff} = b_0^Q \approx b/2$ . Thus, using the effective co-volume we get

$$b_0^Q = (1/3) N_A \pi \sigma^3 \quad (20)$$

where  $b_0^Q$  is the co-volume parameter in the quadrupolar term. When Eq. (20) is employed the value of  $b_0^Q$  should, ideally, be similar to that of  $b_0$ .

115 In this work two variants of the qCPA are evaluated; In the first version it is assumed that the co-volume from CPA can be set equal to the quadrupolar hard sphere co-volume i.e.  $b_0 = b_0^Q$ . Equation (20) is then used to calculate  $\sigma$ , which is used in eq. (14)-(16). In this way no additional adjustable parameters are introduced by the quadrupolar term. In the second version  $b_0^Q$  is employed  
120 as an additional adjustable parameter. The experimental quadrupolar moment of CO<sub>2</sub> (-4.3 DÅ) is employed in both cases.

The terms described in this section are, in many aspects, similar to those employed by Karakatsani et al. [10, 11] in tPC-PSAFT. The main differences between the two equations of state are that the combining rules suggested in this

125 work (Eq. (18)-(19)) are in closer agreement with those employed by Gubbins and Twu [6, 7] and Gross [8] and that the base EoS is CPA rather than PC-SAFT.

### 3. Results and Discussion

#### 3.1. Parameter Estimation

130 CPA has three adjustable parameters for non-associating compounds ( $a_0, b_0, c_1$ ), and two additional parameters for associating compounds ( $\beta^{A_i B_i}, \epsilon^{A_i B_i}$ ). The suggested qCPA can have either the same number of parameters or an additional parameter ( $b_0^Q$ ). For non quadrupolar compounds qCPA reduces to CPA.

The pure compound parameters in CPA are typically fitted to experimental pressures and liquid densities at saturation using a weighted least squares objective function of the type

$$\min \phi = \sum_{i=1}^N \left[ \left( \frac{\rho_{exp,i}^{liq} - \rho_{calc,i}^{liq}}{\rho_{exp,i}^{liq}} \right)^2 + \left( \frac{P_{exp,i}^{sat} - P_{calc,i}^{sat}}{P_{exp,i}^{sat}} \right)^2 \right] \quad (21)$$

135 where  $N$  is the number of experiments,  $P^{sat}$  the saturated pressure and  $\rho^{liq}$  the liquid density. Subscripts indicate experimental and calculated values respectively. The CPA parameters from literature, which are employed in this work, have been estimated using this objective function. The parameters are shown in Appendix A

Often, the 'experimental data' employed for parameter estimation originate 140 from pure compound correlations such as those used by the Design Institute for Physical Properties (DIPPR) [33] or the National Institute of Standards and Technology (NIST) [34]. As the equations of state correlate the pseudo-experimental data with increasing accuracy, however, the risk of fitting parameters to any systematic bias in the correlations increases. Moreover, a common 145 issue with modern equations of state appear to be, that as the number of adjustable parameters increases several parameter sets exist, which represent the pure compound properties within experimental error [12, 13, 20, 35].

We have recently investigated the uncertainty of the pure compound parameters for CO<sub>2</sub> using several different modeling approaches including the new qCPA [36]. To estimate the uncertainty of the model parameters real experimental data are employed, rather than a correlation, and a slightly different objective function is employed, which uses the variance as the weighting factor, rather than the experimental value.

$$\min \phi = \sum_{i=1}^N \left( \frac{y_i^{exp} - y_i^{calc}}{s_i^{exp}} \right)^2 \quad (22)$$

Where  $s^{exp}$  is the standard deviation and  $y_i^{exp}$  is the experimental value of the  $i$ th saturated liquid density or saturated pressure. We demonstrate in Bjørner et al. [36] that the parameter uncertainty is substantial when qCPA is employed with four adjustable parameters. To evaluate the model it was thus decided to select two parameter sets, which are within the uncertainty; one which reproduces well the VLE of CO<sub>2</sub> + propane at a single temperature and one which represents well the LLE of CO<sub>2</sub> + n-dodecane. We stress, however, that we have not directly used any VLE or LLE data in the parameter fitting. Table 1 shows the CO<sub>2</sub> parameters and associated %AADs for the approaches which we employ in this work. Very good agreement is obtained with all approaches. It is noteworthy, however, that the deviations for CO<sub>2</sub> with qCPA are lower than CPA without association (CPA n.a.) even when the same number of adjustable parameters are employed.

Besides CO<sub>2</sub> we employ a few other quadrupolar compounds, namely water, ethane, benzene and acetylene. We assume, however, that the quadrupolar moment of water is negligible compared to the directional forces of the association term, moreover the quadrupole moment of water is a tensor and it is thus difficult to treat with current models. It is also assumed that the small quadrupolar moment of ethane can be ignored. For benzene and acetylene, however, the quadrupole moment cannot be ignored and their parameters have been re-estimated. Table 2 shows these parameters, for the three parameter version of qCPA, as well as CPA parameters for 1-nonanol and acetylene.

The value of the quadrupolar moment of CO<sub>2</sub> is relatively well-defined, rang-

Table 1: CPA and qCPA pure compound parameters for CO<sub>2</sub> together with %AADs between experimental [37–39] and calculated saturated liquid densities and saturated pressures. The quadrupolar moment of CO<sub>2</sub> is fixed at the experimental value of -4.3 DÅ<sup>a</sup> when qCPA is employed. Details on the parameter estimation can be found in [36].

Modeling approach	$b_0$ mL/mol	$\Gamma(a_0/(Rb_0))$ K	$c_1$ -	$\beta \cdot 1000$ -	$\varepsilon/R$ K	$b_0^Q$ mL/mol	$P^{sat}$	%AAD <sup>b</sup> $\rho^{liq}$
CPA, n.a. <sup>c</sup>	27.28	1550	0.77	-	-	-	0.18	0.95
CPA, 4C	28.41	1329	0.66	25.7	513	-	0.07	0.10
qCPA, 3 par	27.93	1284	0.68	-	-	= $b_0$	0.13	0.46
qCPA, 4 par set 1 <sup>d</sup>	28.2	1172	0.64	-	-	23.6	0.49	0.24
qCPA, 4 par, set 2 <sup>d</sup>	28.1	1230	0.64	-	-	25.4	0.42	0.29

<sup>a</sup> 1DÅ = 1Buckingham = 10<sup>-26</sup>esu = 3.3356 · 10<sup>-40</sup>Cm<sup>2</sup>.

<sup>b</sup>%AAD =  $\frac{100}{N^{exp}} \sum_i^{N^{exp}} \left| \frac{x_i^{calc} - x_i^{exp}}{x_i^{exp}} \right|$  where  $x$  stands for  $P^{sat}$  or  $\rho^{liq}$  and  $N^{exp}$  is the number of experimental data.

<sup>c</sup> Not associating.

<sup>d</sup> Parameters based on uncertainties and (indirectly) VLE or LLE data.

ing from -4.1 DÅ to -4.6 DÅ with a typical value of -4.3 DÅ [40–46]. Experimental values of the quadrupolar moment of benzene and acetylene, however, may vary considerably depending on the reference. The experimental quadrupolar moment of benzene range from -9.98 DÅ [40] to -3.6 DÅ [44], although  
175 most of the experimental data is between -9.98 DÅ and -8.5 DÅ [40]. The situation is similar for acetylene which has a large positive quadrupole moment between 3.0 DÅ and 8.4 DÅ [40]. In this work we assume a fixed quadrupolar moment of -9 DÅ for benzene and 4 DÅ for acetylene.

## 3.2. Pure CO<sub>2</sub>

### 3.2.1. Second virial coefficients

A limiting low-density test for a thermodynamic model is to evaluate its capabilities in predicting the second virial coefficient,  $B$ . It may be calculated from the expression

$$\lim_{\rho \rightarrow 0} \left( \frac{\partial Z}{\partial \rho} \right)_T = B \quad (23)$$

Table 2: Correlated CPA and qCPA pure compound parameters and %AAD in saturated liquid densities and saturated pressures. Experimental data from raw DIPPR data [33]. The experimental quadrupole moments of benzene ( $\approx -9$  D $\text{\AA}$ ) and acetylene ( $\approx 4$  D $\text{\AA}$ ) are employed in qCPA.

Compound	Model	Association scheme	$T_r$ (= $T/T_c$ )	$b_0$ mL/mol	$\Gamma(a_0/(Rb_0))$ K	$c_1$ -	$\beta \cdot 1000$ -	$\varepsilon/R$ K	%AAD	
									$P^{sat}$	$\rho^{liq}$
Benzene	qCPA	n.a.	0.5-0.9	75.57	2763.73	0.73	-	-	0.46 (0.68) <sup>a</sup>	0.71 (0.84)
Acetylene	qCPA	n.a.	0.6-0.9	33.54	1469.45	0.65	-	-	0.41 (0.43)	0.52 (0.68)
Acetylene	CPA	n.a.	0.6-0.9	33.28	1576.91	0.7	-	-	0.43	0.68
1-Nonanol	CPA	2B	0.45-0.9	163.79	3508	0.956	0.254	3570.1	0.40	0.79

<sup>a</sup> Numbers in parenthesis are the %AAD with CPA.

where  $Z$  is the compressibility factor and  $\rho$  is the density. The contribution to the second virial coefficient from the SRK term is

$$B^{SRK} = b - \frac{a(T)}{RT} \quad (24)$$

The contribution from the association term can be expressed as

$$B^{assoc} = -S \left[ \exp \left( \frac{\varepsilon^{AB}}{RT} \right) - 1 \right] b\beta^{AB} \quad (25)$$

where the constant  $S$  depend on the association scheme. Its value for several different schemes is shown in table 3.

Table 3: Value of  $S$  in equation (25) for 8 different association schemes. Detailed calculations for scheme 1A, 2B, and 4C can be found in [47]. Nomenclature follows Huang and Radosz [21].

Scheme	1A	2A	2B	3A	3B	4A	4B	4C
S	0.5	2	1	4.5	2	8	3	4

It is straight-forward to show, that the contribution to the second virial coefficient, from the quadrupole term may be expressed as:

$$B^{quad} = \frac{F'_2 - F'_{3,2}}{\rho(1 - F'_{3,2}/F'_2)^2} \quad (26)$$




Figure 1: Model predictions against experimental data for the second virial coefficient of  $\text{CO}_2$  using CPA, n.a., CPA 4C, and the qCPA. Experimental data from [48–54].

where  $F'$  is the Helmholtz energy pr. mole. This expression is similar to that presented by Karakatsani and Economou [11].

185 The second virial coefficient of  $\text{CO}_2$  was predicted with CPA and qCPA using the parameters and schemes presented in table 1. The results are visualized at relatively low temperatures in figure 1 and the %AADs are shown in table 4. All model predictions are in reasonable agreement with the experimental data, although deviations increase for all models at low temperatures, where  
 190 the sharp decrease in the virial coefficient is not fully captured. It is clear from both figure 1 and table 4 that the least convincing result is obtained when  $\text{CO}_2$  is assumed to be self-associating. This may be due to the fact, that  $\text{CO}_2$  is not self-associating and this theoretically unfounded approach may become poor in the low density limit. On the other hand, when qCPA is employed the  
 195 predictions are also slightly worse, than when non associating CPA is employed for  $\text{CO}_2$ . The differences, however, are quite small.

Table 4: Percentage average absolute deviations between experimental and calculated second virial coefficients with the different models.

Model	CPA, n.a.	CPA, 4C	qCPA, 3par	qCPA, 4par, set 1	qCPA, 4par, set 2
%AAD	5.5	13.3	7.3	7.9	7.1

### 3.2.2. Critical Points

Models such as CPA and SAFT, whose parameters are fitted to the saturated liquid density and saturated pressure typically overestimate the critical point. As there are several applications of CO<sub>2</sub> near the critical region, it may be of interest to calculate its critical temperature and pressure, with the different models.

In table 5 the experimental critical temperature and pressure are compared with the calculated critical points for the different CPA modeling approaches. Unfortunately the addition of a quadrupolar term does not improve the prediction of critical points. Both CPA with the 4C scheme and the three versions of qCPA overestimate the critical pressure by approximately 12 bar and the critical temperature by roughly 9 K. The overall best model is CPA n.a., which is essentially equivalent to the SRK with fitted parameters. Comparable critical points were obtained for CO<sub>2</sub> by Diamantonis and Economou [1] using SAFT (309.5 K and 79.2 bar) and PC-SAFT (315.5 K and 90.9 bar) respectively. In both cases CO<sub>2</sub> was treated as a non-associating species.

Table 5: Experimental [55] and predicted critical pressure and temperature of CO<sub>2</sub> using CPA n.a., CPA 4C and qCPA with three different parameter sets.

Model	Exp	CPA n.a.	CPA 4C	qCPA 3par	qCPA 4par, set 1	qCPA 4par, set 2
$T_C$ [K]	304.13	309.7	312.9	312.2	313.5	313.4
$P_C$ [bar]	73.8	81.	86.4	84.9	86.3	85.7



### 3.2.3. Derivative Properties

Despite the industrial need for simultaneous descriptions of phase equilibria  
215 and derivative properties over a wide range of temperatures and pressures, equa-  
tions of state are rarely evaluated for their performance for derivative properties  
[30, 56]. To the best of our knowledge only relatively few systematic evalua-  
tions of the performance of CPA or SAFT for derivative properties have been  
published. Lafitte et al. [57, 58] evaluated the performance of various SAFT  
220 variants, with emphasis on SAFT-VR Mie, for predicting derivative properties  
of alkanes and alcohols and recently, de Villiers et al. [30] predicted derivative  
properties for similar compounds using both CPA, PC-SAFT and SAFT.

Table 6 shows the %AAD values for the density and various derivative prop-  
erties of CO<sub>2</sub> at two temperatures in the compressed liquid region and one in  
225 the critical region. It is gratifying to see that in most cases the deviations for  
each modeling approach are quite similar at the three temperatures. See the  
supporting material for results at saturation.

*Compressed liquid region.* It can be seen from table 6 that in the compressed  
liquid region the predictions with CPA 4C and qCPA are typically similar and  
230 better than CPA n.a. A pronounced exception is the prediction of  $C_V^{res}$  where  
both CPA 4C and qCPA gives poor results and CPA n.a. performs better, at  
least in terms of %AAD. A similar observation was made by de Villiers et al.  
[30] for n-alkanes where the original SAFT performed better than CPA and  
PC-SAFT for the prediction of  $C_V$  but worse for other properties. Using SAFT  
235 and PC-SAFT Diamantonis and Economou [1] calculated derivative properties  
for several small molecules including non-associating CO<sub>2</sub> over an extensive  
temperature range. Both models performed well far from the critical region with  
PC-SAFT being somewhat more accurate. Considering the different models  
and temperature/pressure intervals their results are surprisingly similar to the  
240 results for non-associating CPA. The speed of sound predictions, however, are  
significantly better with the SAFT-type models compared to the CPA-based  
models. This is unsurprising, as it is by now well established that SAFT can

predict the trend of the speed of sound much better than CPA-based models [1, 59].

Table 6: %AAD values of  $\rho^{liq}$ ,  $u$ ,  $C_P^{res}$ ,  $C_V^{res}$ , and  $\mu_{JT}$  for  $\text{CO}_2$  at three reduced temperatures using CPA n.a., CPA 4C and qCPA with three different parameter sets. Pseudo-experimental data from NIST [34].

Models	$T_r$	P range [bar]	% AAD				
			$\rho^{liq}$	$u$	$C_P^{res}$	$C_V^{res}$	$\mu_{JT}$
CPA, n.a.			2.6	12.2	15.2	6.5	17.0
CPA, 4C			1.1	9.8	6.2	42.5	20.1
qCPA, 3par	0.8	100-1000	1.6	10.2	6.3	30.0	12.0
qCPA, 4par, set 1			1.2	9.2	4.6	35.7	13.3
qCPA, 4par, set 2			1.3	9.7	7.1	30.6	14.5
CPA, n.a.			2.7	13.0	8.1	15.9	15.2
CPA, 4C			1.8	10.4	4.7	41.8	12.9
qCPA, 3par	0.9	100-1000	2.0	11.2	3.8	33.9	4.7
qCPA, 4par, set 1			1.7	10.3	2.9	38.7	1.8
qCPA, 4par, set 2			1.7	10.7	4.6	33.8	1.7
CPA, n.a.			3.2	4.9	3.9	32.5	7.3
CPA, 4C			1.6	3.8	6.3	31.4	4.8
qCPA, 3par	1.1	0-250	2.0	4.2	5.4	32.2	5.1
qCPA, 4par, set 1			1.4	4.0	5.9	32.2	4.3
qCPA, 4par, set 2			1.1	4.1	4.7	32.2	4.2

245 Figure 2 shows  $C_V^{res}/R$  (a) and  $C_P^{res}/R$  (b) predictions for  $\text{CO}_2$  at  $T_r = 0.8$  and 0.9. Predictions with CPA 4C and the different qCPA parameter sets are very similar, with qCPA being slightly better. More importantly both model approaches predict quite accurately the intersecting isobaric heat capacities at




Figure 2: Normalized isochoric (a) and isobaric (b) heat capacity predictions of  $\text{CO}_2$  with CPA and qCPA in the compressed liquid region and at  $T_r = 0.8$  and 0.9. Pseudo-experimental data from NIST [34].

approximately 300bar (see figure 2b), whereas CPA n.a. predict this point  
 250 around 800bar. The  $C_V^{res}$  predictions in figure 2a illustrate clearly that most of  
 the models have difficulties with  $C_V^{res}$  predictions.

Notice that while CPA n.a. yields the best  $C_V^{res}$  predictions it is the poorest  
 in terms of  $C_P^{res}$ . However,  $C_P^{res}$  depend on  $C_V^{res}$  as can be seen from equation  
 (B.2) in Appendix B. This suggests that a large part of the improved  $C_P^{res}$   
 255 predictions with qCPA and CPA 4C is due to the over-prediction of  $C_V^{res}$ . In  
 this way at least part of the improvement is in fact due to a cancellation of  
 errors between the  $C_V^{res}$  and the ratio between  $(\partial P/\partial T)_{V,n}^2$  and  $(\partial P/\partial V)_{T,n}$ .

Figure 3a compare the prediction of the liquid density of  $\text{CO}_2$  at  $T_r=0.8$   
 and 0.9 against data from NIST. Figure 3b compares the predictions against  
 260 the experimental data from Brewer et al. [60] in the more restricted pressure  
 range of 200-400 bar and at  $T_r = 0.9$  and  $T_r = 0.93$ . In general the performance  
 is satisfactory for all model approaches, in particular at moderate pressures.  
 The density predictions for CPA n.a. at elevated pressures appear to be worse




Figure 3: Density predictions of CO<sub>2</sub> with CPA and qCPA in the compressed liquid region. At  $T_r = 0.8$  and  $0.9$  (a) compared to pseudo-experimental data from NIST [34] and at  $T_r = 0.9$  and  $T_r = 0.93$  (b) compared to experimental data from Brewer et al.[60].

than for CPA 4C and qCPA. Note from figure 3b, however, that between 200  
 265 and 300 bar the density predictions of CPA n.a. are better than the predictions  
 using CPA 4C. The predictions with qCPA are satisfactory in the whole pres-  
 sure range. See the supporting material for the %AAD values between model  
 predictions and the density values from Brewer et al. [60].

*Critical region.* The critical region is important due to the observed extrema  
 270 and inflection points in the derivative properties, which serves as a valuable test  
 for any EoS. Table 6 shows the %AAD values for the derivative properties of CO<sub>2</sub>  
 at  $T_r = 1.1$  and in a pressure range of 0-250 bar. With the exception of  $C_V^{res}$  the  
 predictions of all model approaches are quite good. Figure 4a clearly shows that  
 the maximum in  $C_V^{res}$  is not predicted by any of the models. A similar result  
 275 was obtained by de Villiers et al [30] for n-alkanes using CPA, SAFT and PC-  
 SAFT. This suggests a fundamental problem with the temperature dependence  
 of both the SRK term and the dispersion term in SAFT. As seen in figure 4b all




Figure 4: Normalized isochoric (a) and isobaric (b) heat capacity predictions of CO<sub>2</sub> with CPA and qCPA in the critical region ( $T_r = 1.1$ ). Pseudo-experimental data from NIST [34].

the model variants evaluated in this work capture the maximum in  $C_P^{res}$  with good accuracy.

280 As the low %AAD in table 6 suggests all model approaches predict the inflection point in the density and the minimum in speed of sound. Especially the density predictions are surprisingly accurate. In both cases CPA n.a. perform slightly worse than the other models.

In summary we have seen that the qualitative difference between the various  
 285 model variants is not particular large for the prediction of derivative properties, and all models typically predicts the same trends. With the clear exception of  $C_V^{res}$  all models generally perform satisfactory considering that the model parameters have not been correlated to any of the properties. While the model differences are smaller than expected, qCPA and CPA with the 4C scheme typically perform slightly better quantitatively than CPA n.a. This may suggest a small improvement in the volume predictions and/or the ratio between  
 290  $(\partial P/\partial T)_{V,n}$  and  $(\partial P/\partial V)_{T,n}$ .

### 3.3. Binary Mixtures

The primary objective of an equation of state is typically the correlation and prediction of the VLE or LLE between mixtures of different compounds. In this section the new qCPA is employed for the prediction ( $k_{ij} = 0$ ) and correlation of several different CO<sub>2</sub> containing binary mixtures. A single temperature independent interaction parameter is employed for the correlations.

#### 3.3.1. CO<sub>2</sub> + n-alkanes

The VLE of several CO<sub>2</sub> + n-alkane mixtures have been predicted using the different modeling approaches. Characteristic results for these predictions are visualized in figures 5-7 and in supplementary materials. The results, in terms of %AAD, are presented in table 7 both for the predictions ( $k_{ij} = 0$ ) as well as when an interaction parameter has been correlated to the VLE data. Moreover the LLE of CO<sub>2</sub> plus a number of heavy alkanes (C<sub>12</sub>, C<sub>14</sub> and C<sub>15</sub>) have been predicted and correlated to experimental data using the different modeling approaches. These results are presented in table 8 and 9.

As described in section 3.1, some of the CO<sub>2</sub>+n-alkane mixtures have been utilized indirectly to obtain the two parameter sets for qCPA when  $b_0^Q$  is used as an additional adjustable. Parameter set 1 for qCPA has been selected partly based on the CO<sub>2</sub> + propane VLE at a single temperature, while parameter set 2 has been selected partly based on the CO<sub>2</sub> + n-dodecane LLE (see supplementary materials for the illustrations).

The CO<sub>2</sub> + ethane VLE is shown in figure 5 at four temperatures. The low temperature azeotrope, observed for this VLE, is predicted very well with qCPA at all temperatures. Particularly when parameter set 1 is employed. CPA n.a. fails to predict the phase behaviour, clearly indicating the need for an improved model for CO<sub>2</sub>.

It should be mentioned that Gross [8] has presented equally good CO<sub>2</sub> + ethane VLE predictions using the PCP-SAFT, although their predictions were compared to simulation results rather than experimental data. Using a group contribution method with a quadrupole term for CO<sub>2</sub> NguyenHuynh et al. [71]

Table 7: Deviations for CPA and qCPA predictions ( $k_{ij} = 0$ ) and correlations ( $k_{ij} \neq 0$ ) of a number of CO<sub>2</sub> + n-alkane VLEs. Including the correlated  $k_{ij}$ . Experimental data from references [61–70].

System	T range [K]	Modeling approach	%AAD in P <sup>a</sup>	% AAD in $y_1$	%AAD in $x_1$	$k_{ij}$	% AAD in P <sup>a</sup>	% AAD in $y_1$	%AAD in $x_1$
$k_{ij} = 0$									
CO <sub>2</sub> (1)+methane(2)	230-293	CPA, n.a.	16.4	11.5	10.2	0.089	1.0	4.6	0.4
		CPA, 4C	4.9	9.0	2.0	0.016	2.3	5.2	0.7
		qCPA, 3par	1.3	8.1	0.9	-0.007	1.2	4.7	0.4
		qCPA, 4par, set 1	12.2	7.4	5.6	-0.057	1.3	4.9	0.5
		qCPA, 4par, set2	6.0	7.6	3.8	-0.032	1.3	4.8	0.4
CO <sub>2</sub> (1)+ethane(2)	213-270	CPA, n.a.	17.5	>100	>100	0.130	3.0	2.7	4.9
		CPA, 4C	9.3	29.7	71.0	0.067	0.3	2.1	1.9
		qCPA, 3par	6.2	18.4	45.0	0.022	0.6	2.5	4.4
		qCPA, 4par, set 1	0.7	2.8	4.4	0.000	0.7	2.8	4.4
		qCPA, 4par, set2	3.1	9.3	22.3	0.017	0.9	3.5	6.2
CO <sub>2</sub> (1)+propane(2)	230-270	CPA, n.a.	29.1	6.4	104.1	0.129	5.0	1.6	11.6
		CPA, 4C	17.7	2.9	50.4	0.074	3.0	2.2	6.6
		qCPA, 3par	12.9	2.1	34.3	0.035	4.3	1.8	10.0
		qCPA, 4par, set 1	3.7	1.9	8.5	0.000	3.7	1.9	8.5
		qCPA, 4par, set2	8.2	1.8	20.0	0.021	4.0	1.9	9.1
CO <sub>2</sub> (1)+butane(2)	250-418	CPA, n.a.	22.6	8.5	59.4	0.124	4.2	11.1	11.5
		CPA, 4C	14.1	6.9	32.7	0.071	2.8	5.3	9.3
		qCPA, 3par	10.3	6.4	24.0	0.045	3.0	5.4	9.8
		qCPA, 4par, set 1	3.3	5.4	10.3	0.000	3.3	5.4	10.3
		qCPA, 4par, set2	7.2	5.9	17.5	0.028	3.3	5.3	9.6
CO <sub>2</sub> (1)+pentane(2)	294-423	CPA, n.a.	19.5	1.9	29.7	0.110	2.1	1.3	2.6
		CPA, 4C	11.8	1.5	16.0	0.065	2.2	1.3	2.7
		qCPA, 3par	7.3	1.4	9.6	0.038	2.1	1.3	2.6
		qCPA, 4par, set 1	2.3	1.3	2.8	0.000	2.3	1.3	2.8
		qCPA, 4par, set2	4.6	1.3	5.9	0.024	2.3	1.3	2.8
CO <sub>2</sub> (1)+hexane(2)	273-303	CPA, n.a.	23.68	-	27.55	0.115	1.83	-	2.44
		CPA, 4C	14.32	-	18.68	0.067	1.02	-	1.4
		qCPA, 3par	8.71	-	11.58	0.037	1.45	-	1.98
		qCPA, 4par, set 1	1.38	-	2.04	0.000	1.38	-	2.04
		qCPA, 4par, set2	5.27	-	7.1	0.021	1.6	-	2.27
CO <sub>2</sub> (1)+decane(2)	277-584	CPA, n.a.	25.82	0.28	34.65	0.103	5.73	0.26	5.87
		CPA, 4C	15.55	0.28	18.79	0.060	2.46	0.26	2.48
		qCPA, 3par	9.1	0.26	10.25	0.028	3.19	0.26	3.34
		qCPA, 4par, set 1	1.93	0.26	1.87	0.000	1.93	0.26	1.87
		qCPA, 4par, set2	5.34	0.27	5.69	0.011	2.96	0.26	3.06

<sup>a</sup> Liquid phase deviation only.




Figure 5: VLE predictions ( $k_{ij} = 0$ ) between  $\text{CO}_2$  and ethane at four temperatures using CPA n.a., CPA 4C, or qCPA with three or four adjustable parameters. Experimental data from [62, 63].

also presented excellent results for the  $\text{CO}_2$  + ethane and  $\text{CO}_2$  + propane systems. These results, however, are not directly comparable to the results in this work, as the authors correlated the  $\text{CO}_2$  parameters to the  $\text{CO}_2$  + propane VLE in addition to the saturated liquid density and vapor pressure. Moreover a non-zero  $k_{ij}$  is employed; its value being obtained from another correlation. The predictions and correlations presented by Tsivintzelis et al. [18] for  $\text{CO}_2$  + n-alkane mixtures with CPA are obviously similar to the CPA 4C and CPA n.a. results presented in this work.

Figures 5-7 illustrate that the series behaviour from  $\text{C}_2$ - $\text{C}_{10}$  is captured quite well with qCPA, and the predictions with qCPA continues to be excellent, especially for parameter set 1. Unfortunately all models overestimate the critical point (see figure 6). As already mentioned this is an unfortunate characteristic of such equations of state. Cross-over approaches exist which may deal with the problem, at the cost of more complex expressions and additional parameters. Figure 6 show the  $\text{CO}_2$  + n-butane VLE prediction. Gross [8] also showed very






Figure 6: VLE predictions ( $k_{ij} = 0$ ) between  $\text{CO}_2$  and n-butane at three temperatures using CPA n.a., CPA 4C, or qCPA with three or four adjustable parameters. Experimental data from [64, 66].

good results for this system as well as for the  $\text{CO}_2$  + n-pentane VLE using the PCP-SAFT. In these cases a  $k_{ij}$  of approximately 0.04 was needed to correlate  
 340 both systems. Using qCPA with parameter set 1, however, the best  $k_{ij}$  is approximately zero. A more fair comparison, considering the number of adjustable parameters, may be to compare the optimum  $k_{ij}$  for the three parameter version of the qCPA (see table 7) with those found for PCP-SAFT. These  $k_{ij}$ 's are of a similar magnitude despite the different base models. Close to the critical point,  
 345 however, PCP-SAFT seem to perform better.

Generally the qCPA approaches with four parameters appear to yield the best VLE predictions for  $\text{CO}_2$  + n-alkanes, followed by the three parameter version of qCPA, then CPA 4C and finally by CPA n.a. An exception to this trend is the  $\text{CO}_2$  + methane system which is shown in figure 8. Except in the vicinity  
 350 of the critical point quite good predictions are obtained when  $\text{CO}_2$  is considered an associating species and when qCPA with three adjustable parameters is employed. However, when qCPA is employed with an additional adjustable parameter rather poor predictions are obtained.




Figure 7: VLE predictions ( $k_{ij} = 0$ ) between CO<sub>2</sub> and n-decane at three temperatures using CPA n.a., CPA 4C, or qCPA with three or four adjustable parameters. Experimental data from [65].




Figure 8: VLE predictions ( $k_{ij} = 0$ ) between CO<sub>2</sub> and methane at two temperatures using CPA n.a., CPA 4C, or qCPA with three or four adjustable parameters. Experimental data from [62].

As the LLE between CO<sub>2</sub> and heavy hydrocarbons is believed to be, partly,  
 355 due to the quadrupolar moment of CO<sub>2</sub> a successful quadrupolar term should  
 improve the models ability to correlate these LLEs. A characteristic correla-  
 tion of the CO<sub>2</sub> + n-pentadecane LLE is visualized in figure 9a. All model  
 approaches can correlate the LLE of CO<sub>2</sub> and the heavy alkane. A much larger  
 interaction parameter, however, is needed when CO<sub>2</sub> is modelled as an inert  
 360 or self-associating compound, compared to when CO<sub>2</sub> is modelled with qCPA.  
 The correlated interaction parameter and the results, in terms of %AAD, are  
 shown in table 8. As is typically the case, the LLE results are very sensitive  
 to the binary interaction parameter, and many models do not even predict the  
 presence of a LLE without this parameter. As figure 9b illustrate, however, the  
 365 four parameter version of qCPA can predict ( $k_{ij} = 0$ ) the LLE of these mixtures  
 fairly well. The %AAD of the predictions are shown in table 9.

Table 8: Correlated  $k_{ij}$  and calculated %AADs for the CO<sub>2</sub> + heavy n-alkane  
 LLE systems using the CPA and qCPA. Experimental data from [72].

System	P range [bar]	Model	$k_{ij}$	%AAD in HC phase	%AAD in CO <sub>2</sub> phase	Average %AAD in CO <sub>2</sub>
CO <sub>2</sub> (1)+n-dodecane (2)	20-29	CPA n.a.	0.102	0.59	0.39	0.49
		CPA 4C	0.054	0.42	0.18	0.30
		qCPA 3par	0.019	0.47	0.3	0.39
		qCPA 4par, 1	-0.025	0.42	0.25	0.34
		qCPA 4par, 2	-0.003	0.44	0.28	0.36
CO <sub>2</sub> (1) +n-tetradecane (2)	31-83	CPA n.a.	0.100	0.31	0.06	0.19
		CPA 4C	0.058	0.29	0.11	0.20
		qCPA 3par	0.023	0.24	0.08	0.16
		qCPA 4par, 1	-0.017	0.35	0.11	0.23
		qCPA 4par, 2	0.003	0.28	0.11	0.19
CO <sub>2</sub> (1) +n-pentadecane (2)	32-80	CPA n.a.	0.105	0.25	0.04	0.14
		CPA 4C	0.059	0.15	0.12	0.13
		qCPA 3par	0.025	0.07	0.06	0.07
		qCPA 4par, 1	-0.017	0.11	0.11	0.11
		qCPA 4par, 2	0.004	0.09	0.09	0.09

### 3.3.2. CO<sub>2</sub> + self-associating compounds - Approach to induced association

When mixtures of CO<sub>2</sub> + self-associating compounds, such as alcohols and  
 water, are considered it raises the important question of whether CO<sub>2</sub> should be  
 370 modelled only as a quadrupolar molecule or as a quadrupolar *and* a solvating

Table 9: %AADs for the predicted LLEs of CO<sub>2</sub> + heavy n-alkane systems using the four parameter version of qCPA. Experimental data from [72].

System	P range [bar]	Model	%AAD in hydrocarbon phase	%AAD in CO <sub>2</sub> phase	Average % AAD in CO <sub>2</sub>
CO <sub>2</sub> (1)+n-dodecane (2)	20-29	qCPA set 1	2.49	$k_{ij} = 0$ 0.53	1.51
		qCPA set 2	0.57	0.34	0.45
CO <sub>2</sub> (1) +n-tetradecane (2)	31-83	qCPA set 1	1.44	0.13	0.79
		qCPA set 2	0.28	0.11	0.16
CO <sub>2</sub> (1) +n-pentadecane (2)	32-80	qCPA set 1	1.37	0.07	0.72
		qCPA set 2	0.35	0.14	0.24




Figure 9: Correlation (a) and prediction (b) of the CO<sub>2</sub>+n-pentadecane LLE, where CO<sub>2</sub> is modelled with CPA n.a., CPA 4C, or qCPA with three or four adjustable parameters. Experimental data from [72].

molecule (i.e. a molecule with electron donor *or* electron acceptor sites). Several studies indicate Lewis acid-Lewis base type interactions for CO<sub>2</sub> with polar, associating molecules such as water and alcohols [18, 73–76]. As an engineering approach the Lewis acid-Lewis base interaction can be viewed as an induced  
375 cross-association (solvation). This kind of cross-association, where one component self-associates but the other only has electron donor or electron acceptor sites, is difficult to treat with current models since the association parameters,  $\varepsilon^{A_i B_i}$  and  $\beta^{A_i B_i}$ , are available only for the self-associating compound. That is, while the compounds cross-associate with each other, the cross-association volume cannot be determined from the CR-1 combining rule (equation (11b)). As  
380 a pragmatic solution the cross-association volume,  $\beta^{A_i B_j}$ , is often fitted to the experimental data (simultaneously with the binary interaction parameter,  $k_{ij}$ ) while the association energy,  $\varepsilon^{A_i B_i}$ , of the solvating compound is set to zero. This approach typically works well, however, the improved correlations may, at  
385 least partially, be attributed to an increased flexibility due to a higher number of adjustable parameters.

A simple alternative to correlating the cross-association volume,  $\beta^{A_i B_j}$ , for solvating mixtures was proposed by Kleiner and Sadowski [77]. The method assumes that the cross-association volume can be set equal to the association  
390 volume of the self-associating compound. In this way equations (11a)-(11b) can be directly employed without the need for any adjustable parameters beyond a  $k_{ij}$ . Clearly improved correlations can be obtained if both the binary interaction parameter and the cross-association volume are fitted to the experimental data. The added model flexibility of such an approach would, however, make it difficult  
395 to critically compare the models.

Another important question is how many solvation sites CO<sub>2</sub> has. Indeed in a recent work NguyenHuynh et al. [78], illustrated, using the GC-PPC-SAFT, that very good results could be obtained for mixtures of CO<sub>2</sub> and alcohols if CO<sub>2</sub> was assumed to have two cross sites. Similarly Figure 10 illustrate the  
400 predictions ( $k_{ij} = 0$ ) with qCPA for the CO<sub>2</sub> + ethanol mixture and the vapor liquid liquid equilibrium (VLLE) of the CO<sub>2</sub>-rich phase of the CO<sub>2</sub> + water




Figure 10: Predicted ( $k_{ij} = 0$ ) VLEs of the CO<sub>2</sub> + ethanol system (a) and the CO<sub>2</sub>-rich vapor phase of the CO<sub>2</sub> + water system (b) with qCPA with three parameters and assuming zero, one or two solvation sites. It is assumed for both systems that  $\beta_{crs} = \beta_{assoc}$ . Experimental data from [79–85].

405 mixture using zero, one or two cross association sites. The cross association parameters were determined using the approach by Kleiner and Sadowski [77]. When cross association is not taken into account, or only one cross association site is considered, the equilibrium pressure in figure 10a is overestimated and a false liquid-liquid phase split is predicted. Similarly the minimum in the water solubility in the CO<sub>2</sub>-rich phase is captured quite accurately when two cross-association sites are assumed, but not when CO<sub>2</sub> has one or zero sites (see figure 10b). That is, when qCPA is employed CO<sub>2</sub> is assumed to have  
 410 two solvation sites, and the cross-association parameters are found using the approach of Kleiner and Sadowski [77].

In the case where we consider CO<sub>2</sub> to be a self-associating compound the cross-association between CO<sub>2</sub> and the associating compound is handled by directly applying the CR-1 mixing rule (see eq. (11a)-(11b)).

415  $CO_2 + \text{alcohols}$ . In general qCPA performs quite well for many of the alcohol systems, even with a zero  $k_{ij}$ , but a non-zero temperature independent interaction parameter is used to better represent the phase equilibria. The results are summarized in terms of %AADs in table 10. Characteristic VLE results are shown for  $CO_2 + \text{methanol}$  in figure 11 at three temperatures. Despite  
420 a very large binary interaction parameter the highest deviations are obtained when  $CO_2$  is considered a self-associating molecule following the 4C association scheme. The model which captures the trend of the VLEs most accurately is clearly qCPA. The correlations with the different qCPA parameter sets for  $CO_2$  are similar, although a smaller interaction parameter (in absolute terms) is  
425 needed when qCPA with three adjustable parameters is employed. In all cases a small negative interaction parameter is necessary, whereas a positive interaction parameter is needed when  $CO_2$  is considered either self-associating or inert. Surprisingly good correlations, with a low interaction parameter, are obtained when  $CO_2$  is considered an inert. The model, however, does not capture the  
430 trend of the experimental data quite as well as qCPA, especially at higher temperatures where the pressure is overestimated and a false liquid-liquid split may occur.

Figure 12 illustrate that the phase behaviour description of the simultaneous VLE and LLE of the  $CO_2 + \text{nonanol}$  system is satisfactory with all models  
435 using a temperature independent interaction parameter. The same is the case for the  $CO_2 + \text{octanol}$  system. Notice that when qCPA with three parameters is employed for the  $CO_2 + \text{octanol}$  system the optimum  $k_{ij}$  is found to be zero, i.e. the model is predictive for this system. It is noteworthy, that as the chain length of the alcohols increases the binary interaction parameter tend to decrease or  
440 increase when  $CO_2$  is treated as either a self-associating or an inert compound respectively. When  $CO_2$  is treated as a quadrupolar molecule, however, the small interaction parameter seem to be almost constant for the various systems.

$CO_2 + \text{water}$ . The capabilities of association theories for modeling the important  $CO_2 + \text{water}$  mixture has been extensively studied in recent literature using




Figure 11: Correlation of the CO<sub>2</sub> + methanol VLE using CPA n.a., CPA 4C, or qCPA with three or four adjustable parameters. Experimental data from [80, 81, 86–88].




Figure 12: Correlation of the CO<sub>2</sub> + nonanol VLE and LLE using CPA n.a., CPA 4C, or qCPA with three or four adjustable parameters. Experimental data from [81, 90].



Table 10: CPA and qCPA deviations and  $k_{ij}$  for CO<sub>2</sub> + alcohol VLE and LLE mixtures. Experimental data from references [79–81, 86–89].

System	T range (K)	Modeling approach	$k_{ij}$	% AAD in P	% AAD in $y_1$	%AAD in $x_1$
CO <sub>2</sub> (1)+methanol(2)	230-313.2	CPA, n.a.	0.027	7.9	0.4	10.5
		CPA, 4C	0.180	18.7	0.5	48.6
		qCPA, 3par	-0.010	4.7	0.2	8.9
		qCPA, 4par, set 1	-0.050	5.6	0.2	9.7
		qCPA, 4par, set2	-0.050	5.6	0.2	10.1
CO <sub>2</sub> (1)+ethanol(2)	291.15-313.2	CPA, n.a.	0.050	9.4	0.4	10.1
		CPA, 4C	0.200	6.1	0.2	7.2
		qCPA, 3par	-0.019	3.9	0.4	8.3
		qCPA, 4par, set 1	-0.057	2.7	0.3	6.5
		qCPA, 4par, set2	-0.030	2.9	0.3	6.5
CO <sub>2</sub> (1)+propanol(2)	313.4	CPA, n.a.	0.058	10.7	0.2	7.4
		CPA, 4C	0.180	6.7	0.1	11.0
		qCPA, 3par	-0.020	4.1	0.2	5.1
		qCPA, 4par, set 1	-0.050	3.4	0.2	3.2
		qCPA, 4par, set2	-0.030	3.0	0.2	3.1
CO <sub>2</sub> (1)+octanol(2) <sup>a</sup>	308.2-328.2	CPA, n.a.	0.070	12.9	0.4	10.8
		CPA, 4C	0.160	14.8	0.3	10.3
		qCPA, 3par	0.000	11.0	0.3	7.4
		qCPA, 4par, set 1	-0.040	10.4	0.3	6.6
		qCPA, 4par, set2	-0.020	10.8	0.3	7.7
CO <sub>2</sub> (1)+nonanol(2) <sup>a</sup>	308.1-328.2	CPA, n.a.	0.070	19.6	0.8	9.1
		CPA, 4C	0.140	23.5	0.5	13.7
		qCPA, 3par	0.000	18.5	0.8	6.4
		qCPA, 4par, set 1	-0.045	17.1	0.7	6.4
		qCPA, 4par, set2	-0.025	18.1	0.7	7.6

<sup>a</sup> Deviations based on both VLE and LLE data.

445 both CPA [16, 18, 91, 92] and SAFT-based models [16, 91, 93–95]. The mixture has also been studied, with mixed success, using some of the multipolar SAFT-based equations of state [96–98]. It is clear from most investigations, however, that the best results for CO<sub>2</sub> + water are almost always obtained when CO<sub>2</sub> is considered to be a solvating molecule.

450 As stated previously we assume that CO<sub>2</sub> cross-associates with two solvation sites and that  $\beta^{A_i B_j} = \beta_{crs} = \beta_{water}$ . Figure 13 illustrates the correlation of the CO<sub>2</sub>-rich vapor phase and the water-rich liquid phase at 323.15K. All models correlate the solubility of CO<sub>2</sub> in the water-rich phase quite well (figure 13a), although a large interaction parameter is needed when CO<sub>2</sub> is assumed to be a self-associating compound. However, as also demonstrated by Tsivintzelis et al. [18], CPA cannot capture the minimum in the solubility of water in the vapor phase when CO<sub>2</sub> is modelled as an inert compound, nor is the trend of the experimental data captured if CO<sub>2</sub> is modeled as a self-associating compound (see figure 13b). When CO<sub>2</sub> is treated as a quadrupolar compound which cross-associates with water the minimum in solubility in the CO<sub>2</sub> phase is, at least  
460 qualitatively, captured.

Instead of treating CO<sub>2</sub> as a purely inert compound a more fair model comparison may be to also assume CO<sub>2</sub> to cross-associate, using the same assumptions as we did for qCPA. Figure 14 compares the model correlations in the  
465 CO<sub>2</sub>-rich vapor phase using either CPA or qCPA with three adjustable parameters where CO<sub>2</sub> is assumed to solvate with water in both cases. It can be seen from the figure that if CO<sub>2</sub> is assumed to be solvating, then the minimum in the water solubility is captured equally well in CPA and qCPA. This suggests that taking cross-association between CO<sub>2</sub> and water into account is in fact the  
470 dominant contribution to modeling the CO<sub>2</sub> + water system. We note that to model both phases simultaneously a relatively high  $k_{ij}$  is needed, when the quadrupolar nature of CO<sub>2</sub> is not explicitly taken into account. Table 11 summarizes the results in terms of %AAD both with and without an interaction parameter.




Figure 13: Correlation of the  $\text{CO}_2$  solubility in the water rich liquid phase (a) and the water solubility in the  $\text{CO}_2$  rich vapor phase (b) for the  $\text{CO}_2 + \text{water}$  system. CPA n.a., CPA 4C, or qCPA with three or four adjustable parameters is employed. Experimental data from [82–85, 99].

Table 11: Deviations for CPA and qCPA predictions ( $k_{ij} = 0$ ) and correlations ( $k_{ij} \neq 0$ ) for the  $\text{CO}_2(1) + \text{water}(2)$  mixture at 323.15 K (both VLE and LLE data). Including the correlated  $k_{ij}$ . Experimental data from references [82–85, 99]

Modeling approach	% AAD in $y_2$	%AAD in $x_1$	$k_{ij}$	% AAD in $y_2$	%AAD in $x_1$
$k_{ij}=0$					
CPA, n.a.	62.5	4.6	0.005	62.9	3.6
CPA with solvation	10.3	>100	0.128	21.5	4.2
CPA, 4C	>100	>100	0.255	>100	4.5
qCPA, 3par	13.7	25.9	0.033	17.1	3.7
qCPA, 4par, set 1	17.0	11.5	-0.015	15.5	3.6
qCPA, 4par, set2	14.7	7.5	0.010	15.8	3.5




Figure 14: Correlation of the water solubility in the  $\text{CO}_2$  rich vapor phase for the  $\text{CO}_2 + \text{water}$  system. Comparing the performance of CPA with solvation, and qCPA with solvation. Experimental data from [82–85].

475 *Mixtures with two quadrupolar compounds.*  $\text{CO}_2$  has so far been the only quadrupolar compound explicitly considered in this work, although both water and ethane have a quadrupole moment, which we have ignored. It is important, however, to evaluate the capabilities of the new model for mixtures of two or more quadrupolar compounds. More specifically such an evaluation may provide an indication  
 480 as to the adequacy of the proposed mixing rules and cross-quadrupolar interactions. Here we investigate mixtures of  $\text{CO}_2$  and either benzene or acetylene.

A fixed experimental quadrupole moment of  $-9 \text{ D}\text{\AA}$  is employed for benzene. See table 2 for the re-estimated qCPA pure compound parameters. The predicted VLE of benzene and  $\text{CO}_2$  is compared to experimental values in figure  
 485 15. All model approaches, except when both  $\text{CO}_2$  and benzene are treated as inert compounds, perform very well and quite predictively. Interestingly CPA performs very well when  $\text{CO}_2$  is assumed to be associating, despite the fact that both  $\text{CO}_2$  and benzene are solvating molecules rather than associating ones.




Figure 15: Prediction of the CO<sub>2</sub> + benzene VLE at two temperatures (upper: 347.25 K and lower: 315.45 K) using CPA n.a., CPA 4C, or qCPA with three or four adjustable parameters. Experimental data from [86, 100].

The results in terms of %AAD are presented in table 12. Gross [8] modelled  
 490 this system using the PCP-SAFT and regular PC-SAFT, however, the results  
 with the former model were worse than with PC-SAFT. To improve the results  
 Gross had to set the cross-quadrupolar interactions to zero.

Acetylene is modelled with a fixed quadrupolar moment of 4 D<sup>2</sup>Å the re-  
 estimated pure compound parameters can be found in table 2. The VLE between  
 495 CO<sub>2</sub> + acetylene results in an uncommon negative azeotrope, which is believed  
 to be caused by the opposite sign of the quadrupole moments for the two com-  
 ponents [102]. Ideally a quadrupolar term would be able to model this effect.  
 Figure 16 shows, however, that a positive azeotrope is predicted by both qCPA  
 and CPA 4C. All models are capable of correlating the azeotrope. The reason  
 500 that this quadrupole-quadrupole pair is so poorly represented by the quadrupolar  
 model is believed to be due to the fact that all quadrupolar moments are  
 raised to even powers (see Eq. (14)-(16)). The sign of the quadrupole moment

Table 12: Deviations for CPA and qCPA predictions ( $k_{ij} = 0$ ) and correlations ( $k_{ij} \neq 0$ ) for the  $\text{CO}_2 + \text{benzene}$  mixture. Including the correlated  $k_{ij}$ . Experimental data from [86, 100].

T range (K)	Modeling approach	% AAD in P	%AAD in $x_1$	$k_{ij}$	% AAD in P	%AAD in $x_1$
$k_{ij}=0$						
298.2-347.3	CPA, n.a.	26.8	23.9	0.068	14.3	6.2
	CPA, 4C	15.8	9.5	0.020	12	3.9
	qCPA, 3par	17.8	13.3	0.008	10.5	3.4
	qCPA, 4par, set 1	11.9	5.7	-0.038	12.0	3.8
	qCPA, 4par, set2	15.8	10.9	-0.014	12.0	3.8




Figure 16: Prediction of the  $\text{CO}_2 + \text{acetylene}$  VLE at 233 K using CPA n.a., CPA 4C, or qCPA with three or four adjustable parameters. Experimental data from [101].

thus becomes inconsequential and  $\text{CO}_2 + \text{acetylene}$  is effectively modeled as two quadrupoles of the same sign. The preferred molecular orientation for two quadrupoles of opposite sign, however, is end to end whereas it is perpendicular if they are of the same sign.

#### 4. Conclusion

In an effort to improve the predictive capabilities of classic thermodynamic models, primarily for mixtures containing  $\text{CO}_2$ , a quadrupolar equation of state contribution, developed from statistical mechanics [5], have been modified and combined with CPA. The resulting qCPA can be used without introducing any other adjustable pure compound parameters or, alternatively, an additional adjustable parameter can be used. The model results were compared with CPA n.a. and CPA 4C.

When the same number of adjustable parameters are used for CPA n.a. and qCPA, a systematic improvement in the correlation of the saturated liquid density and vapor pressure is observed. This is particularly true for the strongly quadrupolar  $\text{CO}_2$  molecule, but applies to benzene and acetylene as well.

The models were employed to predict various pure compound properties for  $\text{CO}_2$  including the second virial coefficient, the critical properties and various derivative properties. qCPA and CPA 4C seem to perform slightly better than CPA n.a. in regards to the derivative properties. On the other hand CPA n.a. is slightly more accurate in representing the second virial coefficient and the critical point. The differences between the modelling approaches, however, are generally small. The main reason for deviations in the derivative properties seem to be caused by a deficiency in the second order derivatives.

It was found that qCPA (all approaches) significantly improves the prediction of binary VLE and the correlation of LLE between  $\text{CO}_2 + \text{n-alkane}$  mixtures. A significantly smaller interaction parameter is needed to correlate the experimental data with qCPA compared to the CPA approaches. When it comes to  $\text{CO}_2 + \text{associating mixtures}$  very good correlations were obtained with qCPA

when solvation was taken into account using the approach suggested by Kleiner and Sadowski [77] and a single small binary interaction parameter. The model was finally applied to two quadrupolar-quadrupolar mixtures, which illustrated  
535 that the model seem to have difficulties for quadrupolar mixtures of opposite sign such as CO<sub>2</sub> + acetylene. Overall explicitly accounting for the quadrupolar forces appears to offer significantly improved predictions, and better (smaller  $k_{ij}$ ) correlations, for CO<sub>2</sub> containing mixtures compared to the two other CPA approaches.

540 The performance of the qCPA approaches is similar with regards to the derivative properties of CO<sub>2</sub> and for mixtures of CO<sub>2</sub> + self-associating compounds. For CO<sub>2</sub> + n-alkanes the four-parameter versions of qCPA both perform somewhat better than the three-parameter version. Nevertheless this modest improvement may not justify the increased model flexibility and uncertainty  
545 in the parameter estimation. Unless the primary focus is binary CO<sub>2</sub> + n-alkane mixtures the most promising approach is thus considered to be qCPA with three adjustable parameters.

## Appendix A. Employed pure compound parameters

The pure compound parameters from literature which has been used in this  
550 work are shown in table A.13.



Table A.13: CPA pure compound parameters from literature employed in this work.

Compound	Association scheme	$b_0$ mL/mol	$\Gamma(a_0/(Rb_0))$ K	$c_1$ -	$\beta \cdot 1000$ -	$\varepsilon/R$ K	Reference
Methane	n.a.	29.10	959.03	0.45	-	-	[103]
Ethane	n.a.	42.90	1544.55	0.58	-	-	[103]
Propane	n.a.	57.83	1896.45	0.63	-	-	[104]
n-Butane	n.a.	72.08	2193.08	0.71	-	-	[104]
n-Pentane	n.a.	91.01	2405.11	0.80	-	-	[104]
n-Hexane	n.a.	107.89	2640.03	0.83	-	-	[104]
n-Decane	n.a.	178.65	3190.54	1.13	-	-	[104]
n-dodecane	n.a.	216.24	3471.04	1.20	-	-	[105]
n-tetradecane	n.a.	250.53	3678.42	1.29	-	-	[105]
n-pentadecane	n.a.	274.53	3751.99	1.34	-	-	[105]
Benzene	n.a.	74.99	2867.19	0.76	-	-	[104]
Water	4C	14.52	1017.34	0.67	69.20	2003.25	[29]
Methanol	2B	30.98	1573.71	0.43	16.10	2957.78	[29]
Ethanol	2B	49.11	2123.83	0.74	8.00	2589.85	[106]
1-Propanol	2B	64.11	2234.52	0.92	8.10	2525.86	[106]
1-Octanol	2B	148.80	3367.99	1.15	0.14	3218.55	[106]

## Appendix B. Equations for the derivative properties

This appendix list expressions for the derivative properties evaluated in this work, namely the residual isochoric- and isobaric heat capacities, the residual enthalpy, the Joule-Thomson coefficient and the speed of sound. The properties  
 555 are given in terms of the reduced residual Helmholtz energy.

*Residual isochoric heat capacity.*

$$C_V^{res}(T, V, \mathbf{n}) = -RT^2 \left( \frac{\partial^2 F}{\partial T^2} \right)_{V, \mathbf{n}} - 2RT \left( \frac{\partial F}{\partial T} \right)_{V, \mathbf{n}} \quad (\text{B.1})$$

*Residual isobaric heat capacity.*

$$C_P^{res}(T, V, \mathbf{n}) = C_V^{res} - T \frac{\left( \frac{\partial P}{\partial T} \right)_{V, \mathbf{n}}^2}{\left( \frac{\partial P}{\partial V} \right)_{T, \mathbf{n}}} - nR \quad (\text{B.2})$$

*Residual enthalpy.*

$$\frac{H^{res}(T, V, \mathbf{n})}{nRT} = Z - \frac{T}{N} \left( \frac{\partial F}{\partial T} \right)_{V, \mathbf{n}} - 1 \quad (\text{B.3})$$

*The Joule-Thomson coefficient.*

$$\mu_{JT} = \left( \frac{\partial T}{\partial P} \right)_{H, \mathbf{n}} = -\frac{1}{C_P} \left[ V + T \left( \frac{\partial P}{\partial T} \right)_{V, \mathbf{n}} / \left( \frac{\partial P}{\partial V} \right)_{T, \mathbf{n}} \right] \quad (\text{B.4})$$

*The speed of sound.*

$$u = \sqrt{-V^2 \frac{C_P}{C_V} \frac{\left( \frac{\partial P}{\partial V} \right)_{T, \mathbf{n}}}{M_W}} \quad (\text{B.5})$$

where

$$\left( \frac{\partial P}{\partial V} \right)_{T, \mathbf{n}} = -RT \left( \frac{\partial^2 F}{\partial V^2} \right)_{T, \mathbf{n}} - \frac{nRT}{V^2} \quad (\text{B.6})$$

$$\left( \frac{\partial P}{\partial T} \right)_{V, \mathbf{n}} = -RT \left( \frac{\partial^2 F}{\partial V \partial T} \right)_{\mathbf{n}} - \frac{P}{T} \quad (\text{B.7})$$

and  $M_W$  is the molecular weight. It is seen that the second order derivatives  $(\partial^2 F / \partial V^2)_{\mathbf{n}, T}$ ,  $(\partial^2 F / \partial V \partial T)_{\mathbf{n}}$  and  $(\partial^2 F / \partial T^2)_{\mathbf{n}, V}$  are crucial in order to obtain accurate derivative property predictions.

The enthalpy and the heat capacities are expressed as a sum of two con-  
560 tributions; an ideal, temperature dependent, contribution (ig) and a residual  
contribution (res) obtained from an EoS. In this work the  $C_P^{ig}$  is calculated from  
the DIPPR correlation [33]. The ideal isochoric heat capacity and enthalpy is  
calculated through their relation to  $C_P^{ig}$ .

### Abbreviations

565 **CPA** cubic plus association.

**CS** Carnahan-Starling.

**DIPPR** Design Institute for Physical Properties.

**EoS** equation of state.

**LLE** liquid liquid equilibrium.

570 **NIST** National Institute of Standards and Technology.

**PC-SAFT** Perturbed-Chain SAFT.

**PCP-SAFT** Perturbed-Chain Polar SAFT.

**qCPA** quadrupolar CPA.

**RDF** Radial distribution function.

575 **SAFT** Statistical Association Fluid Theory.

**SRK** Soave-Redlich-Kwong.

**tPC-PSAFT** truncated Perturbed-Chain Polar SAFT.

**VLE** vapor liquid equilibrium.

**VLLE** vapor liquid liquid equilibrium.

580 **Nomenclature**

$A^r(T, V, \mathbf{n})$  Residual Helmholtz energy.

$A_2^r$  Second order term in the perturbation expansion for quadrupolar interactions.

585  $A_3^r$  Third order term in the perturbation expansion for quadrupolar interactions.

$A_i$  bonding sites on molecule  $i$ .

$B$  Second virial coefficient or expression for co-volume mixing term..

$C_P^{res}$  The residual isobaric heat capacity.

$C_V^{res}$  The residual isochoric heat capacity.

590  $D$  Mixing rule for the energetic part of SRK.

$I_{TQ}$  Three-body correlation integral.

$I_n$  Two-body correlation integral.

$N_A$  Avogadro's constant.

$P_C$  Critical pressure.

595  $P$  Pressure.

$Q$  Quadrupole moment.

$R$  The gas constant.

$S$  Constant in expression for the second virial coefficient of the association term..

600  $T_C$  Critical temperature.

$T_r$  Reduced temperature  $T_r = T/T_C$ .

$T$  Temperature in Kelvin.

$V$  Volume.

$X_{A_i}$  Fraction of sites A on molecule  $i$  which do *not* form bonds with other sites  
 605 (site monomer fraction).

$Z$  compressibility factor.

$\Delta^{A_i B_j}$  Association strength between site A on molecule  $i$  and site B on molecule  
 $j$ .

$\beta^{A_i B_j}$  cross-association volume.

610  $\eta$  Reduced density.

$\mathbf{n}$  Vector of molar composition.

$\mu_{JT}$  The Joule-Thomson coefficient.

$\rho$  Density.

$\sigma$  Hard-sphere diameter.

615  $\varepsilon^{A_i B_j}$  cross-association energy.

$a_0$  Pure compound energetic parameter in the SRK term.

$a_{ij}$  Cross energetic parameter in the SRK term.

$b_0^Q$  co-volume parameter in the quadrupolar term.

$b_0$  Pure compound co-volume.

620  $b_{ii}$  Pure compound co-volume for component  $i$ .

$b_{ij}$  cross-covolume.

$c_1$  Pure compound parameter in the SRK term.

$g$  Radial distribution function.

$k_b$  Boltzmann's constant.

625  $k_{ij}$  Binary interaction parameter.

$n$  Total composition.

$s^{exp}$  standard deviation.

$u$  The speed of sound.

### Acknowledgements

630 The authors gratefully acknowledge the Danish Research Council for Independent Research for funding this work as part of the project; 'CO<sub>2</sub> Hydrates - Challenges and Possibilities'.

### References

- [1] N. I. Diamantonis, I. G. Economou, Evaluation of statistical associating fluid theory (saft) and perturbed chain-saft equations of state for the  
635 calculation of thermodynamic derivative properties of fluids related to carbon capture and sequestration, *Energy & Fuels* 25 (7) (2011) 3334–3343. arXiv:<http://dx.doi.org/10.1021/ef200387p>, doi:10.1021/ef200387p.  
640 URL <http://dx.doi.org/10.1021/ef200387p>
- [2] G. Stell, J. Rasaiah, H. Narang, Thermodynamic perturbation theory for simple polar fluids, i, *Molecular Physics* 23 (2) (1972) 393–406. arXiv:  
<http://www.tandfonline.com/doi/pdf/10.1080/00268977200100381>,  
doi:10.1080/00268977200100381.  
645 URL <http://www.tandfonline.com/doi/abs/10.1080/00268977200100381>
- [3] G. Rushbrooke, G. Stell, J. Hye, Theory of polar liquids, *Molecular Physics* 26 (5) (1973) 1199–1215. arXiv:  
<http://www.tandfonline.com/doi/pdf/10.1080/00268977300102411>,  
650 doi:10.1080/00268977300102411.

URL <http://www.tandfonline.com/doi/abs/10.1080/00268977300102411>

- [4] G. Stell, J. Rasaiah, H. Narang, Thermodynamic perturbation theory for simple polar fluids. ii, *Molecular Physics* 27 (5) (1974) 1393–1414. arXiv:  
655 <http://www.tandfonline.com/doi/pdf/10.1080/00268977400101181>,  
doi:10.1080/00268977400101181.

URL <http://www.tandfonline.com/doi/abs/10.1080/00268977400101181>

- [5] B. Larsen, J. Rasaiah, G. Stell, Thermodynamic perturbation theory for multipolar and ionic liquids, *Molecular Physics* 33 (4) (1977) 987–1027. arXiv:  
660 <http://www.tandfonline.com/doi/pdf/10.1080/00268977700100901>, doi:10.1080/00268977700100901.

URL <http://www.tandfonline.com/doi/abs/10.1080/00268977700100901>

- [6] K. Gubbins, C. Twu, Thermodynamics of polyatomic fluid mixtures: i theory, *Chemical Engineering Science* 33 (7) (1978) 863 – 878.  
665 doi:10.1016/0009-2509(78)85176-8.

URL <http://www.sciencedirect.com/science/article/pii/0009250978851768>

- [7] C. Twu, K. Gubbins, Thermodynamics of polyatomic fluid mixtures: ii: Polar, quadrupolar and octopolar molecules, *Chemical Engineering Science* 33 (7) (1978) 879 – 887. doi:10.1016/0009-2509(78)85177-X.  
670

URL <http://www.sciencedirect.com/science/article/pii/000925097885177X>

- [8] J. Gross, An equation-of-state contribution for polar components: Quadrupolar molecules, *AIChE Journal* 51 (9) (2005) 2556–2568. doi:  
675 10.1002/aic.10502.

URL <http://dx.doi.org/10.1002/aic.10502>

- [9] E. K. Karakatsani, T. Spyriouni, I. G. Economou, Extended statistical associating fluid theory (saft) equations of state for dipolar fluids, *AICHE Journal* 51 (8) (2005) 2328–2342. doi:10.1002/aic.10473.  
URL <http://dx.doi.org/10.1002/aic.10473>
- [10] E. Karakatsani, G. Kontogeorgis, I. Economou, Evaluation of the truncated perturbed chain-polar statistical associating fluid theory for complex mixture fluid phase equilibria, *Industrial & Engineering Chemistry Research* 45 (17) (2006) 6063–6074.
- [11] E. K. Karakatsani, I. G. Economou, Perturbed chain-statistical associating fluid theory extended to dipolar and quadrupolar molecular fluids, *The Journal of Physical Chemistry B* 110 (18) (2006) 9252–9261. arXiv:<http://pubs.acs.org/doi/pdf/10.1021/jp056957b>, doi:10.1021/jp056957b.  
URL <http://pubs.acs.org/doi/abs/10.1021/jp056957b>
- [12] D. NguyenHuynh, J.-P. Passarello, P. Tobaly, J.-C. de Hemptinne, Application of gc-saft eos to polar systems using a segment approach, *Fluid Phase Equilibria* 264 (1-2) (2008) 62 – 75. doi:<http://dx.doi.org/10.1016/j.fluid.2007.10.019>.  
URL <http://www.sciencedirect.com/science/article/pii/S0378381207006838>
- [13] P. K. Jog, S. G. Sauer, J. Blaesing, W. G. Chapman, Application of dipolar chain theory to the phase behavior of polar fluids and mixtures, *Industrial & Engineering Chemistry Research* 40 (21) (2001) 4641–4648.
- [14] P. K. Jog, W. G. Chapman, Application of wertheim’s thermodynamic perturbation theory to dipolar hard sphere chains, *Molecular Physics* 97 (3) (1999) 307–319. arXiv:<http://www.tandfonline.com/doi/pdf/10.1080/00268979909482832>, doi:10.1080/00268979909482832.  
URL <http://www.tandfonline.com/doi/abs/10.1080/00268979909482832>



- [15] N. M. Al-Saifi, Prediction and computation of phase equilibria in polar and polarizable mixtures using theory-based equations of state., Ph.D. thesis, The University of British Columbia (2011).  
710
- [16] C. Perakis, E. Voutsas, K. Magoulas, D. Tassios, Thermodynamic modeling of the vapor-liquid equilibrium of the water/ethanol/co2 system, *Fluid Phase Equilibria* 243 (12) (2006) 142 – 150.  
doi:<http://dx.doi.org/10.1016/j.fluid.2006.02.018>.  
715 URL <http://www.sciencedirect.com/science/article/pii/S0378381206001208>
- [17] E. Voutsas, C. Perakis, G. Pappa, D. Tassios, An evaluation of the performance of the cubic-plus-association equation of state in mixtures of non-polar, polar and associating compounds: Towards a single model for non-polymeric systems, *Fluid Phase Equilibria* 261 (12) (2007) 343 – 350, properties and Phase Equilibria for Product and Process Design 11th International Conference on Properties and Phase Equilibria for Product and Process Design.  
720 doi:<http://dx.doi.org/10.1016/j.fluid.2007.07.051>.  
725 URL <http://www.sciencedirect.com/science/article/pii/S037838120700444X>
- [18] I. Tsvintzelis, G. M. Kontogeorgis, M. L. Michelsen, E. H. Stenby, Modeling phase equilibria for acid gas mixtures using the {CPA} equation of state. part ii: Binary mixtures with {CO<sub>2</sub>}, *Fluid Phase Equilibria* 306 (1) (2011) 38 – 56, 20 years of the {SAFT} equation of state Recent advances and challenges Symposium.  
730 doi:<http://dx.doi.org/10.1016/j.fluid.2011.02.006>.  
URL <http://www.sciencedirect.com/science/article/pii/S0378381211000641>
- [19] M. B. Oliveira, A. J. Queimada, G. M. Kontogeorgis, J. A. Coutinho, Evaluation of the {CO<sub>2</sub>} behavior in binary mixtures with alkanes,  
735

alcohols, acids and esters using the cubic-plus-association equation of state, *The Journal of Supercritical Fluids* 55 (3) (2011) 876 – 892. doi:<http://dx.doi.org/10.1016/j.supflu.2010.09.036>.

740 URL <http://www.sciencedirect.com/science/article/pii/S0896844610003268>

[20] G. M. Kontogeorgis, E. C. Voutsas, I. V. Yakoumis, D. P. Tassios, An equation of state for associating fluids, *Industrial & Engineering Chemistry Research* 35 (1996) 4310–4318. doi:10.1021/ie9600203.

745 URL <http://pubs.acs.org/doi/abs/10.1021/ie9600203>

[21] S. H. Huang, M. Radosz, Equation of state for small, large, polydisperse, and associating molecules, *Industrial & Engineering Chemistry Research* 29 (11) (1990) 2284–2294. arXiv:<http://dx.doi.org/10.1021/ie00107a014>, doi:10.1021/ie00107a014.

750 URL <http://dx.doi.org/10.1021/ie00107a014>

[22] G. M. Kontogeorgis, M. L. Michelsen, G. K. Folas, S. Derawi, N. von Solms, E. H. Stenby, Ten years with the cpa (cubic-plus-association) equation of state. part 1. pure compounds and self-associating systems, *Industrial & Engineering Chemistry Research* 45 (14) (2006) 4855–4868.

755 [23] G. M. Kontogeorgis, M. L. Michelsen, G. K. Folas, S. Derawi, N. von Solms, E. H. Stenby, Ten years with the cpa (cubic-plus-association) equation of state. part 2. cross-associating and multicomponent systems, *Industrial & Engineering Chemistry Research* 45 (14) (2006) 4869–4878. arXiv:<http://dx.doi.org/10.1021/ie051306n>, doi:10.1021/ie051306n.

760 URL <http://dx.doi.org/10.1021/ie051306n>

[24] G. Kontogeorgis, G. Folas, *Thermodynamic Models for Industrial Applications - From classical and advanced mixing rules to association theories*, 1st Edition, John Wiley and Sons, 2010, Ch. 9-12 The Cubic Plus Association Equation of State.

- 765 [25] G. Jackson, W. G. Chapman, K. E. Gubbins, Phase equilibria of associating fluids, *Molecular Physics* 65 (1) (1988) 1–31.
- [26] W. G. Chapman, G. Jackson, K. E. Gubbins, Phase equilibria of associating fluids, *Molecular Physics* 65 (5) (1988) 1057–1079.
- [27] W. G. Chapman, K. E. Gubbins, G. Jackson, M. Radosz, New reference  
770 equation of state for associating liquids, *Industrial & Engineering Chemistry Research* 29 (8) (1990) 1709–1721. arXiv:<http://dx.doi.org/10.1021/ie00104a021>, doi:10.1021/ie00104a021.  
URL <http://dx.doi.org/10.1021/ie00104a021>
- [28] M. L. Michelsen, J. M. Mollerup, *Thermodynamic Models: Fundamentals & Computational aspects*, 2nd Edition, Tie-Line Publications, 2007.  
775
- [29] G. M. Kontogeorgis, I. V. Yakoumis, H. Meijer, E. Hendriks, T. Moorwood, Multicomponent phase equilibrium calculations for water-methanol-alkane mixtures, *Fluid Phase Equilibria* 158160 (0) (1999) 201 – 209. doi:[http://dx.doi.org/10.1016/S0378-3812\(99\)00060-6](http://dx.doi.org/10.1016/S0378-3812(99)00060-6).  
780 URL <http://www.sciencedirect.com/science/article/pii/S0378381299000606>
- [30] A. de Villiers, C. Schwarz, A. Burger, G. Kontogeorgis, Evaluation of the pc-saft, {SAFT} and {CPA} equations of state in predicting derivative properties of selected non-polar and hydrogen-bonding compounds, *Fluid Phase Equilibria* 338 (0) (2013) 1 – 15.  
785 doi:<http://dx.doi.org/10.1016/j.fluid.2012.09.035>.  
URL <http://www.sciencedirect.com/science/article/pii/S0378381212004839>
- [31] J. Gross, J. Vrabec, An equation-of-state contribution for polar components: Dipolar molecules, *AIChE Journal* 52 (3) (2006) 1194–1204.  
790 doi:10.1002/aic.10683.  
URL <http://dx.doi.org/10.1002/aic.10683>

- [32] J. Wong, J. Prausnitz, Comments concerning a simple equation of state of the van der waals form, Chemical Engineering Communications 37 (1-6) (1985) 41–53. `arXiv:http://dx.doi.org/10.1080/00986448508911271`, `doi:10.1080/00986448508911271`.  
URL `http://dx.doi.org/10.1080/00986448508911271`
- [33] Dippr diadem the dippr information and data evaluation manager for the design institute for physical properties, version 8.0.2 (computer program), (2014).
- [34] M. M. E.W. Lemmon, D. Friend, NIST Chemistry WebBook, NIST Standard Reference Database Number 69, National Institute of Standards and Technology, Gaithersburg MD, 20899, `http://webbook.nist.gov`, 2014, Ch. Thermophysical Properties of Fluid Systems.  
URL `http://webbook.nist.gov`
- [35] S. Korden, N. Van Nhu, J. Vrabec, J. Gross, K. Leonhard, On the treatment of electrostatic interactions of non-spherical molecules in equation of state models, Soft Materials 10 (1-3) (2012) 81–105. `arXiv:http://dx.doi.org/10.1080/1539445X.2011.599701`, `doi:10.1080/1539445X.2011.599701`.  
URL `http://dx.doi.org/10.1080/1539445X.2011.599701`
- [36] M. G. Bjørner, G. M. Kontogeorgis, Uncertainty analysis of the cpa and a quadrupolar cpa equation of state - with emphasis on CO<sub>2</sub>, to be submitted to Fluid Phase Equilibria for publication (2015).
- [37] W. Duschek, R. Kleinrahm, W. Wagner, Measurement and correlation of the (pressure, density, temperature) relation of carbon dioxide ii. saturated-liquid and saturated-vapour densities and the vapour pressure along the entire coexistence curve, The Journal of Chemical Thermodynamics 22 (9) (1990) 841 – 864. `doi:http://dx.doi.org/10.1016/0021-9614(90)90173-N`.

URL [http://www.sciencedirect.com/science/article/pii/002196149090173N](http://www.sciencedirect.com/science/article/pii/S002196149090173N)

[38] E. Fernández-Fassnacht, F. D. Ríó, The vapour pressure of  $\text{CO}_2$  from 194 to 243 K, *The Journal of Chemical Thermodynamics* 16 (5) (1984) 469 – 474. doi:[http://dx.doi.org/10.1016/0021-9614\(84\)90204-0](http://dx.doi.org/10.1016/0021-9614(84)90204-0).

URL [http://www.sciencedirect.com/science/article/pii/0021961484902040](http://www.sciencedirect.com/science/article/pii/S0021961484902040)

[39] J. R. Roebuck, T. A. Murrell, E. E. Miller, The joule-thomson effect in carbon dioxide, *Journal of the American Chemical Society* 64 (2) (1942) 400–411. arXiv:<http://dx.doi.org/10.1021/ja01254a048>, doi:10.1021/ja01254a048.

URL <http://dx.doi.org/10.1021/ja01254a048>

[40] C. Gray, K. Gubbins, *Theory of Molecular Fluids: I: Fundamentals*, International Series of Monographs on Chemistry, OUP Oxford, 1984.

URL <http://books.google.dk/books?id=3mz2RcnnMGwC>

[41] A. D. Buckingham, Molecular quadrupole moments, *Q. Rev. Chem. Soc.* 13 (1959) 183–214. doi:10.1039/QR9591300183.

URL <http://dx.doi.org/10.1039/QR9591300183>

[42] A. D. Buckingham, R. L. Disch, The quadrupole moment of the carbon dioxide molecule, *Proceedings of the Royal Society of London A: Mathematical, Physical and Engineering Sciences* 273 (1353) (1963) 275–289. doi:10.1098/rspa.1963.0088.

[43] A. D. Buckingham, R. L. Disch, D. A. Dunmur, Quadrupole moments of some simple molecules, *Journal of the American Chemical Society* 90 (12) (1968) 3104–3107. arXiv:<http://dx.doi.org/10.1021/ja01014a023>, doi:10.1021/ja01014a023.

URL <http://dx.doi.org/10.1021/ja01014a023>

- [44] D. E. Stogryn, A. P. Stogryn, Molecular multipole moments, *Molecular Physics* 11 (4) (1966) 371–393. [arXiv:http://dx.doi.org/10.1080/00268976600101201](https://arxiv.org/abs/http://dx.doi.org/10.1080/00268976600101201), doi:10.1080/00268976600101201.  
850 URL <http://dx.doi.org/10.1080/00268976600101201>
- [45] J. E. Harries, The quadrupole moment of  $\text{CO}_2$ , measured from the far infrared spectrum, *Journal of Physics B: Atomic and Molecular Physics* 3 (12) (1970) L150.  
855 URL <http://stacks.iop.org/0022-3700/3/i=12/a=021>
- [46] C. Graham, J. Pierrus, R. Raab, Measurement of the electric quadrupole moments of  $\text{CO}_2$ ,  $\text{CO}$  and  $\text{N}_2$ , *Molecular Physics* 67 (4) (1989) 939–955. [arXiv:http://dx.doi.org/10.1080/00268978900101551](https://arxiv.org/abs/http://dx.doi.org/10.1080/00268978900101551), doi:10.1080/00268978900101551.  
860 URL <http://dx.doi.org/10.1080/00268978900101551>
- [47] J. Zeuthen, Extension of the CPA EOS to systems containing organic acids, Masters thesis, The Technical University of Denmark (2003).
- [48] B. Mallu, G. Natarajan, D. Viswanath,  $(p, v_m, t, x)$  and virial coefficients of  $x\text{CO}_2 + (1-x)\text{CO}$ , *The Journal of Chemical Thermodynamics* 21 (9) (1989) 989 – 996. doi:[http://dx.doi.org/10.1016/0021-9614\(89\)90159-6](http://dx.doi.org/10.1016/0021-9614(89)90159-6).  
865 URL <http://www.sciencedirect.com/science/article/pii/S0021961489901596>
- [49] R. S. Dadson, E. J. Evans, J. H. King, The second virial coefficient of carbon dioxide, *Proceedings of the Physical Society* 92 (4) (1967) 1115.  
870 URL <http://stacks.iop.org/0370-1328/92/i=4/a=337>
- [50] J. Dymond, E. B. Smith, *The virial coefficients of pure gases and mixtures: a critical compilation*, Oxford University Press, 1980.
- [51] E. G. Butcher, R. S. Dadson, The virial coefficients of the carbon dioxide-ethylene system. i. pure gases, *Proceedings of the Royal Society of London*  
875

A: Mathematical, Physical and Engineering Sciences 277 (1371) (1964)  
448–467. doi:10.1098/rspa.1964.0034.

[52] T. L. Cottrell, R. A. Hamilton, The second virial coefficient of gases and  
mixtures. part 1.-carbon dioxide + helium mixtures, Trans. Faraday Soc.  
880 52 (1956) 156–160. doi:10.1039/TF9565200156.

URL <http://dx.doi.org/10.1039/TF9565200156>

[53] P. J. McElroy, L. L. Kee, C. A. Renner, Excess second virial coeffi-  
cients for binary mixtures of carbon dioxide with methane, ethane, and  
propane, Journal of Chemical & Engineering Data 35 (3) (1990) 314–  
885 317. arXiv:<http://dx.doi.org/10.1021/je00061a024>, doi:10.1021/  
je00061a024.

URL <http://dx.doi.org/10.1021/je00061a024>

[54] B. Schramm, E. Elias, L. Kern, G. Natour, A. Schmitt, C. Weber,  
Precise measurements of second virial coefficients of simple gases and  
890 gas mixtures in the temperature range below 300 k, Berichte der Bun-  
sengesellschaft fr physikalische Chemie 95 (5) (1991) 615–621. doi:  
10.1002/bbpc.19910950513.

URL <http://dx.doi.org/10.1002/bbpc.19910950513>

[55] R. Span, W. Wagner, A new equation of state for carbon dioxide covering  
895 the fluid region from the triple-point temperature to 1100 k at pressures  
up to 800 mpa, Journal of Physical and Chemical Reference Data 25 (6)  
(1996) 1509–1596. doi:<http://dx.doi.org/10.1063/1.555991>.

URL [http://scitation.aip.org/content/aip/journal/jpcrd/25/6/  
10.1063/1.555991](http://scitation.aip.org/content/aip/journal/jpcrd/25/6/10.1063/1.555991)

[56] E. Hendriks, G. M. Kontogeorgis, R. Dohrn, J.-C. de Hemptinne, I. G.  
900 Economou, L. F. ilnik, V. Vesovic, Industrial requirements for thermo-  
dynamics and transport properties, Industrial & Engineering Chemistry  
Research 49 (22) (2010) 11131–11141. arXiv:<http://dx.doi.org/10.1021/ie100877a011>.

1021/ie101231b, doi:10.1021/ie101231b.

905 URL <http://dx.doi.org/10.1021/ie101231b>

[57] T. Lafitte, D. Bessieres, M. M. Pieiro, J.-L. Daridon, Simultaneous estimation of phase behavior and second-derivative properties using the statistical associating fluid theory with variable range approach, *The Journal of Chemical Physics* 124 (2) (2006) -. doi:<http://dx.doi.org/10.1063/1.2140276>.

910 URL <http://scitation.aip.org/content/aip/journal/jcp/124/2/10.1063/1.2140276>

[58] T. Lafitte, M. M. Piñeiro, J.-L. Daridon, D. Bessières, A comprehensive description of chemical association effects on second derivative properties of alcohols through a soft-vr approach, *The Journal of Physical Chemistry B* 111 (13) (2007) 3447–3461, pMID: 17388508. arXiv:<http://dx.doi.org/10.1021/jp0682208>, doi:10.1021/jp0682208.

915 URL <http://dx.doi.org/10.1021/jp0682208>

[59] X. Liang, B. Maribo-Mogensen, K. Thomsen, W. Yan, G. M. Kontogeorgis, Approach to improve speed of sound calculation within pcsoft framework, *Industrial & Engineering Chemistry Research* 51 (45) (2012) 14903–14914. arXiv:<http://dx.doi.org/10.1021/ie3018127>, doi:10.1021/ie3018127.

920 URL <http://dx.doi.org/10.1021/ie3018127>

[60] P. G. Brewer, G. Friederich, E. T. Peltzer, F. M. Orr, Direct experiments on the ocean disposal of fossil fuel co<sub>2</sub>, *Science* 284 (5416) (1999) 943–945. arXiv:<http://www.sciencemag.org/content/284/5416/943.full.pdf>, doi:10.1126/science.284.5416.943.

925 URL <http://www.sciencemag.org/content/284/5416/943.abstract>

[61] L. A. Webster, A. J. Kidnay, Vapor-liquid equilibria for the methane-propane-carbon dioxide systems at 230 k and 270 k, *Journal of Chemical & Engineering Data* 46 (3) (2001) 759–764. arXiv:<http://dx.doi.org/>

930



10.1021/je000307d, doi:10.1021/je000307d.

URL <http://dx.doi.org/10.1021/je000307d>

935 [62] M. S.-W. Wei, T. S. Brown, A. J. Kidnay, E. D. Sloan, Vapor + liquid equilibria for the ternary system methane + ethane + carbon dioxide at 230 k and its constituent binaries at temperatures from 207 to 270 k, *Journal of Chemical & Engineering Data* 40 (4) (1995) 726–731. arXiv:<http://dx.doi.org/10.1021/je00020a002>, doi:10.1021/je00020a002.

940 URL <http://dx.doi.org/10.1021/je00020a002>

[63] T. Brown, A. Kidnay, E. Sloan, Vapor-liquid equilibria in the carbon dioxide-ethane system, *Fluid Phase Equilibria* 40 (12) (1988) 169 – 184. doi:[http://dx.doi.org/10.1016/0378-3812\(88\)80028-1](http://dx.doi.org/10.1016/0378-3812(88)80028-1).

945 URL <http://www.sciencedirect.com/science/article/pii/S0378381288800281>

[64] A. Q. Clark, K. Stead, (vapour + liquid) phase equilibria of binary, ternary, and quaternary mixtures of ch<sub>4</sub>, c<sub>2</sub>h<sub>6</sub>, c<sub>3</sub>h<sub>8</sub>, c<sub>4</sub>h<sub>10</sub>, and {CO<sub>2</sub>}, *The Journal of Chemical Thermodynamics* 20 (4) (1988) 413 – 427. doi:[http://dx.doi.org/10.1016/0021-9614\(88\)90178-4](http://dx.doi.org/10.1016/0021-9614(88)90178-4).

950 URL <http://www.sciencedirect.com/science/article/pii/S0021961488901784>

[65] H. H. Reamer, B. H. Sage, Phase equilibria in hydrocarbon systems. volumetric and phase behavior of the n-decane-co<sub>2</sub> system., *Journal of Chemical & Engineering Data* 8 (4) (1963) 508–513. arXiv:<http://dx.doi.org/10.1021/je60019a010>, doi:10.1021/je60019a010.

955 URL <http://dx.doi.org/10.1021/je60019a010>

[66] J. J. C. Hsu, N. Nagarajan, R. L. Robinson, Equilibrium phase compositions, phase densities, and interfacial tensions for carbon dioxide + hydrocarbon systems. 1. carbon dioxide + n-butane, *Journal of Chemical & Engineering Data* 30 (4) (1985) 485–491. arXiv:<http://dx.doi.org/>

960

10.1021/je00042a036, doi:10.1021/je00042a036.

URL <http://dx.doi.org/10.1021/je00042a036>

[67] M. E. Pozo de Fernandez, J. A. Zollweg, W. B. Streett, Vapor-liquid equilibrium in the binary system carbon dioxide + n-butane, Journal of Chemical & Engineering Data 34 (3) (1989) 324–328. arXiv:<http://dx.doi.org/10.1021/je00057a019>, doi:10.1021/je00057a019.

URL <http://dx.doi.org/10.1021/je00057a019>

[68] T. Brown, V. Niesen, E. Sloan, A. Kidnay, Vapor-liquid equilibria for the binary systems of nitrogen, carbon dioxide, and n-butane at temperatures from 220 to 344 k., Fluid Phase Equilibria 53 (0) (1989) 7 – 14, proceedings of the Fifth International Conference. doi:[http://dx.doi.org/10.1016/0378-3812\(89\)80067-6](http://dx.doi.org/10.1016/0378-3812(89)80067-6).

URL <http://www.sciencedirect.com/science/article/pii/0378381289800676>

[69] H. Cheng, M. E. Pozo de Fernandez, J. A. Zollweg, W. B. Streett, Vapor-liquid equilibrium in the system carbon dioxide + n-pentane from 252 to 458 k at pressures to 10 mpa, Journal of Chemical & Engineering Data 34 (3) (1989) 319–323. arXiv:<http://dx.doi.org/10.1021/je00057a018>, doi:10.1021/je00057a018.

URL <http://dx.doi.org/10.1021/je00057a018>

[70] G.-I. Kaminishi, C. Yokoyama, T. Shinji, Vapor pressures of binary mixtures of carbon dioxide with benzene, n-hexane and cyclohexane up to 7 {MPa}, Fluid Phase Equilibria 34 (1) (1987) 83 – 99. doi:[http://dx.doi.org/10.1016/0378-3812\(87\)85052-5](http://dx.doi.org/10.1016/0378-3812(87)85052-5).

URL <http://www.sciencedirect.com/science/article/pii/0378381287850525>

[71] D. Nguyen-Huynh, J.-P. Passarello, P. Tobaly, J.-C. de Hemptinne, Modeling phase equilibria of asymmetric mixtures using a group-contribution saft (gc-saft) with a kij correlation method based on londons theory. 1.

- 990 application to  $\text{CO}_2$  + n-alkane, methane + n-alkane, and ethane + n-alkane  
systems, *Industrial & Engineering Chemistry Research* 47 (22) (2008)  
8847–8858. arXiv:<http://dx.doi.org/10.1021/ie071643r>, doi:10.  
1021/ie071643r.  
URL <http://dx.doi.org/10.1021/ie071643r>
- 995 [72] J. D. Hottovy, K. D. Luks, J. P. Kohn, Three-phase liquid-liquid-vapor  
equilibria behavior of certain binary carbon dioxide-n-paraffin sys-  
tems, *Journal of Chemical & Engineering Data* 26 (3) (1981) 256–  
258. arXiv:<http://dx.doi.org/10.1021/je00025a009>, doi:10.1021/  
je00025a009.  
1000 URL <http://dx.doi.org/10.1021/je00025a009>
- [73] J. Reilly, C. Bokis, M. Donohue, An experimental investigation of lewis  
acid-base interactions of liquid carbon dioxide using fourier transform in-  
frared (ft-ir) spectroscopy, *International Journal of Thermophysics* 16 (3)  
(1995) 599–610. doi:10.1007/BF01438845.  
1005 URL <http://dx.doi.org/10.1007/BF01438845>
- [74] P. Lalanne, T. Tassaing, Y. Danten, F. Cansell, S. C. Tucker, M. Besnard,  
 $\text{CO}_2$ -ethanol interaction studied by vibrational spectroscopy in supercrit-  
ical  $\text{CO}_2$ , *The Journal of Physical Chemistry A* 108 (14) (2004) 2617–  
2624. arXiv:<http://dx.doi.org/10.1021/jp037802b>, doi:10.1021/  
1010 jp037802b.  
URL <http://dx.doi.org/10.1021/jp037802b>
- [75] Y. Danten, T. Tassaing, M. Besnard, Ab initio investigation of vibrational  
spectra of  $\text{water}(\text{CO}_2)_n$  complexes ( $n = 1, 2$ ), *The Journal of Physical  
Chemistry A* 109 (14) (2005) 3250–3256, PMID: 16833656. arXiv:<http://dx.doi.org/10.1021/jp0503819>,  
1015 //dx.doi.org/10.1021/jp0503819, doi:10.1021/jp0503819.  
URL <http://dx.doi.org/10.1021/jp0503819>
- [76] J. Sadlej, P. Mazurek, Ab initio calculations on the water-carbon dioxide  
system, *Journal of Molecular Structure: {THEOCHEM}* 337 (2) (1995)

- 129 – 138. doi:[http://dx.doi.org/10.1016/0166-1280\(95\)04132-P](http://dx.doi.org/10.1016/0166-1280(95)04132-P).  
1020 URL <http://www.sciencedirect.com/science/article/pii/S016612809504132P>
- [77] M. Kleiner, G. Sadowski, Modeling of polar systems using pcp-saft: An approach to account for induced-association interactions, *The Journal of Physical Chemistry C* 111 (43) (2007) 15544–15553. arXiv:<http://dx.doi.org/10.1021/jp072640v>, doi:10.1021/jp072640v.  
1025 URL <http://dx.doi.org/10.1021/jp072640v>
- [78] D. NguyenHuynh, J.-P. Passarello, J.-C. de Hemptinne, F. Volle, P. Tobaly, Simultaneous modeling of vle, {LLE} and {VLLE} of {CO<sub>2</sub>} and 1, 2, 3 and 4 alkanol containing mixtures using gc-ppc-saft {EOS}, *The Journal of Supercritical Fluids* 95 (0) (2014) 146 – 157.  
1030 doi:<http://dx.doi.org/10.1016/j.supflu.2014.07.022>.  
URL <http://www.sciencedirect.com/science/article/pii/S089684461400223X>
- [79] K. Suzuki, H. Sue, M. Itou, R. L. Smith, H. Inomata, K. Arai, S. Saito,  
1035 Isothermal vapor-liquid equilibrium data for binary systems at high pressures: carbon dioxide-methanol, carbon dioxide-ethanol, carbon dioxide-1-propanol, methane-ethanol, methane-1-propanol, ethane-ethanol, and ethane-1-propanol systems, *Journal of Chemical & Engineering Data* 35 (1) (1990) 63–66. arXiv:<http://dx.doi.org/10.1021/je00059a020>,  
1040 doi:10.1021/je00059a020.  
URL <http://dx.doi.org/10.1021/je00059a020>
- [80] J. H. Yoon, M. K. Chun, W. H. Hong, H. Lee, High-pressure phase equilibria for carbon dioxide-methanol-water system: experimental data and critical evaluation of mixing rules, *Industrial & Engineering Chemistry Research* 32 (11) (1993) 2881–2887. arXiv:<http://dx.doi.org/10.1021/ie00023a061>, doi:10.1021/ie00023a061.  
1045 URL <http://dx.doi.org/10.1021/ie00023a061>

- 1050 [81] C. J. Chang, K.-L. Chiu, C.-Y. Day, A new apparatus for the determination of p-x-y diagrams and henry's constants in high pressure alcohols with critical carbon dioxide, *The Journal of Supercritical Fluids* 12 (3) (1998) 223 – 237. doi:[http://dx.doi.org/10.1016/S0896-8446\(98\)00076-X](http://dx.doi.org/10.1016/S0896-8446(98)00076-X).  
URL <http://www.sciencedirect.com/science/article/pii/S089684469800076X>
- 1055 [82] J. Briones, J. Mullins, M. Thies, B.-U. Kim, Ternary phase equilibria for acetic acid-water mixtures with supercritical carbon dioxide, *Fluid Phase Equilibria* 36 (0) (1987) 235 – 246. doi:[http://dx.doi.org/10.1016/0378-3812\(87\)85026-4](http://dx.doi.org/10.1016/0378-3812(87)85026-4).  
URL <http://www.sciencedirect.com/science/article/pii/0378381287850264>
- 1060 [83] A. Bamberger, G. Sieder, G. Maurer, High-pressure (vapor+liquid) equilibrium in binary mixtures of (carbon dioxide+water or acetic acid) at temperatures from 313 to 353 k, *The Journal of Supercritical Fluids* 17 (2) (2000) 97 – 110. doi:[http://dx.doi.org/10.1016/S0896-8446\(99\)00054-6](http://dx.doi.org/10.1016/S0896-8446(99)00054-6).  
1065 URL <http://www.sciencedirect.com/science/article/pii/S0896844699000546>
- [84] A. D. King, C. R. Coan, Solubility of water in compressed carbon dioxide, nitrous oxide, and ethane. evidence for hydration of carbon dioxide and nitrous oxide in the gas phase, *Journal of the American Chemical Society* 93 (8) (1971) 1857–1862. arXiv:<http://dx.doi.org/10.1021/ja00737a004>, doi:10.1021/ja00737a004.  
1070 URL <http://dx.doi.org/10.1021/ja00737a004>
- [85] R. Dohrn, A. Bünz, F. Devlieghere, D. Thelen, Experimental measurements of phase equilibria for ternary and quaternary systems of glucose, water, co<sub>2</sub> and ethanol with a novel apparatus, *Fluid Phase Equilibria* 83 (1993) 149–158.  
1075

- [86] K. Ohgaki, T. Katayama, Isothermal vapor-liquid equilibrium data for binary systems containing carbon dioxide at high pressures: methanol-carbon dioxide, n-hexane-carbon dioxide, and benzene-carbon dioxide systems, *Journal of Chemical & Engineering Data* 21 (1) (1976) 53–55. arXiv:<http://dx.doi.org/10.1021/je60068a015>, doi:10.1021/je60068a015.  
URL <http://dx.doi.org/10.1021/je60068a015>
- [87] J. H. Hong, R. Kobayashi, Vapor-liquid equilibrium studies for the carbon dioxide-methanol system, *Fluid Phase Equilibria* 41 (3) (1988) 269 – 276. doi:[http://dx.doi.org/10.1016/0378-3812\(88\)80011-6](http://dx.doi.org/10.1016/0378-3812(88)80011-6).  
URL <http://www.sciencedirect.com/science/article/pii/S0378381288800116>
- [88] K. Bezanhtak, G. B. Combes, F. Dehghani, N. R. Foster, D. L. Tomasko, Vapor-liquid equilibrium for binary systems of carbon dioxide + methanol, hydrogen + methanol, and hydrogen + carbon dioxide at high pressures, *Journal of Chemical & Engineering Data* 47 (2) (2002) 161–168. arXiv:<http://dx.doi.org/10.1021/je010122m>, doi:10.1021/je010122m.  
URL <http://dx.doi.org/10.1021/je010122m>
- [89] D. Kodama, M. Kato, High-pressure phase equilibrium for carbon dioxide + ethanol at 291.15 k, *Journal of Chemical & Engineering Data* 50 (1) (2005) 16–17. arXiv:<http://dx.doi.org/10.1021/je034243t>, doi:10.1021/je034243t.  
URL <http://dx.doi.org/10.1021/je034243t>
- [90] O. Pfohl, A. Pagel, G. Brunner, Phase equilibria in systems containing o-cresol, p-cresol, carbon dioxide, and ethanol at 323.15–473.15 k and 10–35 {MPa}, *Fluid Phase Equilibria* 157 (1) (1999) 53 – 79. doi:[http://dx.doi.org/10.1016/S0378-3812\(99\)00019-9](http://dx.doi.org/10.1016/S0378-3812(99)00019-9).  
URL <http://www.sciencedirect.com/science/article/pii/S0378381299000199>

- [91] C. A. Perakis, E. C. Voutsas, K. G. Magoulas, D. P. Tassios, Thermodynamic modeling of the water + acetic acid + co<sub>2</sub> system: The importance of the number of association sites of water and of the nonassociation contribution for the cpa and saft-type models, *Industrial & Engineering Chemistry Research* 46 (3) (2007) 932–938. `arXiv:http://dx.doi.org/10.1021/ie0609416`, `doi:10.1021/ie0609416`.  
URL `http://dx.doi.org/10.1021/ie0609416`
- [92] S. Kim, Y. Kim, B. H. Park, J. H. Lee, J. W. Kang, Measurement and correlation of solubility of water in carbon dioxide-rich phase, *Fluid Phase Equilibria* 328 (0) (2012) 9 – 12. `doi:http://dx.doi.org/10.1016/j.fluid.2012.05.013`.  
URL `http://www.sciencedirect.com/science/article/pii/S0378381212002117`
- [93] Z.-Y. Zhang, J.-C. Yang, Y.-G. Li, Prediction of phase equilibria for co<sub>2</sub>-c<sub>2</sub>h<sub>5</sub>oh-h<sub>2</sub>o system using the {SAFT} equation of state, *Fluid Phase Equilibria* 169 (1) (2000) 1 – 18. `doi:http://dx.doi.org/10.1016/S0378-3812(99)00341-6`.  
URL `http://www.sciencedirect.com/science/article/pii/S0378381299003416`
- [94] X. Ji, S. P. Tan, H. Adidharma, M. Radosz, Saft1-rpm approximation extended to phase equilibria and densities of co<sub>2</sub>-h<sub>2</sub>o and co<sub>2</sub>-h<sub>2</sub>o-nacl systems, *Industrial & Engineering Chemistry Research* 44 (22) (2005) 8419–8427. `arXiv:http://dx.doi.org/10.1021/ie050725h`, `doi:10.1021/ie050725h`.  
URL `http://dx.doi.org/10.1021/ie050725h`
- [95] M. C. dos Ramos, F. J. Blas, A. Galindo, Modelling the phase equilibria and excess properties of the water-carbon dioxide binary mixture, *Fluid Phase Equilibria* 261 (12) (2007) 359 – 365, properties and Phase Equilibria for Product and Process Design 11th International Conference

- 1135 on Properties and Phase Equilibria for Product and Process Design.  
doi:<http://dx.doi.org/10.1016/j.fluid.2007.07.012>.  
URL <http://www.sciencedirect.com/science/article/pii/S0378381207003664>
- [96] X. Tang, J. Gross, Modeling the phase equilibria of hydrogen sulfide  
1140 and carbon dioxide in mixture with hydrocarbons and water using the  
pcp-saft equation of state, Fluid Phase Equilibria 293 (1) (2010) 11 – 21.  
doi:<http://dx.doi.org/10.1016/j.fluid.2010.02.004>.  
URL <http://www.sciencedirect.com/science/article/pii/S0378381210000798>
- 1145 [97] N. I. Diamantonis, I. G. Economou, Modeling the phase equilibria of a  
h<sub>2</sub>o-co<sub>2</sub> mixture with pc-saft and tpc-psaft equations of state, Molecular  
Physics 110 (11-12) (2012) 1205–1212. arXiv:<http://dx.doi.org/10.1080/00268976.2012.656721>, doi:10.1080/00268976.2012.656721.  
URL <http://dx.doi.org/10.1080/00268976.2012.656721>
- 1150 [98] D. Nguyen-Huynh, J.-C. de Hemptinne, R. Lugo, J.-P. Passarello, P. To-  
baly, Modeling liquid-liquid and liquid-vapor equilibria of binary systems  
containing water with an alkane, an aromatic hydrocarbon, an alcohol  
or a gas (methane, ethane, co<sub>2</sub> or h<sub>2</sub>s), using group contribution po-  
lar perturbed-chain statistical associating fluid theory, Industrial & En-  
1155 gineering Chemistry Research 50 (12) (2011) 7467–7483. arXiv:<http://dx.doi.org/10.1021/ie102045g>, doi:10.1021/ie102045g.  
URL <http://dx.doi.org/10.1021/ie102045g>
- [99] B. Rumpf, H. Nicolaisen, C. cal, G. Maurer, Solubility of carbon diox-  
ide in aqueous solutions of sodium chloride: Experimental results and  
1160 correlation, Journal of Solution Chemistry 23 (3) (1994) 431–448. doi:  
10.1007/BF00973113.  
URL <http://dx.doi.org/10.1007/BF00973113>



- [100] P. G. Bendale, R. M. Enick, Use of carbon dioxide to shift benzene/acetonitrile and benzene/cyclohexane azeotropes, Fluid Phase Equilibria 94 (0) (1994) 227 – 253. doi:[http://dx.doi.org/10.1016/S0378-3812\(94\)87059-4](http://dx.doi.org/10.1016/S0378-3812(94)87059-4).  
URL <http://www.sciencedirect.com/science/article/pii/S0378381294870594>
- [101] S. Dougill, W. Sajik, S. Hwang, Preparing a liquified mixture of carbon dioxide and acetylene by mixing the pressurized gases and cooling to obtain a liquid/gas or solid/gas mixture; nonexplosive, uS Patent 5,766,514 (Jun. 16 1998).  
URL <http://www.google.com/patents/US5766514>
- [102] C. G. Gray, K. E. Gubbins, C. G. Joslin, Theory of Molecular Fluids Volume 2: Applications, Vol. 53, Oxford University Press, 2011. arXiv:<http://dx.doi.org/10.1080/00107514.2012.720286>, doi:10.1080/00107514.2012.720286.  
URL <http://dx.doi.org/10.1080/00107514.2012.720286>
- [103] I. Tsivintzelis, G. M. Kontogeorgis, M. L. Michelsen, E. H. Stenby, Modeling phase equilibria for acid gas mixtures using the cpa equation of state. i. mixtures with h<sub>2</sub>s, AIChE Journal 56 (11) (2010) 2965–2982. doi:10.1002/aic.12207.  
URL <http://dx.doi.org/10.1002/aic.12207>
- [104] I. V. Yakoumis, G. M. Kontogeorgis, E. C. Voutsas, D. P. Tassios, Vapor-liquid equilibria for alcoholhydrocarbon systems using the {CPA} equation of state, Fluid Phase Equilibria 130 (12) (1997) 31 – 47. doi:[http://dx.doi.org/10.1016/S0378-3812\(96\)03200-1](http://dx.doi.org/10.1016/S0378-3812(96)03200-1).  
URL <http://www.sciencedirect.com/science/article/pii/S0378381296032001>
- [105] I. Tsivintzelis, B. Maribo-Mogensen, The cubic-plus-association eos, Tech. rep., CERE, Center for Energy Resources Engineering Department of

Chemical and Biochemical Engineering, Technical University of Denmark  
(September 2012).

- [106] G. Folas, Modeling of complex mixtures containing hydrogen bonding molecules, Ph.D. thesis (2007).

1195