

Technical University of Denmark



Feeding chicory (*Cichorium intybus*) selectively reduces *Ostertagia ostertagi* infection levels in cattle

Pena-Espinoza, Miguel Angel; Desrues, Oliver; Hansen, Tina; Thamsborg, Stig M.; Enemark, Heidi

Published in:

25th International Conference of the World Association for the Advancement of Veterinary Parasitology

Publication date:

2015

Document Version

Publisher's PDF, also known as Version of record

[Link back to DTU Orbit](#)

Citation (APA):

Pena-Espinoza, M. A., Desrues, O., Hansen, T., Thamsborg, S. M., & Enemark, H. (2015). Feeding chicory (*Cichorium intybus*) selectively reduces *Ostertagia ostertagi* infection levels in cattle. In 25th International Conference of the World Association for the Advancement of Veterinary Parasitology: Abstract Book (pp. 299-299). Liverpool, United Kingdom.

DTU Library

Technical Information Center of Denmark

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Oral poster session 1: We can work it out

Tuesday 18 August
10:00-10:30

Ref: 0642

Poster Number: P2C328

Feeding chicory (*Cichorium intybus*) selectively reduces *Ostertagia ostertagi* infection levels in cattle

Miguel Peña-Espinoza^{1,2}, Olivier Desrues², Tina Hansen², Stig Thamsborg², Heidi Enemark^{1,3}

¹National Veterinary Institute, Technical University of Denmark, Copenhagen, Denmark,

²Department of Veterinary Disease Biology, University of Copenhagen, Copenhagen, Denmark,

³Norwegian Veterinary Institute, Oslo, Norway

Objectives: Studies were conducted to test the potential use of chicory against gastrointestinal nematode infections in cattle. **Methods:** In study 1, fifteen 2-4 months-old dairy calves were allocated into a chicory (CHI, n=9) or control (CTL, n=6) group. CHI and CTL were stabled and fed with chicory silage or hay, resp., *ad lib* for 56 days. Protein/energy intakes were equalized between groups throughout the study. After 14 days on the diet all calves were infected with 10,000 *Ostertagia ostertagi* and 66,000 *Cooperia oncophora* third-stage (L3) larvae. In study 2, twenty 4-6 months-old dairy calves grazed a second-year, pure chicory sward (CHI, n=10) or a ryegrass/white clover pasture (CTL, n=10) for 43 days. After 7 days on the diet all calves were infected with 20,000 *O. ostertagi* L3. In both studies, individual live weights were recorded and faecal egg counts were calculated as number of eggs per g of dried feces (FECDM). At day 56 (study 1) calves were killed for worm recovery. Live weights and log-transformed FECDM were analysed by ANOVA using repeated measurements. Log-transformed worm counts were analysed by t-test. **Results:** In study 1 daily live weight gains were 500 and 329 g/day in CHI and CTL animals, resp. (p=0.02). Mean FECDM were not significantly different between groups (p=0.19). *O. ostertagi* geo mean worm counts (SD) were 1599 (886) and 3752 (633) in CHI and CTL groups, resp. (p<0.001). *C. oncophora* geo mean worm counts were not statistically different between groups (p=0.10). In study 2 daily live weight gains were 366 and 748 g/day in CHI and CTL calves, resp. (p<0.001). Mean FECDM of CHI and CTL at day 20 post-infection were 1589 and 1474, resp. (p>0.05). From this point, egg excretion in CHI calves was significantly reduced and by day 36 post-infection FECDM was decreased by 48-65% compared to CTL (P<0.05). **Discussion:** Feeding on a chicory diet demonstrated a marked anthelmintic effect against *O. ostertagi* in both trials, whereas *C. oncophora* in study 1 was unaffected. Adult worm counts and egg excretion of *O. ostertagi* were significantly reduced in chicory-fed calves, suggesting that the survival of adult worms is reduced and/or worm expulsion is enhanced. The lower weight gains of CHI in study 2 probably reflect lower energy consumption and suggest that duration of grazing of pure chicory should be limited to selectively target established *O. ostertagi* adult populations.