

Technical University of Denmark



Continuum Navier-Stokes modelling of water ow past fullerene molecules

Walther, Jens Honore; Popadic, A. ; Koumoutsakos, P.; Praprotnik, M.

Publication date:
2015

Document Version
Peer reviewed version

[Link back to DTU Orbit](#)

Citation (APA):

Walther, J. H., Popadic, A., Koumoutsakos, P., & Praprotnik, M. (2015). Continuum Navier-Stokes modelling of water ow past fullerene molecules. Abstract from 68th Annual Meeting of the American Physical Society's Division of Fluid Dynamics (DFD) , Boston, United States.

DTU Library

Technical Information Center of Denmark

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Abstract Submitted
for the DFD15 Meeting of
The American Physical Society

Continuum Navier-Stokes modelling of water flow past fullerene molecules J. H. WALTHER, Technical University of Denmark, Denmark, A. POPADIC, National Institute of Chemistry, Slovenia, P. KOUMOUTSAKOS, ETH Zurich, Switzerland, M. PRAPROTNIK, National Institute of Chemistry, Slovenia — We present continuum simulations of water flow past fullerene molecules. The governing Navier-Stokes equations are complemented with the Navier slip boundary condition with a slip length that is extracted from related molecular dynamics simulations. We find that several quantities of interest as computed by the present model are in good agreement with results from atomistic and atomistic-continuum simulations at a fraction of the computational cost. We simulate the flow past a single fullerene and an array of fullerenes and demonstrate that such nanoscale flows can be computed efficiently by continuum flow solvers, allowing for investigations into spatiotemporal scales inaccessible to atomistic simulations.

J. H. Walther
Technical University of Denmark, Denmark

Date submitted: 29 Jul 2015

Electronic form version 1.4