

Technical University of Denmark



Azido-RGO as a universal platform for RGO based sensors

Olsen, Gunnar; Ulstrup, Jens; Chi, Qijin

Publication date:
2015

Document Version
Publisher's PDF, also known as Version of record

[Link back to DTU Orbit](#)

Citation (APA):

Olsen, G., Ulstrup, J., & Chi, Q. (2015). Azido-RGO as a universal platform for RGO based sensors. Abstract from DTU Chemistry PhD School Symposium 2015, Kgs. Lyngby, Denmark.

DTU Library

Technical Information Center of Denmark

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Azido-RGO as a universal platform for RGO based sensors


Gunnar Olsen, Jens Ulstrup and Qijin Chi

Technical University of Denmark, Department of chemistry, 2800 Kgs Lyngby, Denmark



Reduced graphene oxide (RGO) is an interesting candidate for support material in chemical sensors [1]. RGO exhibits sufficient electric conductivity; high surface area and can be functionalized [2] by multifarious functional surface groups. This is attractive in selective introduction of enzymes or supramolecular host complexes for bio- and chemosensing, respectively [3-4]. Functionalization is mostly effected at the graphene oxide (GO) step during processing of RGO, However, these systems can be unstable under normal reduction conditions. To address this issue several mild “green” reduction methods of GO can be used, but these methods lead to insufficient reduction of GO which in turn results in diminished conductivity [5].

In our work we have instead focused our attention on synthesis of azide functionalized RGO. This material offers highly reduced RGO with possibility of post-reduction functionalization through Copper(I)-catalyzed Azide-Alkyne Cycloaddition (CuAAC) “click-chemistry” tested on numerous complex and fragile systems [6].



Scheme. 1. Azido-RGO showing function Azide groups on both surface and edges of RGO-sheet.

Acknowledgments

We are grateful for financial support by the Danish Research Council for Technology and Product Science (Project No. 12-127447) and by the IRSES FP7 Foreign exchange Program ELECTRONANOMAT .

References

- [1] S. Sharma and B. G. Pollet, *J. Power Sources*, **208**, 96-119 (2012)
- [2] Y. Zhu, *et al.*, *Adv. Mater.*, **22**, 3906-3924 (2010)
- [3] S. Wu, *et al.*, *Small*, **9**, 1160-1172 (2013)
- [4] L. You, *et al.*, *Chem. Rev.* **115**, 7840-7892 [2015]
- [5] S. Thakur; N. Karak, *Carbon*, **94**, 224-242, (2015)
- [6] M. Meldel; C. W. Tornøe, *Chem. Rev.* **108**, 2952-3015 (2008)