brought to you by I CORE

Technical University of Denmark



Speciation analysis of 129I in atmosphere by AMS and its applications for studies of environmental processes

Hou, Xiaolin; Zhang, Luyuan; Xu, Sheng; Zhou, Weijian

Published in:

Final Program of the 6th East Asia Accelerator Mass Spectrometry Symposium (EA-AMS 6)

Publication date: 2015

Document Version Publisher's PDF, also known as Version of record

Link back to DTU Orbit

Citation (APA):

Hou, X., Zhang, L., Xu, S., & Zhou, W. (2015). Speciation analysis of 129I in atmosphere by AMS and its applications for studies of environmental processes. In Final Program of the 6th East Asia Accelerator Mass Spectrometry Symposium (EA-AMS 6) (pp. 52-53). National Taiwan University.

DTU Library

Technical Information Center of Denmark

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Speciation analysis of 129I in atmosphere by AMS and its applications for studies of environmental processes

Xiaolin Hou1,2*, Luyuan Zhang1, 2, Sheng Xu3, Weijian Zhou1

- 1. Shaanxi Provincial Key Laboratory of Accelerator Mass Spectrometry and Application, Xi'an AMS Center, SKLLGQ, Institute of Earth Environment, CAS, Xi'an, 710061, China
- 2. Technical University of Denmark, Center for Nuclear Technologies, Roskilde, 4000, Denmark
- 3. Scottish Universities Environmental Research Center, East Kilbride, G75 0QF, United Kingdom *Correspondence to: Xiaolin Hou; E-mail: houxl@ieecas.cn*

Keywords: AMS, iodine-129, aerosol, rainwater, speciation,

The long-lived 129I has been mainly released to the environment from nuclear reprocessing plants, nuclear weapons testing and nuclear accidents. The special source term of 1291, unique chemical properties and environmental behavior of iodine make 129I an ideal tracer to investigation of the pathway of gaseous pollution in the atmosphere, atmospheric chemistry and geochemical cycle of iodine. However, the very low concentration of 129I and its species make their determination a challenge for these investigations. In the past year, our laboratory has developed a series of methods for speciation analysis of 129l in air, aerosol and rainwater (snow) and applied them in the studies of environmental processes. This work aims to give an overview and summary of the methodologies of speciation analysis for 129l in atmosphere and their applications. 129l in iodide and iodate form as well as total iodine in rainwater, particulate associated iodine, gaseous inorganic iodine and gaseous organic iodine species in air samples, water soluble iodide, water soluble iodate, water soluble organic iodine, alkali soluble iodine, and insoluble iodine species in aerosol samples were separated using different physical and chemical methods, and the separated 129I was then measured by AMS and 127I was measured using ICP-MS. The atmospheric behaviors of radioactive iodine released from Fukushima accident have been investigated by speciation analysis of 129I and 127I in time series of aerosol samples and rainwater collected at Fukushima, Japan. The aerosol samples collected in Denmark during the Fukushima accident period were also analyzed for species of 129l and 127l to investigate the sources of 129I in the Europe and the atmospheric behavior of 129I from different sources. Time series of rainwater samples collected in Denmark and seawater samples collected in the North Sea were analyzed for chemical species of 129I and 127I to investigate the interaction of iodine between seawater and atmosphere. Particulate associated iodine, gaseous inorganic iodine and gaseous organic iodine were separately collected at locations with different distance to the North Sea from 50 meter to 1000 km to investigate sources of iodine in the terrestrial, transport pathway of iodine in the atmosphere. A series of snow and seawater samples collected in the Antarctic have been analyzed for 1291, the sources of 1291 in the Antarctic and transport pathway of gaseous pollutant to the Antarctic were investigated.