



## **A risk modelling approach for setting microbiological criteria: using enterococci as indicator for Salmonella in pork**

**Bollerslev, Anne Mette; Hansen, Tina Beck; Nauta, Maarten; Aabo, Søren**

*Publication date:*  
2015

*Document Version*  
Publisher's PDF, also known as Version of record

[Link back to DTU Orbit](#)

*Citation (APA):*

Bollerslev, A. M., Hansen, T. B., Nauta, M., & Aabo, S. (2015). A risk modelling approach for setting microbiological criteria: using enterococci as indicator for Salmonella in pork National Food Institute, Technical University of Denmark. [Sound/Visual production (digital)]. 9th International Conference on Predictive Modelling in Food, Rio de Janeiro, Brazil, 08/09/2015

## **DTU Library** Technical Information Center of Denmark

---

### **General rights**

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.


# A risk modelling approach for setting microbiological criteria: using enterococci as indicator for *Salmonella* in pork

Anne Mette Bollerslev

Tina Beck Hansen

Maarten Nauta


Søren Aabo



# Introduction

- *Salmonella* is considered a leading cause of human enteric disease in the world
- Around 30 % are considered to be caused by pork, in Europe
- From slaughter to retail, growth and cross-contamination increase consumer risk – the process hygiene is important
- Often a hygiene criterion is used to control process hygiene

# Correlation between enterococci and *Salmonella*





**Is it possible to estimate the consumer risk of salmonellosis from eating pork by measuring the level of a hygiene indicator?**




# Objective

- To develop an approach which can be used to define risk-based microbiological criteria for enterococci, to control *Salmonella* in pork cutting plants and retail butchers
- How?
  - By combining the observed positive correlation between *Salmonella* and enterococci with consumer risk modelling\*
  - By associating a consumer risk to a certain level of enterococci

\*Duarte ASR, Nauta MJ, Aabo S. Variation in the effect of carcass decontamination impacts the risk for consumers. Food Control 2016;59:12-19.

# Applied risk-based approach



## Primary study - method


- 5,310 pork samples
- 18 Cutting plants and 414 retail butchers in DK
- Cuttings and minced meat
- Analyses:
  - *Salmonella* (qualitatively, semi-quantitatively)
  - enterococci (quantitatively)





# Results from primary sampling – correlation between *Salmonella* and enterococci

	Conc. [CFU/g]
K	< 0.04
A	0.04 – 0.4
B	0.4 – 4
C	4 – 40
D	40 – 400
E	> 400



Concentration of *Salmonella*

$N_{\text{total}} = 5,310$

$N = 5,215$

61

20

11




2

1


# Growth experiment - method

- Media: BHI broth
- Bacteria:
  - 3 *E. faecium*
  - 3 *E. faecalis*
  - 3 *S. enterica* subsp. *enterica*Serovars: Typhimurium, Infantis and Derby
- Temperatures: 5, 7 and 12° C
- Duration: 5 days

# Results from growth experiment



# Applied risk-based approach



# The risk model - method

- Based on the quantitative model developed by Duarte et al. (2016)\*
- Input:
  - The level of enterococci
  - The corresponding level of *Salmonella* is assumed to be the mean of the semi-quantitative interval
- Output:
  - Relative mean population risk of salmonellosis
  - Number of salmonellosis cases associated to each enterococci level
- Assumptions:
  - Enterococci and *Salmonella* have common growth characteristics
  - The reduction factor  $a$ , is based on 507,222,432 servings of 100 g pork yearly in DK

\*Duarte ASR, Nauta MJ, Aabo S. Variation in the effect of carcass decontamination impacts the risk for consumers

## Scenarios tested in the risk model

	Unit	Sc. 1	Sc. 2	Sc. 3
Limit for enterococci	Log CFU/g	4.0	3.0	2.0

# Results from the risk model

	Unit	Sc. 1	Sc. 2	Sc. 3
Limit for enterococci	Log CFU/g	4.0	3.0	2.0
Relative risk reduction	%	75.7	86.6	88.2
Samples with a level of enterococci > the limit value	%	0.36	3.99	15.8

# Perspectives

- To use this risk-based approach for defining microbiological limits for a hygiene indicator
- To develop risk-based process hygiene criteria in pork cutting plants and retail butchers



## Sum up

- Positive correlation between the quantitative level of enterococci and *Salmonella* in pork
- By use of risk modelling the level of enterococci was associated to a relative consumer risk of salmonellosis
- The approach can be used to define microbiological limits in a risk-based manner

**Thank you for your attention!**