

Technical University of Denmark



Swine influenza viruses in circulation in European pigs exhibit an increasing genetic diversity since last pandemic 2009

Simon, G.; Larsen, Lars Erik; Dürrwald, R.; Foni, E.; Harder, T; Markowska-Daniel, I.; Dan, A.; Maldonado, J.; Huovilainen, A.; Billinis, Charalambos; Davidson, I.; Reid, S. M.; Brown, I. H.; Loeffen, Willie

Published in:

Proceedings of the 7th european symposium of porcine health management

Publication date:
2015

Document Version
Publisher's PDF, also known as Version of record

[Link back to DTU Orbit](#)

Citation (APA):

Simon, G., Larsen, L. E., Dürrwald, R., Foni, E., Harder, T., Markowska-Daniel, I., ... Loeffen, W. (2015). Swine influenza viruses in circulation in European pigs exhibit an increasing genetic diversity since last pandemic 2009. In Proceedings of the 7th european symposium of porcine health management (pp. 80-80). [014] Nantes, France.

DTU Library
Technical Information Center of Denmark

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

014

SWINE INFLUENZA VIRUSES IN CIRCULATION IN EUROPEAN PIGS EXHIBIT AN INCREASING GENETIC DIVERSITY SINCE LAST PANDEMIC IN 2009.

G. Simon (1), L. E. Larsen (2), R. Dürrwald (3), E. Foni (4), T. Harder (5), K. Van Reeth (6), I. Markowska-Daniel (7), A. Dan (8), J. Maldonado (9), A. Huovilainen (10), C. Billinis (11), I. Davidson (12), S. M. Reid (13), I. H. Brown (13), W. Loeffen (14)

(1) Anses, Ploufragan, France; (2) DTU, Copenhagen, Denmark; (3) IDT-Biologika GmbH, Dessau-Rosslau, Germany; (4) IZSLER, Parma, Italy; (5) FLI, Greifswald-Insel Reims, Germany; (6) Ghent University, Ghent, Belgium; (7) NVRI, Pulawy, Poland; (8) Nebih, Budapest; (9) HIPRA, Gerona, Spain; (10) EVIRA, Helsinki, Finland; (11) University of Thessaly, Karditsa, Greece; (12) KVL, Rishon L'Tzion, Israel; (13) AHVLA, Weybridge, UK; (14) CVI, Lelystad, Netherlands

Swine influenza causes concern for veterinary and public health. Thus, ongoing knowledge of influenza A viruses circulating in pigs is necessary for the prevention and control of the disease in pigs, but also to detect reassortant viruses that may pose a threat to humans, as was the case for the H1N1 virus responsible for the last pandemic in 2009 (H1N1pdm).

In continuation to two previous coordination actions (2001-2008) that initiated the surveillance for swine influenza viruses (SIVs) circulating in European pigs, a third phase of the European Surveillance Network for Influenza in Pigs (ESNIP3) aimed to expand the knowledge of European SIVs from 2010 to 2013. ESNIP3 stimulated surveillance in European countries and supported the coordination of appropriate diagnostic tools and subtyping methods. Thus, in an extensive virological monitoring, mainly conducted through passive surveillance programs, more than 9 000 herds in 17 countries were examined over a period of three years.

Influenza A viruses were detected in 31% of herds, from which 1887 viruses were subtyped preliminary. Viruses of the 3 subtypes enzootic in swine (H1N1, H3N2, H1N2) were identified but several genetic lineages were distinguished depending on the origin of genomic segments. Thus, the dominating lineages were identified to represent the 3 previously known European enzootic SIVs, i.e. the avian-like swine H1N1 (53.6%), the human-like reassortant swine H1N2 (13%) and the human-like reassortant swine H3N2 (9.1%), but also the H1N1pdm lineage (10.3%). Viruses from these four lineages co-circulated in several countries, but with very different relative levels of incidence. The H3N2 lineage was missing from some areas whereas it was still prevalent in other parts of Europe. Interestingly, regions free of H3N2 were those that exhibited the highest frequencies of H1N2 viruses in circulation. H1N1pdm viruses were detected at an increasing rate in some countries throughout the three year period, indicating that this subtype has become established in the European pig population. Finally, based on HA and NA identification, 13.9% of the viruses represented reassortants between these four lineages. Among them, reassortants between older enzootic SIVs and H1N1pdm emerged in several countries and were detected at an increasing frequency in 2012-2013.

Surveillance of SIV infections on a European scale revealed striking differences between countries participating in the ESNIP3 network regarding the prevalence of ancient and newly emerged SIV lineages. Some novel reassortants might become established in pig herds in the near future and potentially provide implications for zoonotic infections.

NOTES

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....