

Technical University of Denmark



## Small-scale distribution of Baltic sprat: Statistical properties of school height, length and biomass imply school formation and species interaction mechanisms

Neuenfeldt, Stefan; Gloes, Dominik; Möller, Klas Ove; Möllmann, Christian

*Publication date:*  
2015

*Document Version*  
Publisher's PDF, also known as Version of record

[Link back to DTU Orbit](#)

*Citation (APA):*  
Neuenfeldt, S., Gloes, D., Möller, K. O., & Möllmann, C. (2015). Small-scale distribution of Baltic sprat: Statistical properties of school height, length and biomass imply school formation and species interaction mechanisms. Abstract from ICES Annual Science Conference 2015, Copenhagen, Denmark.

**DTU Library**  
Technical Information Center of Denmark

---

### General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

**Small-scale distribution of Baltic sprat: Statistical properties of school height, length and biomass imply school formation and species interaction mechanisms.**

Stefan Neuenfeldt, Dominik Gloes, Klas Ove Möller and Christian Möllmann

A 10 km acoustic transect in the central Bornholm Basin of the Baltic Sea has been sampled constantly sailing back and forth during a 48 h period. Sprat schools were identified and their height, width and acoustic backscatter as approximation for the fish density in each school were recorded using the Echoview software (Myriax) with the school detection module. School height, length and backscatter (nautical area backscattering coefficient) were all clearly log-normally distributed. Here, we present a simple mechanisms leading to the log-normal distributions, based on Gibrat's law of proportional growth, and discuss alternative causes for the observed apparent universality in the size distributions of the sprat schools. Furthermore, the observed aggregation pattern gives rise to several testable hypotheses on the school formation process and related species interactions during dusk and dawn which are developed and discussed.