

Optical tweezers theory near a flat surface: a perturbative method

Flyvbjerg, Henrik; Dutra, Rafael de Sousa; Neto, Paulo A. Maia; Nussenzveig, H. Moyses

Publication date:
2016

Document Version
Peer reviewed version

[Link back to DTU Orbit](#)

Citation (APA):

Flyvbjerg, H., Dutra, R. D. S., Neto, P. A. M., & Nussenzveig, H. M. (2016). Optical tweezers theory near a flat surface: a perturbative method. Abstract from APS March Meeting 2016, Baltimore, MD, United States.

DTU Library

Technical Information Center of Denmark

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Abstract Submitted
for the MAR16 Meeting of
The American Physical Society

Sorting Category: 06.16 (T)

Optical tweezers theory near a flat surface: a perturbative method¹ HENRIK FLYVBJERG, RAFAEL S. DUTRA, Department of Micro- and Nanotechnology, Technical University of Denmark, PAOLO A. MAIA NETO, H. MOYSES NUSSENZVEIG, Institute of Physics, Universidade Federal do Rio de Janeiro — We propose a perturbative calculation of the optical force exercised by a focused laser beam on a microsphere of arbitrary radius that is localized near a flat glass surface in a standard optical tweezers setup. Starting from the Mie-Debye representation for the electric field of a Gaussian laser beam, focused by an objective of high numerical aperture, we derive a recursive series that represents the multiple reflections that describe the reverberation of laser light between the microsphere and the glass slide. We present numerical results for the axial component of the optical force and the axial trap stiffness. Numerical results for a configuration typical in biological applications a microsphere of 0.5 μm radius at a distance around 0.25 μm from the surface show a 37% [1] Viana N B, Rocha M S, Mesquita O N, et al. (2007) Towards absolute calibration of optical tweezers. Phys Rev E 75:021914-114. [2] Dutra R S, Viana N B, Maia Neto P A, et al. (2014) Absolute calibration of forces in optical tweezers. Phys Rev A 90:013825-113.

¹Rafael S. Dutra thanks the Brazilian "Science without Borders" program for a postdoctoral scholarship.

Prefer Oral Session
 Prefer Poster Session

Henrik Flyvbjerg
henrik.flyvbjerg@nanotech.dtu.dk
Department of Micro- and Nanotechnology

Date submitted: 05 Nov 2015

Electronic form version 1.4