

## RESEARCH

## Open Access

# Maternal educational level and children's healthy eating behaviour: role of the home food environment (cross-sectional results from the INPACT study)

Wilke JC van Ansem<sup>1,2\*</sup>, Carola TM Schrijvers<sup>1,2</sup>, Gerda Rodenburg<sup>1,2</sup> and Dike van de Mheen<sup>1,2,3</sup>

## Abstract

**Background:** The aims of this study are 1) to investigate the association between maternal educational level and healthy eating behaviour of 11-year-old children (fruit, vegetables and breakfast consumption), and 2) to examine whether factors in the home food environment (parental intake of fruit, vegetables and breakfast; rules about fruit and vegetables and home availability of fruit and vegetables) mediate these associations.

**Methods:** Data were obtained from the Dutch INPACT study. In total, 1318 parent-child dyads were included in this study. Multilevel regression models were used to investigate whether factors of the home food environment mediated the association between maternal educational level and children's healthy eating behaviour.

**Results:** Children of mothers with a high educational level consumed more pieces of fruit per day ( $B = 0.13$ , 95% CI: 0.04-0.22), more grams of vegetables per day ( $B = 23.81$ , 95% CI = 14.93-32.69) and were more likely to have breakfast on a daily basis ( $OR = 2.97$ , 95% CI: 1.38-6.39) than children of mothers with a low educational level. Home availability, food consumption rules and parental consumption mediated the association between maternal education level and children's fruit and vegetable consumption. Parental breakfast consumption mediated the association between maternal education level and children's breakfast consumption.

**Conclusions:** Factors in the home food environment play an important role in the explanation of socio-economic disparities in children's healthy eating behaviour and may be promising targets for interventions.

**Keywords:** Fruit and vegetable consumption, Breakfast consumption, Socio-economic status, Home food environment, Children, Parents

## Background

Dietary behaviour is important for the development and growth of children and also influences health outcomes later in life. Fruit and vegetables and daily breakfast consumption are important components of a healthy diet and their beneficial effects on health are well documented. Diets rich in fruit and vegetables protect against cardiovascular disease (CVD), some types of cancer, and obesity [1,2]. Regular breakfast consumption is associated with

better cognitive performance and a reduced risk of becoming overweight or obese among children and adolescents [3,4]. Despite the importance of healthy dietary behaviours, the majority of the children in the Netherlands, as in other countries, does not consume the recommended amounts of fruit and vegetables [5-8]. In addition, breakfast skipping is highly prevalent in Europe and the United States [9,10]. Also, because dietary habits track into adulthood, it is important to develop interventions aimed to improve dietary behaviours of children [11,12].

Children and adolescents with a low socio-economic status (SES) consume less fruit and vegetables than children and adolescents with a high SES [13-16]. Furthermore, a

\* Correspondence: [vanansem@ivo.nl](mailto:vanansem@ivo.nl)

<sup>1</sup>IVO Addiction Research Institute, Heemraadssingel 194, Rotterdam, DM 3021, The Netherlands

<sup>2</sup>Erasmus Medical Centre, Postbox 2040, Rotterdam, CA 3000, The Netherlands

Full list of author information is available at the end of the article

Norwegian study found an increase in socio-economic disparities in adolescent's fruit and vegetable consumption between 2001 and 2008 [17]. Studies of socio-economic disparities in breakfast consumption showed inconsistent findings. A literature review found that parental educational level and parental unemployment were unrelated to adolescents and children's breakfast consumption [18]. However, other studies found a positive association between maternal educational level and children's breakfast consumption [9,19,20]. Given the inconsistencies in the findings from previous studies and the relative small part of the literature assessing socio-economic disparities in dietary behaviour of children, the first aim of this study is to investigate socio-economic differences in healthy eating behaviours of children (fruit, vegetable and breakfast consumption).

The home food environment is important in the development of children's dietary behaviour [21]. Parents have an important influence on the dietary behaviour of children because they generally determine which food is available at home, they can set rules about what their children are allowed to eat and they act as role models, also with respect to dietary behaviour [22]. Several literature reviews concluded that aspects of the home environment are associated with children's fruit and vegetable intake [16,23,24]. Home environmental factors found to be positively related to children's fruit and vegetable intake


are home availability, family rules and parental intake. For breakfast consumption, parental breakfast consumption is an important home environmental factor that is positively associated with children's breakfast consumption [18].

As stated before, the first aim of this study is to investigate socio-economic differences (maternal educational level is used as indicator for children's SES) in healthy eating behaviours of children. However, SES does not directly influence dietary behaviour and is not a modifiable correlate of children's dietary behaviour. Thus it is important to identify modifiable determinants that may explain the socio-economic disparities in children's healthy eating behaviour. Therefore, the second aim of this study is to examine whether factors in the home food environment (parental intake of fruit, vegetables and breakfast; rules about fruit and vegetables and home availability of fruit and vegetables) mediate the association between maternal educational level and children's healthy eating behaviours (fruit, vegetable and breakfast consumption). Figure 1 presents the research model.

## Methods

### Study population and design

The data used in this study are derived from the Dutch INPACT study, INPACT being the acronym for IVO Nutrition Physical Activity Child cohort. This longitudinal study among 8 to 12 year olds and their parents



**Figure 1** Research model.

investigated modifiable environmental determinants of children's dietary behaviour. Participants of the INPACT study were recruited through primary schools in the southern part of the Netherlands (Eindhoven and surroundings). The municipal health service invited all general primary schools ( $n = 265$ ) in this area to participate in this study. Ninety one schools (34.3%) agreed. The response rate of schools in rural and urban areas was similar. A sample of 1844 parent-child dyads (62.5%) gave informed consent. Trained research assistants visited the participating primary schools and measured children's height and weight. Children completed a short questionnaire at school and parents completed a questionnaire at home. The questionnaire topics varied annually. The INPACT study was approved by the Medical Ethical Committee at Erasmus Medical Centre, Rotterdam. The present study was based on cross-sectional data collected in the last wave (2011), in which a questionnaire was completed by 1428 primary caregivers. In most cases ( $n = 1312$ , 92.1%) the primary caregiver was the mother.

## Measurements

### *Socio-economic status*

The three most commonly used indicators of SES are educational level, income and occupation [25]. Of these three SES indicators, educational level was found to be the strongest and most consistent in predicting health behaviour [25]. In this study, maternal educational level was used as an indicator of children's SES because several studies found maternal educational level to be a reliable determinant of children's dietary behaviour and childhood obesity [9,19,20]. In addition, traditionally, in the majority of the households the mother provides the food for the family and, therefore, maternal educational level also impacts the dietary behaviour of the other members of the family and the home availability of foods. Maternal educational level was classified into three groups: 'low educational level' (primary school and lower secondary education), 'intermediate educational level' (intermediate vocational level, higher secondary school and pre-university education) and 'high educational level' (higher vocational education and university). Throughout the remainder of this paper we thus refer to 'low SES' (children of mothers with a low educational level), 'intermediate SES' (children of mothers with an intermediate educational level) and 'high SES' (children of mothers with a high educational level).

### *Outcome measures (children's fruit, vegetable and breakfast consumption)*

Children's fruit, vegetable and breakfast consumption were measured with a questionnaire based on a validated Food Frequency Questionnaire [26-29]. Parents reported how many days in a normal week their child consumed 1) fruit

(fresh or canned fruit), 2) cooked, fried, steamed or otherwise heated vegetables, 3) salad or other raw vegetables and 4) breakfast. Answering categories ranged from 'none or less than one day a week' to '7 days a week'.

Additionally, parents reported the numbers of servings of fruit and vegetables consumed by their child on such a day. For fruit, answer categories ranged from '0 pieces a day' to 'more than 3 pieces a day', by increments of half a piece of fruit. Reported fruit consumption of more than 3 pieces a day ( $n = 4$ ) was recoded as '4 pieces a day'. For vegetables, answer categories ranged from '0 serving spoons' to 'more than 4 serving spoons a day', by increments of half a serving spoon. Reported vegetable consumption of more than 4 serving spoons ( $n = 12$ ) was recoded as '5 serving spoons a day'. One serving spoon of vegetables was equivalent to 50 grams of vegetables. Total vegetable consumption was calculated in grams for each child by multiplying consumption frequency (how many days a child consumed vegetables) and serving spoons of vegetables. Subsequently, the vegetable consumption was converted to an amount consumed in a day. Total fruit consumption was calculated in pieces for each child by multiplying consumption frequency (how many days a child consumed fruit) and servings (pieces of fruit). Children's fruit consumption was also converted to an amount consumed in a day.

Breakfast consumption was dichotomized into 'daily' and 'not daily', due to limited variation in the answering categories.

### *Potential mediating variables*

**Parental intake of fruit, vegetables and breakfast** Parental fruit, vegetable and breakfast consumption were measured and calculated in the same way as children's fruit, vegetable and breakfast consumption.

**Rules about fruit and vegetable consumption** We assessed whether parents set rules regarding their child's fruit and vegetable consumption with the following questions: 'Do you have the rule that your child should eat 2 pieces of fruit a day?' and 'Do you have the rule that your child should eat 200 grams of vegetables a day?' These specific amounts of fruit and vegetables are consistent with the Dutch guidelines for fruit and vegetables [30]. Response categories were 'yes' and 'no'. These questions were derived from the ENDORSE study [31].

**Home availability of fruit and vegetables** The availability of fruit and vegetables at home was measured using a questionnaire based on the validated Home Environment Survey [32]. Parents were asked about the availability of 1) fruit and 2) vegetables in their home. Response categories were 'yes, always', 'yes, usually', 'sometimes', 'no, usually not' and 'no, never'. Due to limited variability of

these variables, we dichotomized both variables into ‘always’ (‘yes, always’) and ‘not always’ (‘yes, usually’; ‘sometimes’; ‘no, usually not’; ‘no, never’).

#### Potential confounders

The following variables are considered as potential confounders: age, gender, ethnicity and body mass index (BMI) of the child. Age, gender and ethnicity of the child were reported by the parents. A child’s age (in years) was calculated on the basis of the date of birth and the date of measurement. For the purpose of analysis we dichotomised child’s age into ‘≤ 11 years’ versus ‘> 11 years’. Children’s ethnicity was categorised into ‘Dutch native’ (both parents born in the Netherlands) and ‘immigrants’ (at least one of the parents was born outside the Netherlands). Children’s body mass index (BMI) was calculated on the basis of weight and height, which were measured with clothes but without shoes to the nearest 0.1 kg and 0.1 cm; the measurements were made by trained research assistants. BMI cut-off points for children were used to define overweight and obesity [33]. Subsequently child BMI was dichotomised into ‘overweight’ (‘overweight’ and ‘obesity’) versus no overweight (‘underweight and normal weight’).

#### Data analysis

Respondents who lacked data for maternal educational level were excluded from this study (n = 110, 7.7%). In total, 1318 children and parents were included in this study. Descriptive analyses were performed to describe the characteristics of the study population.

To investigate whether home environmental factors mediated the association between maternal educational level and children’s healthy eating behaviour, we used Baron and Kenny’s four-step approach [34]. According

to Baron and Kenny, there are three criteria for mediation: 1) the predictive variable has to be associated with the outcome variable, 2) the predictive variable has to be associated with the mediator, and 3) the mediator has to be associated with the outcome variable (adjusted for the predictive variable). If all the associations assessed in steps 1–3 are statistically significant, the criteria for mediation have been met. Step four of the approach is to test the mediation model: mediation is supported if the association between the predictive variable and the outcome variable changes after controlling for the mediator.

For each outcome measure (child fruit intake, child vegetable intake and child breakfast consumption) the steps of the mediation approach were conducted separately. Depending on the scale of the outcome measures, logistic regression models or linear regression models were used to test the subsequent steps of the mediation-approach.

Several potential mediators were tested for the outcome measures ‘child’s fruit consumption’ and ‘child’s vegetable consumption’. If it appeared that more than one potential mediator met the criteria for mediation, the unique contribution of each mediator was determined (single mediator model). Next, a multivariate mediation model was tested. Bootstrapping resampling techniques were used to calculate confidence intervals for the mediated effects.

All regression models were adjusted for the potential confounders. Due to the used sample-strategy (children were recruited through schools), the data have a nested structure (children within schools). To take into account potential clustering effects, we investigated the associations using multilevel regression analyses. Analyses were performed using R (2013). Cases with missing values were removed per analysis. Due to missing values the computed

**Table 1 Characteristics of the study population: total sample and sample according to socio-economic status (SES)**

	<i>Total sample</i>	<i>Low SES</i>	<i>Intermediate SES</i>	<i>High SES</i>	<i>P-value</i>
<b>Mean age N (%)</b>	N = 1317	N = 263	N = 628	N = 426	<b>0.00</b>
≤11 years	1119 (85.0)	205 (77.9)	528 (84.1)	386 (90.6)	
> 11 years	198 (15.0)	58 (22.1)	100 (15.9)	40 (9.4)	
<b>Gender %</b>	N = 1318	N = 263	N = 629	N = 426	<b>0.02</b>
Boys	50.8	52.5	46.7	55.6	
Girls	49.2	47.5	53.3	44.4	
<b>Child’s BMI %</b>	N = 1283	N = 252	N = 616	N = 415	<b>0.01</b>
Overweight	11.2	16.7	10.6	8.9	
No overweight	88.8	83.3	89.4	91.9	
<b>Child’s ethnicity %</b>	N = 1318	N = 263	N = 629	N = 426	<b>0.02</b>
Native Dutch	88.8	84.4	90.9	89.4	
Immigrant	11.2	15.6	9.1	10.6	

models for fruit, vegetable and breakfast consumption are based on different numbers of participants.

## Results

Background characteristics of the study population are presented in Table 1. Mean age of the children was 11 years, the majority was native Dutch and not overweight, and about half were boys. Significant differences between the three educational levels were found in the background characteristics: relative to the high and low SES groups there were more girls in the intermediate SES group. More children in the low SES group were overweight, were older than 11 years, and were

immigrants compared with children in the intermediate and high SES groups.

**Table 2 Descriptives of the key study variables**

	Socio-economic status (SES)		
	Low	Intermediate	High
<b>Fruit (N = 1269)</b>	N = 247	N = 607	N = 415
Children's fruit intake, pieces per day (mean, SD)	0.96 (0.65)	0.99 (0.57)	1.07 (0.60)
Parental fruit intake, pieces per day (mean, SD)	0.97 (0.74)	1.04 (0.74)	1.19 (0.73)
Parental rules regarding fruit consumption (%)			
Yes	68.8	72.7	79.5
No	31.2	27.3	20.5
Home availability of fruit %			
Always	88.3	92.8	94.2
Not always	11.7	7.2	5.8
<b>Vegetables (N = 1265)</b>	N = 248	N = 606	N = 411
Children's vegetable intake, grams per day (mean, SD)	94.0 (57.7)	100.5 (53.1)	116.9 (60.6)
Parental vegetable intake, grams per day (mean, SD)	148.3 (68.1)	158.4 (67.9)	176.7 (68.0)
Parental rules regarding vegetable consumption (%)			
Yes	83.5	85.0	92.0
No	16.5	15.0	8.0
Home availability of vegetables			
Always	83.1	89.4	90.5
Not always	16.4	8.0	9.5
<b>Breakfast (N = 1270)</b>	N = 246	N = 610	N = 414
Children's breakfast consumption (%)			
Daily	91.9	94.3	97.3
Not daily	8.1	5.7	2.7
Parental breakfast consumption (%)			
Daily	83.7	91.3	95.7
Not daily	16.3	8.7	4.3

## Fruit consumption

Table 2 provides data on children's fruit consumption and determinants of children's fruit consumption stratified by SES. Children with a low SES had the lowest fruit consumption (on average 0.96 pieces per day) while children with a high SES had the highest fruit consumption (on average 1.07 per day). Table 3 presents data on the association between children's SES and their fruit consumption. Children with a high SES consumed more fruit than children with a low SES (B = 0.13, 95% CI: 0.04-0.22). There was no significant difference in fruit consumption between children with an intermediate SES and those with a low SES.

Table 4 presents data on the association between children's SES and possible mediating variables regarding fruit consumption. Parents with a high SES consumed significantly more fruit (B = 0.25, 95% CI: 0.13-0.36), were more likely to have rules about fruit consumption (OR = 1.78, 95% CI: 1.23-2.56) and were more likely to always have fruit available at home (OR = 2.24, 95% CI: 1.25-4.00) than parents with a low SES. Parents with an intermediate SES were also more likely to always have fruit available at home than parents with a low SES (OR = 1.74, 95% CI: 1.05-2.88).

**Table 3 Associations between socio-economic status (SES) and children's fruit, vegetable and breakfast consumption**

Fruit consumption (N = 1269)	Multivariate regression analyses <sup>†</sup>	
SES	B (95% CI)	P-value
Low (Ref. group)	0.84	
Intermediate	0.04 (-0.05 - 0.13)	0.38
High	<b>0.13 (0.04 - 0.22)</b>	<b>0.01</b>
Vegetable consumption (N = 1265)	Multivariate regression analyses <sup>†</sup>	
SES	B (95% CI)	P-value
Low (Ref. group)	83.89	
Intermediate	<b>8.33 (0.09 - 16.56)</b>	<b>0.05</b>
High	<b>23.81 (14.93 - 32.69)</b>	<b>0.00</b>
Breakfast consumption (N = 1270)	Multivariate regression analyses <sup>†</sup>	
SES	OR (95% CI)	P-value
Low (Ref. group)	1.00	
Intermediate	1.39 (0.78 - 2.49)	0.27
High	<b>2.97 (1.38 - 6.39)</b>	<b>0.01</b>

B = unstandardized coefficient, OR = Odds ratio, 95% CI = 95% Confidence Interval. Bold values represent statistically significant association. <sup>†</sup>Multivariate regression analysis adjusted for: child's age, child's gender, child's ethnicity and child's BMI.



**Table 5 Associations between possible mediating variables and children’s fruit, vegetable and breakfast consumption**

<b>Fruit consumption (N = 1269)</b>	<b>Multivariate regression analyses<sup>†</sup></b>	
	<b>B (95% CI)</b>	<b>P-value</b>
<i>Parental fruit consumption</i>	<b>0.34</b> (0.30 – 0.39)	<b>0.00</b>
<i>Rules about fruit consumption</i>		
No (ref. group)	0.55	
Yes	<b>0.49</b> (0.42 – 0.56)	<b>0.00</b>
<i>Home availability of fruit</i>		
Not always (ref. group)	0.42	
Always	<b>0.48</b> (0.36 – 0.60)	<b>0.00</b>
<b>Vegetable consumption (N = 1265)</b>	<b>Multivariate regression analyses<sup>†</sup></b>	
	<b>B (95% CI)</b>	<b>P-value</b>
<i>Parental vegetable consumption</i>	<b>0.46</b> (0.42 – 0.47)	<b>0.00</b>
<i>Rules about vegetable consumption</i>		
No (ref. group)	63.93	
Yes	<b>24.94</b> (15.20 – 34.68)	<b>0.00</b>
<i>Home availability of vegetables</i>		
Not always (ref. group)	67.66	
Always	<b>18.62</b> (9.72 – 27.51)	<b>0.00</b>
<b>Breakfast consumption (N = 1270)</b>	<b>Multivariate regression analyses<sup>†</sup></b>	
	<b>OR (95% CI)</b>	<b>P-value</b>
<i>Parental breakfast consumption</i>		
Not daily (ref. group)	1.00	
Daily	<b>15.75</b> (9.04 – 27.44)	<b>0.00</b>

B = unstandardized coefficient, OR = Odds ratio, 95% CI = 95% Confidence Interval. Bold values represent statistically significant association. <sup>†</sup>Multivariate regression analysis adjusted for: child’s SES, child’s age, child’s gender, child’s ethnicity and child’s BMI.

### Vegetable consumption

Table 2 provides data on children’s vegetable consumption and determinants of children’s vegetable consumption stratified by SES. Children with a low SES had the lowest vegetable consumption (on average 94.0 grams per day) while children with a high SES had the highest vegetable consumption (on average 116.9 grams per day).

Table 3 shows significant socio-economic differences in children’s vegetables consumption. Children with an intermediate SES and children with a high SES consumed more vegetables than children with a low SES (resp. B = 8.33, 95% CI: 0.09-16.56; B = B = 23.81, 95% CI: 14.93-32.96).

Table 4 presents data on the association between SES and possible mediating variables regarding vegetable consumption. Parents with a high SES consumed more vegetables (B = 28.86, 95% CI: 18.05-39.67), were more likely

to have vegetable consumption rules (OR = 2.47, 95% CI: 1.49-4.10), and were more likely to always have vegetables available at home (OR = 1.93, 95% CI: 1.19-3.11) than parents with a low SES. Parents with an intermediate SES also consumed more vegetables (B = 11.29 95% CI: 1.24-21.34) and were more likely to have rules about vegetable consumption (OR: 1.74, 95% CI: 1.13-2.69) than parents with a low SES.

All potential mediators were significantly associated with children’s vegetable consumption (see Table 5). Children consumed more vegetables when their parents consumed more vegetables (B = 0.46, 95% CI: 0.42-0.47), when their parents had rules about vegetable consumption (OR = 24.94, 95% CI: 15.20-34.68), and when vegetables were always available at home (B = 18.62, 95% CI: 9.72-27.51).

Table 6 presents the mediation models. In the single-mediator models, parental vegetable intake explained 56.1% of the difference in vegetable consumption between children with a low SES and those with a high SES; vegetable consumption rules explained 9.9% and home availability of fruit explained 6.0%. In the multiple-mediator model, all the mediators together explained 58.89% of the difference in vegetable intake between children with a low SES and those with a high SES. Parental vegetable intake, vegetable consumption rules and home availability of vegetables had no significant mediating effect on the difference in vegetable intake between children with an intermediate SES and those with a low SES.

### Breakfast consumption

Table 2 presents data on children’s and parents breakfast consumption. Children and parents with a high SES more often reported to have breakfast on a daily basis than children and parents with a low and intermediate SES. Table 3 reports on the association between SES and children’s breakfast consumption. Children with a high SES were more likely to eat breakfast on a daily basis than children with a low SES (OR = 2.97, 95% CI: 1.38-6.39). There was no significant difference in breakfast consumption between children with an intermediate SES and those with a low SES.

Parents with high and intermediate SES were more likely to consume breakfast on a daily basis than parents with a low SES (see Table 4). Table 5 shows that children were more likely to eat breakfast on a daily basis when their parents ate breakfast on a daily basis (OR = 15.75, 95% CI: 9.04-27.44). Table 6 shows the final mediation model; parental breakfast consumption explained 67.9% of the differences in breakfast consumption between children with a high SES and those with a low SES. Parental breakfast consumption had no significant mediating effect on the difference in breakfast consumption between children with an intermediate and with a high SES.

**Table 6 Mediation analyses**

<b>Fruit consumption (N = 1269)</b>	<b>Direct association between SES and children's fruit consumption B</b>	<b>Mediation models B (95% CI)</b>	<b>P-value</b>	<b>Percentage change</b>	<b>P-value</b>
<b>SES</b>					
<i>Model A</i>					
Low (ref. group)	0.84	0.54			
Intermediate	0.04	0.01 (−0.07 – 0.09)	0.84	−79.49	0.36
High	0.13	0.04 (−0.09 – 0.17)	0.32	−66.01	<b>0.02</b>
<b>SES</b>					
<i>Model B</i>					
Low (ref. group)	0.84	0.55			
Intermediate	0.04	0.02 (−0.06 – 0.10)	0.58	−42.16	0.42
High	0.13	0.08 (−0.01 – 0.16)	0.09	−40.85	<b>0.00</b>
<b>SES</b>					
<i>Model C</i>					
Low (ref. group)	0.84	0.42			
Intermediate	0.04	0.02 (−0.07 – 0.10)	0.69	−56.10	0.40
High	0.13	<b>0.10 (0.01 – 0.19)</b>	<b>0.03</b>	−23.15	<b>0.02</b>
<b>SES</b>					
<i>Model D</i>					
Low (ref. group)	0.84	0.23			
Intermediate	0.04	−0.01 (−0.08 – 0.07)	1.12	−114.44	0.40
High	0.13	0.01 (−0.07 – 0.10)	0.75	−89.53	<b>0.00</b>
<b>Vegetable consumption (N = 1265)</b>	<b>Direct association between SES and children's vegetable consumption B</b>	<b>Mediation models B (95% CI)<sup>†</sup></b>	<b>P-value</b>	<b>Percentage change</b>	<b>P-value</b>
<b>SES</b>					
<i>Model E</i>					
Low (ref. group)	83.89	17.91			
Intermediate	8.33	3.19 (−3.66 – 10.05)	0.36	−61.70	0.14
High	23.81	<b>10.44 (2.97 – 17.93)</b>	<b>0.01</b>	<b>−56.13</b>	<b>0.00</b>
<b>SES</b>					
<i>Model F</i>					
Low (ref. group)	83.89	63.93			
Intermediate	8.33	6.75 (−1.42 – 14.92)	0.11	−19.02	0.08
High	23.81	<b>21.47 (12.63 – 30.31)</b>	<b>0.00</b>	<b>−9.85</b>	<b>0.00</b>
<b>SES</b>					
<i>Model G</i>					
Low (ref. group)	83.89	67.66			
Intermediate	8.33	7.39 (−0.26 – 16.12)	0.06	−4.88	0.56
High	23.81	<b>22.39 (13.55 – 31.23)</b>	<b>0.00</b>	<b>−5.98</b>	<b>0.00</b>
<b>SES</b>					
<i>Model H</i>					
Low (ref. group)	83.89	10.57			
Intermediate	8.33	2.75 (−4.11 – 9.61)	0.43	−66.79	0.06
High	23.81	<b>9.77 (2.27 – 17.27)</b>	<b>0.01</b>	<b>−58.89</b>	<b>0.00</b>



**Table 6 Mediation analyses (Continued)**

<i>Breakfast consumption (N = 1270)</i>	<i>Direct association between SES and children's breakfast consumption OR</i>	<i>Mediation model OR (95% CI)<sup>†</sup></i>	<i>P-value</i>	<i>Percentage change</i>	<i>P-value</i>
<b>SES</b>		<i>Model I</i>			
Low (ref. group)	1.00				
Intermediate	1.39	0.99 (0.52 – 1.90)	0.97	- 102.73	0.27
High	2.97	1.63 (0.71 – 3.67)	0.25	<b>- 67.89</b>	<b>0.02</b>

SES = socioeconomic status; B = unstandardized coefficient, OR = Odds ratio, 95% CI = 95% Confidence Interval. Bold values represent statistically significant association.

Model A: Single mediator model. This model includes the mediator 'parental fruit consumption'.

Model B: Single mediator model. This model includes the mediator 'parental rules regarding fruit consumption'.

Model C: Single mediator model. This model includes the mediator 'home availability of fruit'.

Model D: Multiple mediation model. This model includes the mediators: parental fruit consumption, parental rules regarding fruit consumption and home availability of fruit.

Model E: Single mediator model. This model includes the mediator 'parental vegetable intake'.

Model F: Single mediator model. This model includes the mediator 'parental rules regarding vegetable consumption'.

Model G: Single mediator model. This model includes the mediator 'home availability of vegetables'.

Model H: Multiple mediator model. This model includes the mediators: parental vegetable consumption, parental rules regarding vegetable consumption and home availability of vegetables.

Model I: Single mediator model. This model includes the mediator 'parental breakfast consumption'.

All models are adjusted for: child's age, child's gender, child's ethnicity and child's BMI.

## Discussion

The first aim of this study was to examine the association between SES and children's fruit, vegetable and breakfast consumption. We found that children with a high SES consumed more fruit and vegetables and consumed more often breakfast on a daily basis, than children with a low SES. These findings are in line with those from the majority of similar studies [9,13-15,17,19,35] and emphasise that children from low socio-economic groups can be considered an important target for interventions to improve dietary behaviour.

However, maternal education level, (and other measures of SES), are not considered to have a direct effect on dietary behaviour and are not easily modifiable. To explain socio-economic disparities in children's dietary behaviour, several studies examined socio-economic differences in the home food environment of children. These studies showed that the home food environment of children of mothers with a low educational level was less supportive than the home food environment of children of mothers with a high educational level [36,37]. For example, adolescents of mothers with a low educational level were more likely to report that unhealthy foods were always or usually available at home, while adolescents of mothers with a high educational level were more likely to report that fruit was always or usually available at home and that vegetables were always served at dinner time [38]. In addition, a study among 5–6 year old children found comparable results; households of mothers with a low educational level were more likely to watch television while eating dinner and mothers with a low educational level were more likely to have negative perceptions about the quality and variety of fresh fruit and vegetables at their local shops [39]. Furthermore, Hupkens *et al.* found that

mothers with a high educational level more often limited their children's intake of unhealthy foods (e.g. sweets, soft drinks, chips). These differences in the number of restricted foods by educational level were partly explained in health and taste considerations between mothers with a low and high educational level [40]. A more recent study also found socio-economic differences in food parenting practices; frequent consumption of fruit and vegetables, restrictive rules, verbal praise, negotiation and restraint from negative modelling were all more common among mothers with a high educational level [41]. The present study also shows that aspects of the home food environment differed by SES, where low SES had the less supportive home environment.

However, socio-economic differences in determinants of the home food environment do not necessarily account for socio-economic differences in children's dietary behaviour. Therefore, a second aim of this study was to investigate modifiable factors of the home food environment that mediate the association between SES and children's fruit, vegetable and breakfast consumption. We included parental intake, home availability and parental rules about children's fruit and vegetable consumption as possible mediating variables in the association between SES and children's fruit and vegetable intake. Our results indicate that all the studied home environmental factors mediate the association between SES and children's fruit and vegetable intake. Moreover, our results indicate that the difference in fruit and vegetable consumption between children with a low and high SES is explained in particular by parental intake of fruit and vegetables. Very few studies have assessed mediators of the association between socio-economic status and children's fruit and vegetables intake. Vereecken *et al.* found that differences in children's fruit and vegetable

consumption by mother's educational level were completely explained by mother's consumption and parenting practices [41]. Bere *et al.* concluded that home accessibility was the strongest mediator of the association between maternal educational level and adolescent's fruit and vegetable consumption [13]. Furthermore, Hilsen *et al.* also found that accessibility of fruit and vegetables mediates part of the association between socio-economic status and adolescent's fruit and vegetable intake [17]. In addition, they found that accessibility of fruit and vegetables explains part of the increase in SES disparities in fruit and vegetable consumption between 2001 and 2008.

To our knowledge, ours is the first study to assess possible explanatory variables of socio-economic disparities in children's breakfast consumption. We found that the difference in breakfast consumption between high SES children and low SES children was mediated by parental breakfast consumption. However, we included only one possible mediator in our analyses (parental breakfast consumption), while other potentially mediating variables were not included. For example, parenting practices are associated with children's breakfast consumption and may also be an explanatory variable of socio-economic differences in children's breakfast consumption.

It is known that aspects of the home environment are associated with children's dietary behaviour. This study indicates that home environmental factors also play a role in the explanation of socio-economic disparities in children's healthy eating behaviour. Given that parental intake was the strongest mediator and that parents shape the home food environment (e.g. they decide which food is available at home and can set food rules), parents play an important role in the development of children's dietary behaviour. Therefore, parents can be important targets for interventions. Moreover, it is necessary to reach parents with a low SES and to increase their own consumption of fruit, vegetables and breakfast, to increase the home availability of healthy products and to set food rules for their children. Campbell *et al.* found that maternal nutrition knowledge was associated with children's fruit and vegetable consumption and also with the home availability of fruit and vegetables [42]. Therefore, targeting parental nutritional knowledge (especially among those with a low educational level) may be an effective way to improve the home food environment. Besides interventions that aim at the importance of family involvement, also multiple-setting interventions are effective in changing children's dietary behaviour. In the latter case, children receive the messages in more than one setting (e.g. at home, school, and the sports club) thereby increasing the chance that such an intervention will be more effective than a single-setting approach [43]. However, interventions aiming to improve children's dietary behaviour, such as children's fruit and vegetable consumption are also necessary for

children from higher socio-economic backgrounds since the majority of all children (including children of higher educational background) does not consume the recommended amount of fruit and vegetables.

The present study has some limitations. First, this study has a cross-sectional design, which does not allow to draw conclusions about causal relationships. However, as educational level is a consistent factor over time, it is highly unlikely that children's food consumption will affect a mother's educational level. Although it is possible that children's fruit and vegetable consumption contributes to the amount of fruit and vegetables available at home, or to parental consumption rules regarding fruit and vegetables, we believe that the impact of the home availability and the consumption rules of fruit and vegetables on children's fruit and vegetable intake are larger. Therefore, we expect the directions of the associations we found to be as presented in Figure 1. Second, assessments of child's fruit, vegetable and breakfast consumption were based on parent's reports instead of child reports. Child reports might be more valid, although this remains unclear. Nevertheless, Tak *et al.* [44] concluded that parents' reports could be considered as a valid method to measure children's fruit and vegetable consumption, although the use of parent's reports may evoke socially desirable answers. Finally, we measured breakfast frequency and not breakfast quality, which is associated with the nutrient adequacy of diets [45].

## Conclusion

This study shows that children of mothers with low educational level have less healthy eating habits than children of mothers with a high educational level. Our study adds to the knowledge on possible mechanisms underlying socio-economic differences in healthy eating behaviour of children. Parent's food intake, home availability of healthy foods and parental rules about children's fruit and vegetable intake mediated the association between maternal educational level and children's healthy eating behaviour. Interventions to improve children's dietary behaviour and to reduce socio-economic disparities in children's eating habits, may benefit by focusing on the role of parents in the development of children's dietary behaviour.

## Competing interests

The authors declare that they have no competing interests.

## Authors' contribution

WJCvA, CTMS and DvdM were involved in the design of this study. WJCvA was responsible for data collection, performed the statistical analyses and drafted the manuscript. CTMS was the daily supervisor of the project. CTMS, DvdM and GR helped with the interpretation of the data. GR revised the manuscript. CTMS and DvdM helped to draft the manuscript. All authors read and approved the final manuscript.

## Acknowledgements

The authors thank V. Dam, S. de Schepper, I. de Vlieger, Y. Pierik, S. Pooyé and R. Roubéhié-Fissa for their help in the data collection. This project was financed by the Netherlands Organisation for Health Research and Development (ZonMw; project no. 115100004).

## Author details

<sup>1</sup>IVO Addiction Research Institute, Heemraadssingel 194, Rotterdam, DM 3021, The Netherlands. <sup>2</sup>Erasmus Medical Centre, Postbox 2040, Rotterdam, CA 3000, The Netherlands. <sup>3</sup>Department of Health Education and Promotion, Maastricht University, Maastricht, The Netherlands.

Received: 15 October 2013 Accepted: 2 September 2014

Published online: 12 September 2014

## References

1. Johnsen SP: Intake of fruit and vegetables and risk of stroke: an overview. *Curr Opin Clin Nutr Metab Care* 2004, **7**:665–670.
2. Cancer Research Fund/American Institute for Cancer: *Food, Nutrition, Physical Activity and the Prevention of Cancer: A Global Perspective*. DC: AICR; 2007.
3. Hoyland A, Dye L, Lawton CL: A systematic review of the effect of breakfast on the cognitive performance of children and adolescents. *Nutr Res Rev* 2009, **22**:220–243.
4. Szajewska H, Ruszczyński M: Systematic review demonstrating that breakfast consumption influences body weight outcomes in children and adolescents in Europe. *Crit Rev Food Sci Nutr* 2010, **50**:113–119.
5. Diethelm K, Jankovic N, Moreno LA, Huybrechts I, De Henauw S, De Vriendt T, González-Gross M, Leclercq C, Gottrand F, Gilbert CC, Dallongeville J, Cuenca-Garcia M, Manios Y, Kafatos A, Plada M, Kersting M: Food intake of European adolescents in the light of different food-based dietary guidelines: results of the HELENA (Healthy Lifestyle in Europe by Nutrition in Adolescence) Study. *Public Health Nutr* 2012, **15**:386–398.
6. Huybrechts I, Matthys C, Vereecken C, Maes L, Temme EH, Van Oyen H, De Backer G, De Henauw S: Food intakes by preschool children in Flanders compared with dietary guidelines. *Int J Environ Res Public Health* 2008, **5**:243–257.
7. Ocke M, van Rossum C, Franses H, Buurma E, de Boer E, Brants H, Niekerk E, van der laan J, Drijvers J, Ghameshloou Z: *Dutch National Food Consumption Survey Young Children 2005/2006*. Bilthoven: RIVM; 2008. Report No.: RIVM rapport 350070001.
8. Van Ansem WJ, Schrijvers CT, Rodenburg G, van de Mheen D: Is there an association between the home food environment, the local food shopping environment and children's fruit and vegetable intake? Results from the Dutch INPACT study. *Public Health Nutr* 2013, **16**(7):1206–1214.
9. Vereecken C, Dupuy M, Rasmussen M, Kelly C, Nansel TR, Al Sabbah H, Baldassari D, Jordan MD, Maes L, Niclasen BV-L, Ahluwalia N: Breakfast consumption and its socio-demographic and lifestyle correlates in schoolchildren in 41 countries participating in the HBSC study. *Int J Public Health* 2009, **54**(Suppl 2):180–190.
10. Rampersaud GC, Pereira MA, Girard BL, Adams J, Metzler JD: Breakfast habits, nutritional status, body weight, and academic performance in children and adolescents. *J Am Diet Assoc* 2005, **105**:743–760. quiz 761–2.
11. Kelder SH, Perry CL, Klepp KI, Lytle LL: Longitudinal tracking of adolescent smoking, physical activity, and food choice behaviors. *Am J Public Health* 1994, **84**:1121–1126.
12. Craigie AM, Lake A, Kelly S, Adamson AJ, Mathers JC: Tracking of obesity-related behaviours from childhood to adulthood: a systematic review. *Maturitas* 2011, **70**:266–284.
13. Bere E, van Lenthe F, Klepp K-I, Brug J: Why do parents' education level and income affect the amount of fruits and vegetables adolescents eat? *Eur J Public Health* 2008, **18**:611–615.
14. Riediger ND, Shooshitari S, Moghadasian MH: The influence of sociodemographic factors on patterns of fruit and vegetable consumption in Canadian adolescents. *J Am Diet Assoc* 2007, **107**:1511–1518.
15. Hanson MD, Chen E: Socioeconomic status and health behaviors in adolescence: a review of the literature. *J Behav Med* 2007, **30**:263–285.
16. Rasmussen M, Krølner R, Klepp K-I, Lytle L, Brug J, Bere E, Due P: Determinants of fruit and vegetable consumption among children and adolescents: a review of the literature. Part I: Quantitative studies. *Int J Behav Nutr Phys Act* 2006, **3**:22.
17. Hilsen M, van Stralen MM, Klepp K-I, Bere E: Changes in 10–12 year old's fruit and vegetable intake in Norway from 2001 to 2008 in relation to gender and socioeconomic status - a comparison of two cross-sectional groups. *Int J Behav Nutr Phys Act* 2011, **8**:108.
18. Pearson N, Biddle SJH, Gorely T: Family correlates of breakfast consumption among children and adolescents. A systematic review. *Appetite* 2009, **52**:1–7.
19. Hallström L, Vereecken C, Ruiz JR, Patterson E, Gilbert CC, Catasta G, Díaz L-E, Gómez-Martínez S, González Gross M, Gottrand F, Hegyi A, Lehoux C, Mouratidou T, Widham K, Aström A, Moreno LA, Sjöström M: Breakfast habits and factors influencing food choices at breakfast in relation to socio-demographic and family factors among European adolescents. The HELENA Study. *Appetite* 2011, **56**:649–657.
20. Hallström L, Vereecken CA, Labayen I, Ruiz JR, Le Donne C, García MC, Gilbert CC, Martínez SG, Grammatikaki E, Huybrechts I, Kafatos A, Kersting M, Manios Y, Molnár D, Patterson E, Widham K, De Vriendt T, Moreno LA, Sjöström M: Breakfast habits among European adolescents and their association with sociodemographic factors: the HELENA (Healthy Lifestyle in Europe by Nutrition in Adolescence) study. *Public Health Nutr* 2012, **15**:1879–1889.
21. Tinsley B: *How Children Learn to Be Healthy*. Cambridge: Cambridge University Press; 2003.
22. Lindsay AC, Sussner KM, Kim J, Gortmaker S: The role of parents in preventing childhood obesity. *Future Child* 2006, **16**:169–186.
23. Krølner R, Rasmussen M, Brug J, Klepp K-I, Wind M, Due P: Determinants of fruit and vegetable consumption among children and adolescents: a review of the literature. Part II: qualitative studies. *Int J Behav Nutr Phys Act* 2011, **8**:112.
24. Pearson N, Biddle SJH, Gorely T: Family correlates of fruit and vegetable consumption in children and adolescents: a systematic review. *Public Health Nutr* 2009, **12**:267–283.
25. Winkleby MA, Jatulis DE, Frank E, Fortmann SP: Socioeconomic status and health: how education, income, and occupation contribute to risk factors for cardiovascular disease. *Am J Public Health* 1992, **82**:816–820.
26. Bogers RP, Van Assema P, Kester ADM, Westerterp KR, Dagnelie PC: Reproducibility, validity, and responsiveness to change of a short questionnaire for measuring fruit and vegetable intake. *Am J Epidemiol* 2004, **159**:900–909.
27. Haraldsdóttir J, Thórsdóttir I, de Almeida MDV, Maes L, Pérez Rodrigo C, Elmadfa F, Frost Andersen L: Validity and reproducibility of a precoded questionnaire to assess fruit and vegetable intake in European 11- to 12-year-old schoolchildren. *Ann Nutr Metab* 2005, **49**:221–227.
28. Dutman AE, Stafleu A, Kruijzinga A, Brants HAM, Westerterp KR, Kistemaker C, Meuling WJA, Goldbohm RA: Validation of an FFQ and options for data processing using the doubly labelled water method in children. *Public Health Nutr* 2011, **14**:410–417.
29. Brants H, Stafleu A, Ter Doest D, Hulshof K: Ontwikkeling van een voedsel frequentie vragenlijst: energie-inneming van kinderen van 2 tot en met 12 jaar. *Voeding Nu* 2006, **2**:25–28.
30. The Netherlands Nutrition Centre Foundation: *Richtlijnen Voedselkeuze (Dietary Guidelines)*. The Hague: 2011.
31. Van der Horst K, Oenema A, van de Looij-Jansen P, Brug J: The ENDORSE study: research into environmental determinants of obesity related behaviors in Rotterdam schoolchildren. *BMC Public Health* 2008, **8**:142.
32. Gattshall ML, Shoup JA, Marshall JA, Crane LA, Estabrooks PA: Validation of a survey instrument to assess home environments for physical activity and healthy eating in overweight children. *Int J Behav Nutr Phys Act* 2008, **5**:3.
33. Cole TJ, Bellizzi MC, Flegal KM, Dietz WH: Establishing a standard definition for child overweight and obesity worldwide: international survey. *BMJ* 2000, **320**:1240–1243.
34. Baron RM, Kenny DA: The moderator-mediator variable distinction in social psychological research: conceptual, strategic, and statistical considerations. *J Pers Soc Psychol* 1986, **51**:1173–1182.
35. Johansen A, Rasmussen S, Madsen M: Health behaviour among adolescents in Denmark: influence of school class and individual risk factors. *Scand J Public Health* 2006, **34**:32–40.
36. Sandvik C, Gjestad R, Samdal O, Brug J, Klepp K-I: Does socio-economic status moderate the associations between psychosocial predictors and fruit intake in schoolchildren? The Pro Children study. *Health Educ Res* 2010, **25**:121–134.
37. Neumark-Sztainer D, Wall M, Perry C, Story M: Correlates of fruit and vegetable intake among adolescents. *Prev Med (Baltim)* 2003, **37**:198–208.

38. MacFarlane A, Crawford D, Ball K, Savige G, Worsley A: **Adolescent home food environments and socioeconomic position.** *Asia Pac J Clin Nutr* 2007, **16**:748–756.
39. Campbell K, Crawford D, Jackson M, Cashel K, Worsley A, Gibbons K, Birch LL: **Family food environments of 5-6-year-old-children: does socioeconomic status make a difference?** *Asia Pac J Clin Nutr* 2002, **11**(Suppl 3):S553–S561.
40. Hupkens CL, Knibbe RA, Van Otterloo AH, Drop MJ: **Class differences in the food rules mothers impose on their children: a cross-national study.** *Soc Sci Med* 1998, **47**:1331–1339.
41. Vereecken C, Keukelier E, Maes L: **Influence of mother's educational level on food parenting practices and food habits of young children.** *Appetite* 2004, **43**:93–103.
42. Campbell KJ, Abbott G, Spence AC, Crawford DA, McNaughton SA, Ball K: **Home food availability mediates associations between mothers' nutrition knowledge and child diet.** *Appetite* 2013, **71**:1–6.
43. Hendrie GA, Brindal E, Corsini N, Gardner C, Baird D, Golley RK: **Combined home and school obesity prevention interventions for children: what behavior change strategies and intervention characteristics are associated with effectiveness?** *Health Educ Behav* 2012, **39**:159–171.
44. Tak NI, te Velde SJ, de Vries JHM, Brug J: **Parent and child reports of fruit and vegetable intakes and related family environmental factors show low levels of agreement.** *J Hum Nutr Diet* 2006, **19**:275–285.
45. Matthys C, De Henauw S, Bellemans M, De Maeyer M, De Backer G: **Breakfast habits affect overall nutrient profiles in adolescents.** *Public Health Nutr* 2007, **10**:413–421.

doi:10.1186/s12966-014-0113-0

**Cite this article as:** van Ansem *et al.*: Maternal educational level and children's healthy eating behaviour: role of the home food environment (cross-sectional results from the INPACT study). *International Journal of Behavioral Nutrition and Physical Activity* 2014 **11**:113.

**Submit your next manuscript to BioMed Central and take full advantage of:**

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at  
[www.biomedcentral.com/submit](http://www.biomedcentral.com/submit)

