

# Guillain-Barré Syndrome and Adjuvanted Pandemic Influenza A (H1N1) 2009 Vaccines: A Multinational Self-Controlled Case Series in Europe

Silvana Romio<sup>1,9</sup>, Daniel Weibel<sup>1,16\*</sup>, Jeanne P. Dieleman<sup>1</sup>, Henning K. Olberg<sup>2</sup>, Corinne S. de Vries<sup>3</sup>, Cormac Sammon<sup>3</sup>, Nick Andrews<sup>4</sup>, Henrik Svanström<sup>5</sup>, Ditte Mølgaard-Nielsen<sup>5</sup>, Anders Hviid<sup>5</sup>, Maryse Lapeyre-Mestre<sup>6</sup>, Agnès Sommet<sup>6</sup>, Christel Saussier<sup>7</sup>, Anne Castot<sup>8</sup>, Harald Heijbel<sup>9</sup>, Lisen Arnheim-Dahlström<sup>10</sup>, Par Sparen<sup>10</sup>, Mees Mosseveld<sup>1</sup>, Martijn Schuemie<sup>1</sup>, Nicoline van der Maas<sup>11</sup>, Bart C. Jacobs<sup>12</sup>, Tuija Leino<sup>13</sup>, Terhi Kilpi<sup>13</sup>, Jann Storsaeter<sup>14</sup>, Kari Johansen<sup>15</sup>, Piotr Kramarz<sup>15</sup>, Jan Bonhoeffer<sup>16,17</sup>, Miriam C. J. M. Sturkenboom<sup>1</sup>

**1** Department of Medical Informatics, Erasmus University Medical Center, Rotterdam, The Netherlands, **2** Department of Neurology, Haukeland University Hospital, Bergen, Norway, **3** Department of Pharmacy & Pharmacology, University of Bath, Bath, United Kingdom, **4** Health Protection Agency, London, United Kingdom, **5** Department of Epidemiology Research, Statens Serum Institut, Copenhagen, Denmark, **6** Department of Clinical Pharmacology, Toulouse University Hospital and Institut national de la santé et de la recherche médicale, Toulouse, France, **7** French National Agency for Medicines and Health Products Safety, Saint Denis, France, **8** Medical Doctor - Consultant, Paris, France, **9** Department of Vaccinology, Swedish Institute for Infectious Disease Control, Solna, Sweden, **10** Department Medical Epidemiology and Biostatistics, Karolinska Institute, Stockholm, Sweden, **11** National Institute for Public Health and the Environment, Bilthoven, The Netherlands, **12** Departments of Neurology and Immunology, Erasmus University Medical Center, Rotterdam, The Netherlands, **13** Department of Vaccination and Immune Protection, National Institute for Health and Welfare, Helsinki, Finland, **14** Department of Vaccines, Norwegian Institute of Public Health, Oslo, Norway, **15** Surveillance and Response Support Unit, European Centre for Disease Prevention and Control, ECDC, Stockholm, Sweden, **16** Brighton Collaboration Foundation, Basel, Switzerland, **17** Department of Infectious Diseases and Vaccinology, University Children's Hospital, Basel, Switzerland

## Abstract

**Background:** The risk of Guillain-Barré syndrome (GBS) following the United States' 1976 swine flu vaccination campaign in the USA led to enhanced active surveillance during the pandemic influenza (A(H1N1)pdm09) immunization campaign. This study aimed to estimate the risk of GBS following influenza A(H1N1)pdm09 vaccination.

**Methods:** A self-controlled case series (SCCS) analysis was performed in Denmark, Finland, France, Netherlands, Norway, Sweden, and the United Kingdom. Information was collected according to a common protocol and standardised procedures. Cases classified at levels 1–4a of the Brighton Collaboration case definition were included. The risk window was 42 days starting the day after vaccination. Conditional Poisson regression and pooled random effects models estimated adjusted relative incidences (RI). Pseudo likelihood and vaccinated-only methods addressed the potential contraindication for vaccination following GBS.

**Results:** Three hundred and three (303) GBS and Miller Fisher syndrome cases were included. Ninety-nine (99) were exposed to A(H1N1)pdm09 vaccination, which was most frequently adjuvanted (Pandemrix and Focetria). The unadjusted pooled RI for A(H1N1)pdm09 vaccination and GBS was 3.5 (95% Confidence Interval (CI): 2.2–5.5), based on all countries. This lowered to 2.0 (95% CI: 1.2–3.1) after adjustment for calendartime and to 1.9 (95% CI: 1.1–3.2) when we accounted for contraindications. In a subset (Netherlands, Norway, and United Kingdom) we further adjusted for other confounders and there the RI decreased from 1.7 (adjusted for calendar month) to 1.4 (95% CI: 0.7–2.8), which is the main finding.

**Conclusion:** This study illustrates the potential of conducting European collaborative vaccine safety studies. The main, fully adjusted analysis, showed that the RI of GBS was not significantly elevated after influenza A(H1N1)pdm09 vaccination (RI = 1.4 (95% CI: 0.7–2.8)). Based on the upper limits of the pooled estimate we can rule out with 95% certainty that the number of excess GBS cases after influenza A(H1N1)pdm09 vaccination would be more than 3 per million vaccinated.

**Citation:** Romio S, Weibel D, Dieleman JP, Olberg HK, de Vries CS, et al. (2014) Guillain-Barré Syndrome and Adjuvanted Pandemic Influenza A (H1N1) 2009 Vaccines: A Multinational Self-Controlled Case Series in Europe. PLoS ONE 9(1): e82222. doi:10.1371/journal.pone.0082222

**Editor:** Nicole M. Bouvier, Mount Sinai School of Medicine, United States of America

**Received:** May 22, 2013; **Accepted:** October 24, 2013; **Published:** January 3, 2014

**Copyright:** © 2014 Romio et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

**Funding:** The study was funded by European Centre for Disease Prevention and Control (ECDC) and co-funded by all participating centers from Denmark, Finland, France, Netherlands, Norway, Sweden, and the United Kingdom as specified in the Authors' affiliations. Funders have been involved in designing the study, collecting the data, analysis, decisions to publish, and in reviewing the manuscript.

**Competing Interests:** Silvana Romio, Daniel Weibel, Henning K Olberg, Nick Andrews, Henrik Svanström, Ditte Mølgaard-Nielsen, Anders Hviid, Maryse Lapeyre, Agnès Sommet, Christel Saussier, Anne Castot, Harald Heijbel, Lisen Arnheim-Dahlström, Mees Mosseveld, Noline van der Maas, Bart C Jacobs, Tuija Leino, Jann Storsaeter, Kari Johansen, and Piotr Kramarz have no conflicts of interest. Jeanne P Dieleman has been involved in studies for pharmaceutical companies (i.e., GSK, Sanofi, Astra-Zeneca, Pfizer). None of these had any conflict with the present study. The affiliation (i.e. University of Bath) of Corinne S de Vries and Cormac Sammon has research and consulting contracts in place with Novartis vaccines and with GSK pharmaceuticals. The authors do not personally benefit from these contracts; all financial compensation is to the University of Bath and not to the authors. Martijn Schuemie was employed at the Erasmus University Medical Center, Rotterdam at the time of the study. Since January 1st 2013 (after completion of the study), he is employed at Janssen R&D. Terhi Kilpi is a principal investigator of a nationwide Finnish effectiveness study of the 10-valent pneumococcal conjugate vaccine, a collaborative study, for which her institute has received funding from GSK. Par Sparen received a grant from Glaxo Smith Kline in 2010 to for a retrospective, observational register based cohort study to evaluate the safety of GSK Biological's H1N1 pandemic vaccine administered in Sweden according to local vaccination policy. Miriam CJM Sturkenboom: is head of a research group that occasionally conducts research for pharmaceutical companies including Pfizer, Eli Lilly, Boehringer, AstraZeneca and Novartis. None was related to this topic. This does not alter our adherence to all the PLOS ONE policies on sharing data and materials.

\* E-mail: d.weibel@erasmusmc.nl

These authors contributed equally to this work.

## Introduction

During the influenza A (H1N1) 2009 pandemic, new monovalent adjuvanted and non-adjuvanted influenza A(H1N1)pdm09 vaccines were introduced in Europe. Immunogenicity and safety was in line with the “Committee for medicinal products for human use (CHMP) Note for Guidance”, but safety data were limited [1–3]. Vaccination campaigns started in autumn 2009 at the peak of the pandemic in Europe.

A key safety concern identified in planning the pandemic vaccination campaigns was the potential association between Guillain-Barré syndrome (GBS) and influenza vaccines; this concern stemmed from an association observed in the USA in 1976 between swine flu vaccination and GBS [4]. Subsequent prospective surveillance studies and retrospective epidemiological studies on seasonal influenza vaccines used in 1978, 1992, 1993, and beyond showed no or modest increases in the risk of GBS [5–8]. Despite this, the US Food and Drug Administration (FDA), the World Health Organization (WHO) and the European Medicines Agency (EMA) recommended active monitoring of a potential association between the influenza A(H1N1)pdm09 vaccine and GBS.

In Europe, GBS primarily presents as an acute inflammatory demyelinating polyradiculoneuropathy (AIDP) [9]. Three to ten per cent of GBS patients die and an estimated 20% experience continued disability for more than six months [10]. Prospective studies in developed countries have estimated an incidence rate of 2 per 100,000 population per year with an increased risk with age and in males [11]. GBS is thought to be primarily triggered by a preceding respiratory or gastrointestinal infection [12].

The European Centre for Disease prevention and Control (ECDC) commissioned the VAESCO (Vaccine Adverse Events Surveillance and Communication) consortium to study the potential association between influenza A(H1N1)pdm09 vaccine and GBS. A case control study was conducted for a rapid initial assessment with a large-scale more extensive prospective SCCS study carried out in parallel. The VAESCO case control study was based on 104 cases in five European countries and showed no association between A(H1N1)pdm09 vaccine (mostly adjuvanted with AS03) and GBS [13]. In this paper we present the results from the VAESCO SCCS study which included three times the amount of cases.

## Methods

### Setting and design

The VAESCO consortium conducted a prospective self-controlled case series (SCCS) study to investigate the association between influenza A(H1N1)pdm09 vaccination and GBS. A SCCS is a case-only study comparing the incidence of disease

during risk and non-risk periods within the same person, inherently controlling for measured and unmeasured confounding factors that remain stable over time [14].

The VAESCO consortium was initiated and core funded by ECDC with the aim of improving post licensure vaccine safety in Europe. It is coordinated by the Brighton Collaboration Foundation and includes partners from public health organizations, regulatory authorities and academic research institutions in Europe.

Centers from Denmark (DK), Finland (FI), France (FR), Netherlands (NL), Norway (NO), Sweden (SE), and the United Kingdom (UK) contributed to the study. All centers used a common protocol and applied the standardised Brighton Collaboration GBS case definition for case classification [9]. Implementation of the protocol and data collection differed per country based on ethical requirements and the healthcare structure. Data harmonization, transformation, and pooling used methods and tools derived from the EU-ADR (Exploring and Understanding Adverse Drug Reactions) project [15]. Centers created harmonized input files according to well-defined instructions. These data files were generated directly from automated resources or manually using customized electronic case report forms. The harmonized input files were transformed using a standardized JAVA-based program (Jerboa<sup>®</sup> version 2.6.0, September 2010, Erasmus University Medical Center, Rotterdam, Netherlands). Only anonymous and aggregated de-identified information without dates of disease or exposure were shared for individual patient level data pooling and centralised analysis. Consent forms, original data and Jerboa input files were retained at the local centers. Quality control and verification of transmitted data was done at the central data management and analysis center (Erasmus University) in close collaboration with the other centers. All centers commented on the data and results prior to release.

### Source and study population

The total source population exceeded 50 million (M) subjects, with most countries recruiting cases on a national level (NO (4.4 M), SE (9 M), FI (5.5 M), DK (5 M), NL (16 M)). In the UK, the General Practice Research Database (GPRD) (5 M) was used and in France specialized hospitals with a large but undefined catchment area participated. Case recruitment started on 1st November 2009 and lasted maximally until 1st November 2010.

The study population encompassed all cases with GBS or its variant Miller Fisher syndrome with onset of disease during the study period.

Case recruitment-procedures are described in Table 1. Completeness of recruitment was verified retrospectively at the end of the study period by comparing recruited cases with diagnosed case lists (see Table 1). Additional cases identified in this way were

**Table 1.** Sources of cases, exposure and covariate information per country.

	Cases recruitment	Exposure Information	Covariates during follow-up	Potential bias
<b>DK</b>	Cases were identified from the National Patient Register using primary discharge diagnoses only (ICD-10: G61.0). Case validation based on retrospective chart review.	Vaccination registry	None (only from case hospital charts)	Cases: not all charts available No ability to control for time varying confounders
<b>FI</b>	From hospital Discharge and hospital outpatient records, primary diagnoses (ICD-10 G61.0). Case validation based on retrospective chart review	Vaccination registry	None (only from case hospital charts)	Cases: not all charts available No ability to control for time varying confounders
<b>FR</b>	Cases were identified prospectively through neurologists in 7 reference hospitals in FR. Patients needed to provide informed consent. Completeness was verified against pharmacy data (immunoglobulin prescriptions) and showed incomplete reporting (<50%), Vaccination status of non-reported cases could not be verified since linkage to vaccination registry required consent.	Ad hoc A(H1N1)pdm09 vaccination registry	Hospital charts and interview, only for period prior to GBS	Incompleteness and potential selection bias cannot be excluded. No ability to control for time varying confounders
<b>NL</b>	Cases were identified prospectively through neurologists. Completeness was verified retrospectively by checking against the claims codes in each of the reporting hospitals. Missing patients were included retrospectively in hospitals that were reporting at least one case prospectively.	GP medical record	GP medical record	Small potential for misclassification of exposure since A(H1N1)pdm09 vaccination could also be provided through public health agency for parents of young children
<b>NO</b>	Nationwide neurologist reporting network, group of neurologists. Case validation based on review of GBS experts	Vaccination registry	Neurologists, Hospitals, and GPs	Potential selection due to incompleteness Information on co-variables collected differently for period prior to GBS.
<b>SE</b>	Cases of GBS were identified through seven neurology assessment labs where GBS cases are laboratory confirmed for a population of 9.4 million. Informed consent needed to be obtained from all cases. Completeness of cases was checked in the National Patient Registry for part of the country. Recruitment was incomplete because of delays in consent and non-consent. It was not possible to assess whether this non-response differed by vaccination status and hence selection bias cannot be excluded.	By interview at end of follow-up, recall bias cannot be excluded.	By interview for cases at the end of follow up. change in region over time. Should not be used for adjustment	Consent required, potential selection bias. Recall bias (differential recall over time)
<b>UK</b>	Each case was identified in the General Practice Research Database by using appropriate READ codes (F370.00, F370000, F370100, F370200, F370z00). Case verification was done using any hospital letters, discharge summaries and GPs' notes recorded as free text. No major selection to be expected	Automated GP records, no recall bias. Non-differential misclassification possible since some persons might have been vaccinated outside of GP office.	GP records	Misclassification of cases due to lack of information on test results

doi:10.1371/journal.pone.0082222.t001

included retrospectively where possible. For each subject, follow-up started at the beginning of the study period or date of birth if born after the start of the study period. Follow-up ended with the end of the study period or death occurring prior to the end of the study period.

The earliest date of onset of neurological symptoms was the index date. If the date of first symptoms could not be retrieved the date of diagnosis or hospitalization was used. Informed consent was required in SE and FR. Case characteristics were obtained from neurologists or from discharge letters and used to classify cases according to the Brighton Collaboration GBS Case Classification using the Automated Brighton Classification (ABC) tool ([www.brightoncollaboration.org](http://www.brightoncollaboration.org)).


### Vaccine Exposure

The primary exposure of interest was vaccination with adjuvanted or non-adjuvanted A(H1N1)pdm09 vaccine as recorded in vaccination registries (FR, DK, FI, NO), General Practitioners' (GP) records (NL, UK), or patient interview (SE). The risk period began the day after vaccination and ended 42 days later. If two doses were administered, the risk period of the first dose ended

when the second dose was administered. Brand specific information was collected for each influenza A(H1N1)pdm09 vaccination.

### Covariates

Information on several time varying risk factors for GBS was collected during follow-up including seasonal influenza vaccination, influenza-like illness (ILI), upper respiratory tract infections (URTI), and gastrointestinal infections (GI). Each of these covariates was assigned a 42-day risk period. The risk period began on day one of onset of ILI, URTI, or GI or of seasonal influenza vaccine receipt and ended 42 days after onset or exposure. Covariate data were not collected in DK and FI. In FR, covariate data were collected from neurologists at case occurrence for the period prior to GBS only, whereas in SE data on covariates were collected by interview at the end of follow-up. In the UK, NL, and NO general practitioner records were used to collect information on covariates throughout the follow up period; NO also assessed covariates reported by neurologists at the time of case data collection, leading to a potential for differential data collection over time. To adjust for seasonal effects, changes in circulation of the wild type influenza A(H1N1)pdm09 virus and




**Figure 1. Flowchart of case inclusion.**  
doi:10.1371/journal.pone.0082222.g001

differences in case inclusion over the observation period we considered calendar month as a time varying covariate.

### Statistical Analysis


The RI for the association between A(H1N1)pdm09 vaccine and GBS was estimated using a conditional Poisson regression analysis. This was done for each country separately. Adjustment for calendar month was possible in all countries, whereas further adjustment for ILI, URTI, GI, and seasonal influenza vaccination was only possible in NL, UK, and NO. Sensitivity analyses were

used to assess the effects of misclassification of exposure and confounding. An analysis using vaccinated cases only and an analysis using the pseudo-likelihood approach explored confounding by contra-indication to influenza A(H1N1)pdm09 vaccination [14]. A sub analysis was done to assess the impact of residual confounding by ILI, URTI, seasonal influenza vaccination, and GI infections. Misclassification of the risk period was investigated by applying risk periods smaller than 42 days. In order to study effect modification by infections occurring just prior to GBS onset, stratified analyses were carried out for age, sex, history of GBS,



**Figure 2. Inclusion of GBS cases (DK, FI, FR, NL, NO, SE, UK), influenza A(H1N1)pdm09 immunization period (influenza A(H1N1)vac), and percentage of flu positive cases among all tested per country (Flu pos. DK, ..., Flu pos. UK; Source: ECDC 2011) over total study period.**

doi:10.1371/journal.pone.0082222.g002



**Figure 3. Inclusion of vaccinated cases (% of population) over study period.**  
doi:10.1371/journal.pone.0082222.g003

and prior infections (ILI, URTI, GI) in UK, NL, and NO. The country specific estimates were pooled applying a random effects model. All analysis used SAS v9.1 (Cary, North Carolina).

## Results

In total 730 potential GBS cases were identified during the study period. Of these, 427 cases were excluded (see figure 1), leaving 303 GBS cases in the study population. Case inclusion declined over time from 133 cases in the first three months to 18 in the last three months (Figure 2). The percentage of influenza A (H1N1) pdm09 vaccinated cases did not change significantly over time ( $R^2 = 0.094$ ; Figure 3).

Cases had a mean age of 50 years (SD: 4.1) ranging from 45 (SD: 20.8) years in the NL to 56 (SD: 19.5) years in NO, less than 10% were younger than 20 years. On average the follow-up period was 321 days. Case classification differed by country, primarily depending on the type of data source used for case recruitment. Of all cases, 36% were classified as Brighton Collaboration level 1, 26% as level 2, 13% as level 3, and 25% as level 4a. In 69 cases electrophysiology (mostly AIDP) had either not been performed for diagnosis or was not recorded. On a scale from 0 to 6, with 0 meaning complete physical fitness and 6 meaning death, the disability score was most frequently 4 (30.6%) (Table 2).

Overall, 99 cases (33%) received influenza A(H1N1)pdm09 vaccination, mostly adjuvanted with AS03, before symptom onset (Table 3). Of these, 36 (37%) cases developed GBS within 42 days after a first dose of influenza A(H1N1)pdm09 vaccination whereas 7 cases occurred within the exposure risk window but after a second dose of influenza A(H1N1)pdm09 vaccination.

Few countries could collect data on time-varying covariates over the entire follow-up period. Most countries assessed covariates at the time of case collection, but not afterwards, and therefore these data could not be utilized for adjustments but could be used for stratification. Based on the information collected at case occurrence, 15 cases developed GBS within 42 days after seasonal influenza vaccination and 79 cases developed GBS within 42 days after onset of ILI or URTI (Table 3).

## Risk ratio of GBS

The crude country specific RI of GBS during the influenza A(H1N1)pdm09 vaccination risk period compared to the non-risk period varied from a low of 1.6 in FI to a high of 7.7 in DK (based

on two exposed cases only), with an overall pooled estimate of 3.5 (95% CI: 2.2 to 5.5). Adjustment for calendar month had a significant impact (RI: 2.0, 95% CI: 1.2 to 3.1). Sensitivity analyses accounting for contra-indication after GBS onset showed a minor change in the calendar month adjusted pooled RI from 2.0 to 1.9 (95% CI: 1.1 to 3.2) when the pseudolikelihood method was used, and 1.8 (95% CI: 0.7 to 4.7) when considering vaccinated cases only (Table 4).

In NL, NO, and the UK where further adjustment for infections, seasonal influenza vaccination, and other time dependent covariates was possible, the RI for the association between influenza A(H1N1)pdm09 vaccination and GBS decreased from the unadjusted pooled RI of 3.2 (95% CI: 1.8 to 5.6) to 1.7 (95% CI: 0.8 to 3.4) after adjustment for calendar month, and to 1.4 (95% CI: 0.7 to 2.8) upon further adjustment for ILI, URTI, and GI.

Sensitivity analyses using different post-exposure risk periods resulted in a calendar month-adjusted pooled RI of 2.3 (95% CI: 1.4 to 3.8) for the first four weeks. The RI was 2.3 (95% CI: 1.2 to 4.4) in the first two weeks and 2.6 (95% CI 1.4 to 4.9) during weeks three to four.

We did not observe statistically significant interactions between age, infections, or seasonal influenza vaccination and the association between the influenza A(H1N1)pdm09 vaccination and GBS (Table 5).

## Discussion

Based on a source population of more than 25 million subjects from NL, UK, and NO we found no significant elevated association between the risk of GBS following immunization with an adjuvanted influenza A(H1N1)pdm09 vaccine, when adjusted for all known measurable confounders (RI 1.4, 95% CI: 0.7 to 2.8). This result is very similar to that of the VAESCO consortium case control study, published previously using one third of the cases from fewer countries [13]. In DK, FI, FR and SE we could not adjust for time varying confounders such as infections since data were not collected over the entire follow up period. Pooling data from all seven countries yielded a crude RI of 3.5, which reduced to 2.0 (95% CI: 1.2 to 3.1) after adjustment for calendar month: this pooled estimate still comprises residual confounding by infections. The effect of calendar month may be explained by it being a good proxy for circulation of the wild-type influenza A(H1N1)pdm09 virus (see figure 2).



**Table 2.** Characteristics of Guillain-Barré syndrome cases.

Characteristic	DK		FI		FR		NL		NO		SE		UK		Total	
	N	%	N	%	N	%	N	%	N	%	N	%	N	%	n	%
Cases in study period	31	10.2	29	9.6	41	13.5	80	26.4	50	16.5	32	10.6	40	13.2	303	100
Females	14	45.2	12	41.4	20	48.8	32	40.0	25	50.0	12	37.5	17	42.5	132	43.6
<b>Mean age (SD)<sup>1</sup> [years]</b>	49.2 (20.2)		54.4 (20.8)		50.0 (21.9)		45.0 (20.8)		55.5 (19.5)		51.5 (20.2)		45.4 (20.4)		50.1 (4.1)	
Age ≤4	0	0.0	0	0.0	1	2.4	2	2.5	1	2.0	0	0.0	2	5.0	6	2.0
Age 5–19 years	3	9.7	3	10.3	4	9.8	10	12.5	0	0.0	2	6.3	3	7.5	25	8.3
Age 20–59 years	18	58.1	10	34.5	18	43.9	44	55.0	21	42.0	15	46.9	24	60.0	15	49.5
Age ≥60	10	32.3	16	55.2	18	43.9	24	30.0	28	56.0	15	46.9	11	27.5	122	40.3
<b>Brighton Classification<sup>2</sup></b>																
1	10	32.3	17	58.6	13	31.7	28	35.0	21	42.0	19	59.4	0	0.0	108	35.6
2	8	25.8	3	10.3	16	39.0	30	37.5	14	28.0	8	25.0	0	0.0	79	26.1
3	4	12.9	7	24.1	7	17.1	11	13.8	5	10.0	5	15.6	0	0.0	39	12.9
4a	9	29.0	2	6.9	5	12.2	10	12.5	10	20.0	0	0.0	40	100.0	76	25.1
Unknown	0	0.0	0	0.0	0	0.0	1	1.3	0	0.0	0	0.0	0	0.0	1	0.3
<b>Electrophysiology</b>																
AIDP <sup>3</sup>	23	74.2	16	55.2	15	36.6	36	45.0	29	58.0	23	71.9	0	0.0	142	46.9
AMAN <sup>4</sup>	0	0.0	0	0.0	1	2.4	6	7.5	3	6.0	0	0.0	0	0.0	10	3.3
AMSAN <sup>5</sup>	1	3.2	2	6.9	0	0.0	4	5.0	1	2.0	6	18.8	2	5.0	16	5.3
Equivocal	0	0.0	0	0.0	6	14.6	9	11.3	7	14.0	1	3.1	0	0.0	23	7.6
Normal	2	6.5	0	0.0	2	4.9	3	3.8	6	12.0	0	0.0	0	0.0	13	4.3
Not performed	5	16.1	11	37.9	12	29.3	20	25.0	3	6.0	2	6.3	0	0.0	53	17.5
Unresponsive nerves	0	0.0	0	0.0	0	0.0	1	1.3	1	2.0	0	0.0	0	0.0	2	0.7
Unknown	0	0.0	0	0.0	5	12.2	1	1.3	0	0.0	0	0.0	38	95.0	44	14.5
<b>GBS disability score<sup>6</sup></b>																
0	0	0	0	0.0	6	14.6	0	0.0	0	0.0	0	0.0	0	0	6	2.0
1	0	0	0	0.0	0	0.0	5	6.3	7	14.0	1	3.1	0	0	13	4.3
2	0	0	9	31.0	2	4.9	19	23.8	11	22.0	6	18.6	0	0	47	15.5
3	0	0	4	13.8	10	24.4	21	26.3	6	12.0	7	21.9	0	0	48	15.8
4	0	0	13	44.8	11	26.8	20	25.0	17	34.0	10	31.3	0	0	71	23.4
5	0	0	2	6.9	2	4.9	13	16.3	8	16.0	7	21.9	0	0	32	10.6
6	0	0	1	3.4	0	0.0	1	1.3	1	2.0	1	3.1	0	0	4	1.3
Unknown	31	100.0	0	0.0	10	24.4	1	1.3	0	0.0	0	0.0	40	100.0	82	27.1
<b>Index month</b>																
Nov 2009	9	29.0	4	13.8	9	22.0	5	6.3	5	10.0	8	25.0	6	15.0	46	15.2
Dec 2009	1	3.2	5	17.2	8	19.5	22	27.5	12	24.0	6	18.8	4	10.0	58	19.1
Jan 2010	6	19.4	5	17.2	9	22.0	8	10.0	8	16.0	3	9.4	5	12.5	44	14.5
Feb 2010	5	16.1	6	20.7	6	14.6	9	11.3	1	2.0	5	15.6	8	20.0	40	13.2
Mar 2010	4	12.9	3	10.3	3	7.3	5	6.3	4	8.0	3	9.4	6	15.0	28	9.2
Apr 2010	3	9.7	5	17.2	4	9.8	7	8.8	4	8.0	2	6.3	3	7.5	28	9.2
May 2010	3	9.7	1	3.4	1	2.4	8	10.0	3	6.0	2	6.3	1	2.5	19	6.3
Jun 2010	0	0.0	0	0.0	1	2.4	2	2.5	3	6.0	3	9.4	3	7.5	12	4.0
Jul 2010	0	0.0	0	0.0	0	0.0	3	3.8	3	6.0	0	0.0	1	2.5	7	2.3
Aug 2010	0	0.0	0	0.0	0	0.0	4	5.0	2	4.0	0	0.0	2	5.0	8	2.6
Sep 2010	0	0.0	0	0.0	0	0.0	4	5.0	4	8.0	0	0.0	0	0.0	8	2.6
Oct 2010	0	0.0	0	0.0	0	0.0	3	3.8	1	2.0	0	0.0	1	2.5	5	1.7
TOTAL	31		29		41		80		50		32		40		303	

<sup>1</sup>Standard Deviation.<sup>2</sup>Sejvar J. J. et al. 2011, Guillain-Barre syndrome and Fisher syndrome: case definitions and guidelines for collection, analysis, and presentation of immunization safety data. Vaccine 29(3).<sup>3</sup>AIDP: acute inflammatory demyelinating polyradiculoneuropathy.

**Table 2.** Cont.<sup>4</sup>AMAN: acute motor axonal neuropathy.<sup>5</sup>AMSAN: acute motor and sensory axonal neuropathy.<sup>6</sup>Current disability score at the time of case assessment and/or inclusion into the study.

Abbreviations: DK: Denmark; FI: Finland; FR: France; NL: Netherlands; NO: Norway; SE: Sweden; UK: United Kingdom; GBS: Guillain-Barré syndrome.

doi:10.1371/journal.pone.0082222.t002

This study is unique as it directly pools data on individual patients from seven European countries, using a common protocol, common case definition, common infrastructure, and common data elaboration. The impact of methodological issues that occurred due to differences in implementation of the protocol could be assessed by comparing the association across countries; the consistency observed is reassuring. Beyond the effect of the influenza A(H1N1)pdm09 vaccination on GBS this study underlines the advantages of collaborative transnational vaccine safety studies. They not only increase the scale of a study, but also allow for consistency-checks across sources in the absence of bias from differences in design and methods. The use of common methods and subsequent pooling reaches far beyond the traditional approach of meta-analyses where rather heterogeneous estimates resulting from different designs, methods, and settings are being pooled.

The data from this VAESCO study are in line with other results from Europe with studies from FR (RI 0.9, 95% CI: 0.1 to 7.6) [16], SE (RI 1.1 95% CI: 0.6 to 1.9) [17], and the UK (RI 1.05, 95% CI: 0.37–2.24) [18], all showing no association. In contrast, a recent report from Germany, where AS03 adjuvanted vaccine was used, showed an increased risk of GBS after vaccination (RI 4.65, 95% CI: 2.17 to 9.98) [19]. German investigators had already started a separate SCCS study and thus elected not to participate in VAESCO. They did not adjust for infections or calendar-time and selection bias could not be excluded since cases originated from a reporting network. Pooling of calendar-month adjusted RI estimates with the VAESCO study would be possible through meta-analysis to enlarge the scale of the current EU based study. Five studies from the US, where non-adjuvanted influenza A(H1N1)pdm09 vaccines were used, have recently been published. Each of the initial observational studies found an increased RI ranging from 1.6 (95% CI: 1.0 to 2.2) [20], to 2.1 (95% CI: 1.2 to 3.5) [21], to 2.5 (95% CI: 0.42 to 15.0) [22], and to 4.4 (95% CI: 1.3 to 14.2) [23]. Three studies used self-controlled designs but without further adjusting for time-varying confounders [21–23]. The study assessing the lowest RI (1.6 (95% CI: 1.0 to 2.2)) was a cohort study adjusting for age and sex [20]. The highest RI of 4.4 (95% CI: 1.3 to 14.2) was based on data from the US Vaccine Safety Datalink (VSD) project, which was based on 13 vaccinated cases [23]. Salmon et al. recently published a meta-analysis of US studies on the association between influenza A(H1N1)pdm09 vaccines including two unpublished studies and reported a pooled estimate of 2.35 (95% CI: 1.42–4.01) [24]. A SCCS study from Quebec, Canada adjusted for seasonality and contraindication using vaccinated cases only reported a relative risk of 1.9 (95% CI: 1.0 to 3.5) [25]. After the first VSD study, a second VSD study was recently published, investigating specifically the effect of antecedent infections on the relative incidence of GBS following influenza A(H1N1)pdm09 vaccines, using a case centered analysis. This analysis showed the impact of infections as a confounding factor [26]. After adjusting for antecedent infections, there was no evidence for an elevated GBS risk following 2009–10 monovalent/2010–11 trivalent influenza vaccines. However, the association between GBS and antecedent infection was strongly elevated. The

effect of infections on the risk of GBS and the potential preventive effect of vaccination on the risk of GBS by preventing influenza was recently discussed by Stowe and Poland [27,28]. This recent evidence underlines the need to adjust for infections as we could do in part of the countries in our analyses.

Owing to its observational nature, our study suffers from limitations that should be considered when interpreting data. In NL and SE, where reporting networks were used, completeness of recruitment was verified by retrospectively comparing included cases with claims made for GBS. In FR and SE informed consent was required which could be another reason for non-inclusion. Finally, since cases were included only if charts/medical records could be reviewed, lack of data could be another source of selection bias. The distribution of vaccinated cases over time showed no significant trend, suggesting changes in the number of cases included over time were not related to exposure and selection bias may be limited (Figure 3).

Information bias may arise from misclassification of the outcome as well as the exposure. Cases recruited directly from neurologists (i.e., FR, NL, NO, and SE) generally had higher levels of diagnostic certainty. In the UK all cases were classified with the lowest Brighton Collaboration case certainty level as information was retrieved retrospectively from GP medical records, which capture information from specialist letters but often lack information on specific test results. In DK cases were classified based on retrospective review of specialist charts resulting in partially missing information. As standardized criteria were used for case classification, misclassification of the outcome will be minimal. In all countries prospectively collected health care records were used to obtain information on exposure, except in SE, which relied on interviews and may have suffered from recall bias. In the NL exposure may have been misclassified in young children (<5 years) who were participating in mass vaccination campaigns, but this will be non-differential and there were very few paediatric cases. Exposure might be misclassified due to misspecification of the risk period. Sensitivity analysis showed no difference in the RI when the risk window was restricted to 15 to 28 days after vaccination (RI 2.6, 95% CI: 1.4 to 4.9); compared to the first two weeks (RI 2.3, 95% CI 1.2 to 4.4) and the risk in a 4-week risk window (RI 2.3, 95% CI 1.4 to 3.8).

We addressed confounding both by design (SCCS controls for time-constant confounders), through adjustments, and sensitivity analyses. GBS could be a contra-indication for influenza A(H1N1)pdm09 vaccine as a similar vaccine had been associated with GBS in the past. To investigate this issue we carried out analyses including only vaccinated subjects and analyses applying the pseudo-likelihood method [14]. The pseudo-likelihood method reduced the calendar-adjusted pooled RI from 2.0 to 1.9 and if only vaccinated cases were included to 1.8, indicating that contra-indications were a minor issue. Calendar month acted as an important confounding factor, not because time itself is a risk factor, but because it may serve as a proxy for influenza A(H1N1)pdm09 circulation, which was highly time-dependent and co-occurring with the mass vaccination campaigns (see figure 2). Adjustment for additional time-varying confounders

**Table 3.** Guillain-Barré syndrome occurrence during follow-up and during the 6-week (42 days) risk periods following influenza A(H1N1)pdm09 vaccination and infection.

Exposure	DK		FI		FR		NL		NO		SE		UK		TOTAL	
	n	%	n	%	n	%	n	%	n	%	n	%	n	%	n	%
<b>Cases in study period</b>	31	10.2	29	9.6	41	13.5	80	26.4	50	16.5	32	10.6	40	13.2	303	100
<b>Follow-up total (mean) in days</b>	11,286 (364.1)		6,127 (211.3)		11,421 (278.6)		26,322 (329)		17,845 (356.9)		11,471 (358.5)		12,666 (316.7)		97,138 (320.6)	
<b>Cases exposed to influenza A(H1N1)pdm09 vaccine anytime during follow-up period</b>	4	12.9	13	44.8	5	12.2	29	36.3	23	46.0	22	68.8	3	7.5	99	32.7
<b>GBS in influenza A(H1N1)pdm09 vaccination risk period</b>																
1 <sup>st</sup> dose <sup>1)</sup>	2	6.5	4	13.8	2	4.9	10	12.5	8	16.0	9	28.1	1	2.5	36	11.9
2 <sup>nd</sup> dose <sup>1)</sup>	0	0.0	0	0.0	2	4.9	5	6.3	0	0.0	0	0.0	0	0	7	2.3
<b>Cases during risk period following seasonal influenza vaccination</b>	0	0	1	3.4	1	2.4	4	5	4	8	0	0	5	12.5	15	0.05
<b>Cases during risk period following infections (ILI, URTI) risk period</b>																
ILI <sup>1)</sup>	5	16.1	1	3.4	2	4.9	6	7.5	13	26.0	1	3.1	1	2.5	29	9.6
URTI <sup>1)</sup>	1	3.2	5	17.2	8	19.5	10	12.5	17	34.0	7	21.9	2	5	50	16.5

<sup>1)</sup>% per number of cases included per country.Abbreviations: ILI: influenza like illness; URTI: Upper respiratory tract infection; UK: United Kingdom; NL: Netherlands; FR: France; SE: Sweden; FI: Finland; NO: Norway; DK: Denmark.  
doi:10.1371/journal.pone.0082222.t003



**Table 4.** Relative incidence estimates for the association between infections, influenza A(H1N1)pdm09 vaccination, seasonal influenza vaccination and Guillain-Barré syndrome.

	DK		FI		FR		NL		NO		SE		UK		Pooled	
	RI	95% CI	RI	95% CI	RI	95% CI	RI	95% CI	RI	95% CI	RI	95% CI	RI	95% CI	RI	95% CI
<b>Covariates</b>																
ILI	NA		NA		NA		10.5	3.0–36.3	30.6	8.6–108	NA		1.8	0.2–16.0	10.4	2.6–41.1
URTI	NA		NA		NA		13.0	4.3–39.2	17.7	6.2–34.7	NA		2.2	0.4–10.6	8.51	3.0–24.0
GI	NA		NA		NA		11.6	2.8–49.4	53.31	6.56–433	NA		2.3	0.2–22.6	11.9	2.5–55.6
Seasonal influenza vaccination	NA		NA		NA		1.2	0.4–4.0	5.5	1.6–18.9	NA		6.0	1.8–19.7	3.9	1.8–8.3
<b>Any influenza A(H1N1)pdm09 vaccination</b>																
Unadjusted	7.7	1.1–54.4	1.6	0.5–5.4	6.4	1.0–40.4	2.7	1.3–5.9	3.9	1.6–9.3	4.8	2.1–11.1	3.3	0.3–36.5	3.5	2.2–5.5
Adjusted for calendar month	3.9	0.5–32.2	1.6	0.5–5.4	2.9	0.4–19.6	1.4	0.6–3.4	1.9	0.7–5.2	2.7	1.0–7.8	2.3	0.2–27.7	2.0	1.2–3.1
<b>Adjustment effect any influenza A(H1N1)pdm09 vaccination in NL, NO, UK</b>																
Adjusted for calendar month only							1.4	0.6–3.4	1.9	0.7–5.2			2.3	0.2–27.7	1.7	0.8–3.4
Fully adjusted (month, ILI/URTI, GI)							1.2	0.5–3.3	1.5	0.5–4.6			1.5	0.1–23.1	1.4	0.7–2.8
<b>Sensitivity analysis on influenza A(H1N1)pdm09 vaccination for contra-indication</b>																
<b>Pseudolikelihood</b>																
1st dose	3.6	0.4–29.5	3.2	0.7–14.6	0.6	0.1–6.7	1.3	0.4–4.0	1.6	0.6–4.3	2.4	0.8–6.9	4.8	0.3–83.9	1.9	1.1–3.2
2nd dose	NA		NA		2.2	0.2–26.3	1.2	0.4–3.4	NA		NA		NA		1.3	0.5–3.4
<b>Vaccinated cases only</b>																
	NE		2.6	0.2–32.5	NE		1.2	0.2–8.3	1.6	0.3–7.9	2.5	0.4–16.0	NE		1.8	0.7–4.7

Abbreviations: NA: not available or not valid; NE = Not estimable due to small numbers or absence RI: relative incidence; ILI: influenza like illness; URTI: upper respiratory tract infection, GI: gastrointestinal infection, UK: United Kingdom; NL: Netherlands; FR: France; SE: Sweden; FI: Finland, NO: Norway, DK: Denmark.  
doi:10.1371/journal.pone.0082222.t004

(mainly infections) lowered the pooled calendar-month adjusted RI in NL, NO, and UK from 1.7 to 1.4. This is in line with the effect of control for infections seen by Greene et al [26]. The effect

of infections on the risk of GBS differed substantially between countries due to differences in timing and type of data collection methods. In future studies, standardization of covariate exposure

**Table 5.** Stratified analyses for association between influenza A(H1N1)pdm09 vaccination and Guillain-Barré Syndrome.

	DK		FI		FR		NL		NO		SE		UK		Pooled (random effects)	
	RI	95% CI	RI	95% CI	RI	95% CI	RI	95% CI	RI	95% CI	RI	95% CI	RI	95% CI	RI	95% CI
<b>Changing risk windows</b>																
1–28 days	4.4	(0.5 to 35.6)	1.0	(0.2–4.6)	1.3	(0.1–12.6)	2.5	(1.0–6.4)	2.2	(0.8–6.1)	2.7	(0.9–7.8)	4.2	(0.4–50.2)	2.3 <sup>4)</sup>	(1.4–3.8)
1–14 days	7.6	(0.9–61.7)	2.3	(0.5–10.6)	3.4	(0.3–33.3)	2.5	(0.7–9.3)	1.3	(0.3–5.9)	1.0	(0.2–4.7)	10.8	(0.9–133.2)	2.3 <sup>5)</sup>	(1.2–4.4)
15–28 days	NE <sup>3)</sup>		0.0		0.0		1.9	(0.7–5.5)	2.5	(0.8–7.8)	3.7	(1.2–11.1)			2.6	(1.4–4.9)
<b>42 day risk window</b>																
19–59 years old	0.0		3.3	(0.5–19.3)			1.0	(0.1–10.7)	0.6	(0.1–5.5)	1.0	(0.2–6.6)			1.3	(0.5–3.6)
older than 59 years	2.3	(0.1–38.0)	0.0		0.0		1.1	(0.3–4.9)	3.5	(1.0–12.6)	7.6	(1.6–35.8)	11.9	(0.4–365.5)	3.2	(1.5–6.9)
Co-morbidities <sup>1)</sup>	0.0		2.5	(0.2–35.5)	0.24		0.0		3.2	(0.6–17.0)	0.0				3.0	(0.7–12.3)
No co-morbidities <sup>1)</sup>	0.0		1.7	(0.4–6.7)	1.7	(0.1–19.6)	1.9	(0.6–6.6)	1.4	(0.4–5.3)	0.0				1.7	(0.8–3.4)
Seasonal influenza vaccination	0.0		3.0	(0.2–50.4)	0.2		0.5	(0.1–3.6)	2.1	(0.2–19.0)	0.0				1.2	(0.3–4.5)
No seasonal influenza vaccination	0.0		1.6	(0.4–6.4)	4.8	(0.3–83.6)	2.2	(0.4–11.2)	1.7	(0.6–5.4)	0.0				1.9	(0.9–4)
ILI, URTI infection	NE		1.1	(0.1–10.6)	2.9	(0.2–51.9)	1.1	(0.1–11.4)	1.4	(0.4–4.8)	3.2	(0.8–14.0)			1.8	(0.8–3.9)
No ILI, URTI infection	2.5	(0.2–34.4)	2.2	(0.5–10.3)	0.0		1.5	(0.4–5.8)	3.6	(0.5–24.3)	2.7	(0.6–13.2)			2.2	(1.1–4.7)

<sup>1)</sup>Malignancy, immune suppression, or autoimmune disorder NE = Not estimable due to small numbers.

Abbreviations: RI, relative incidence; ILI, influenza like illness; URTI, Upper respiratory tract infection; UK, United Kingdom; NL, Netherlands; FR, France; SE, Sweden; FI, Finland, NO, Norway, DK, Denmark.

doi:10.1371/journal.pone.0082222.t005

reporting will have to be addressed in more detail. Given the variation in the RI of GBS among other countries' A(H1N1)pdm09 vaccinees, these results, as well as the pooled estimate that was adjusted for calendar month only, are likely affected by residual confounding by infections.

## Conclusion

This large, multinational SCCS study confirms the results from the initial much smaller VAESCO case control study. In each country, the unadjusted association between influenza A(H1N1)pdm09 vaccine and GBS suggests a possible increase in risk, and adjustment for confounders consistently lowered this risk. Further adjustment for infections could only be carried out in some countries and demonstrated the effect of confounding by ILI, GI and URTI, which themselves were strong risk factors for GBS. After adjustment we did not observe an association between influenza A(H1N1)pdm09 vaccine and GBS. Based on the upper limit of the confidence interval of both the partially and fully adjusted RI estimates we can rule out with 95% certainty that adjuvanted influenza A(H1N1)pdm09 vaccines (mainly AS03 adjuvanted) would have resulted in more than 2 or 3 excess cases of GBS per 1 million vaccinated persons.

## Acknowledgments

National Institute for Health and Welfare (THL), Helsinki, FI: Jukka Ollgren; National Institute for Public Health and the Environment (RIVM), Bilthoven, NL: Hester de Melker. Haukeland University Hospital, Bergen, NO: Anette Storstein, Christian Vedeler.

All information and results concerning the UK in this study is based in part on data from the Full Feature General Practice Research Database obtained under licence from the UK Medicines and Healthcare products

## References

- Lu CY, Shao PL, Chang LY, Huang YC, Chiu CH, et al. (2010) Immunogenicity and safety of a monovalent vaccine for the 2009 pandemic influenza virus A (H1N1) in children and adolescents. *Vaccine* 28: 5864–5870.
- Lu W, Tambyah PA (2010) Safety and immunogenicity of influenza A H1N1 vaccines. *Expert Rev Vaccines* 9: 365–369.
- Wijnans L, de Bie S, Dieleman J, Bonhoeffer J, Sturkenboom M (2011) Safety of pandemic H1N1 vaccines in children and adolescents. *Vaccine* 29: 7559–7571.
- Schonberger LB, Bregman DJ, Sullivan-Bolyai JZ, Keenlyside RA, Ziegler DW, et al. (1979) Guillain-Barré syndrome following vaccination in the National Influenza Immunization Program, United States, 1976–1977. *Am J Epidemiol* 110: 105–123.
- Hurwitz ES, Schonberger LB, Nelson DB, Holman RC (1981) Guillain-Barré syndrome and the 1978–1979 influenza vaccine. *N Engl J Med* 304: 1557–1561.
- Juurink DN, Stukel TA, Kwong J, Kopp A, McGeer A, et al. (2006) Guillain-Barré syndrome after influenza vaccination in adults: a population-based study. *Arch Intern Med* 166: 2217–2221.
- Kaplan JE, Katona P, Hurwitz ES, Schonberger LB (1982) Guillain-Barré syndrome in the United States, 1979–1980 and 1980–1981. Lack of an association with influenza vaccination. *JAMA* 248: 698–700.
- Lasky T, Terracciano GJ, Magder L, Koski CL, Ballesteros M, et al. (1998) The Guillain-Barré syndrome and the 1992–1993 and 1993–1994 influenza vaccines. *N Engl J Med* 339: 1797–1802.
- Sejvar JJ, Kohl KS, Gidudu J, Amato A, Bakshi N, et al. (2011) Guillain-Barré syndrome and Fisher syndrome: case definitions and guidelines for collection, analysis, and presentation of immunization safety data. *Vaccine* 29: 599–612.
- van Doorn PA, Ruts L, Jacobs BC (2008) Clinical features, pathogenesis, and treatment of Guillain-Barré syndrome. *Lancet Neurol* 7: 939–950.
- McGrogan A, Madle GC, Seaman HE, de Vries CS (2009) The epidemiology of Guillain-Barré syndrome worldwide. A systematic literature review. *Neuroepidemiology* 32: 150–163.
- Govoni V, Granieri E (2001) Epidemiology of the Guillain-Barré syndrome. *Curr Opin Neurol* 14: 605–613.
- Dieleman J, Romio S, Johansen K, Weibel D, Bonhoeffer J, et al. (2011) Guillain-Barré syndrome and adjuvanted pandemic influenza A (H1N1) 2009 vaccine: multinational case-control study in Europe. *BMJ* 343: d3908.
- Farrington CP, Whitaker HJ, Hoxby MN (2009) Case series analysis for censored, perturbed, or curtailed post-event exposures. *Biostatistics* 10: 3–16.
- Coloma PM, Schuemie MJ, Trifiro G, Gini R, Herings R, et al. (2011) Combining electronic healthcare databases in Europe to allow for large-scale drug safety monitoring: the EU-ADR Project. *Pharmacoepidemiol Drug Saf* 20: 1–11.
- Grimaldi-Bensouda L, Alperovitch A, Besson G, Vial C, Cuisset JM, et al. (2011) Guillain-Barré syndrome, influenzalike illnesses, and influenza vaccination during seasons with and without circulating A/H1N1 viruses. *Am J Epidemiol* 174: 326–335.
- Bardage C, Persson I, Ortvist A, Bergman U, Ludvigsson JF, et al. (2011) Neurological and autoimmune disorders after vaccination against pandemic influenza A (H1N1) with a monovalent adjuvanted vaccine: population based cohort study in Stockholm, Sweden. *BMJ* 343: d5956.
- Andrews N, Stowe J, Al-Shahi Salman R, Miller E (2011) Guillain-Barré syndrome and H1N1 (2009) pandemic influenza vaccination using an AS03 adjuvanted vaccine in the United Kingdom: self-controlled case series. *Vaccine* 29: 7878–7882.
- Paul-Ehrlich-Institut (2012) Zusammenhang zwischen pandemischer Influenza A/H1N1v-Impfung und Guillain-Barré syndrome/Miller-Fisher syndrome in Deutschland.
- Wise ME, Viray M, Sejvar JJ, Lewis P, Baughman AL, et al. (2012) Guillain-Barré Syndrome During the 2009–2010 H1N1 Influenza Vaccination Campaign: Population-based Surveillance Among 45 Million Americans. *Am J Epidemiol* 175: 1110–1119.
- Tokars JJ, Lewis P, DeStefano F, Wise M, Viray M, et al. (2012) The risk of Guillain-Barré syndrome associated with influenza A (H1N1) 2009 monovalent vaccine and 2009–2010 seasonal influenza vaccines: results from self-controlled analyses. *Pharmacoepidemiol Drug Saf* 21: 546–552.
- Yih WK, Lee GM, Lieu TA, Ball R, Kulldorff M, et al. (2012) Surveillance for Adverse Events Following Receipt of Pandemic 2009 H1N1 Vaccine in the Post-Licensure Rapid Immunization Safety Monitoring (PRISM) System, 2009–2010. *Am J Epidemiol* 175: 1120–1128.
- Greene SK, Rett M, Weintraub ES, Li L, Yin R, et al. (2012) Risk of Confirmed Guillain-Barré syndrome Following Receipt of Monovalent Inactivated Influenza A (H1N1) and Seasonal Influenza Vaccines in the Vaccine Safety Datalink Project, 2009–2010. *Am J Epidemiol* 175: 1100–1109.
- Salmon DA, Proschan M, Forshee R, Gargiullo P, Bleser W, et al. (2013) Association between Guillain-Barre syndrome and influenza A (H1N1) 2009 monovalent inactivated vaccines in the USA: a meta-analysis. *Lancet*.

Regulatory Agency and covers the data collection time period up to February 2011. However, the interpretation and conclusions contained in this report are those of the authors alone.

### Ethical approval

In **France**, the case-control study was approved by the ethics committee (i.e., Comité de Protection des Personnes Sud Ouest et Outre Mer I et II) and participants gave informed consent. In the **Netherlands**, the study was provided with a declaration of no objection from the medical ethics committee of the Erasmus University Medical Center in Rotterdam (MEC-2009-404). Subsequent amendments to collect data entirely anonymously through the GP allowed the inclusion of study subjects without requiring informed consent. In **Denmark** ethics approval was not required, though the National Board of Health approved chart review. In **Sweden** the study was approved by the regional ethics committee, Karolinska Institute, Stockholm, and participants gave written informed consent. In the **UK**, the GPRD Group has obtained ethical approval from a multicenter research ethics committee (MREC) for all purely observational research using GPRD data. Individual studies must be granted approval by an Independent Scientific Advisory Committee (ISAC). This study received ISAC approval (protocol No 10\_058). In **Norway**, the Regional Committee for Medical and Health Research Ethics in Western Norway approved national study participation without need for patient consent.

## Author Contributions

Wrote the paper: SR DW JPD HKO MCJMS. Study design, data management, data analysis, interpretation of data and results, and reviewing of manuscript: SR DW JPD HKO CSDV CS NA HS DMN AH MLM AS CS AC HH LAD PS MM MS NVDM BCJ TL TK JS KJ PK JB MCJMS. Main pooled statistical data analysis: SR. Main pooled statistical data analysis, advisory role: NA. Scientific coordination of VAESCO consortium: DW JB MCJMS. Development of software for data collection, harmonization, and aggregation across databases: MS MM. European Centre for Disease Prevention and Control, project leader for the GBS and pandemic vaccines: KJ PK.

25. De Wals P, Decuninck G, Toth E, Boulianne N, Brunet D, et al. (2012) Risk of Guillain-Barré syndrome following H1N1 influenza vaccination in Quebec. *JAMA* 308: 175–181.
26. Greene SK, Rett MD, Vellozzi C, Li L, Kulldorff M, et al. (2013) Guillain-Barre Syndrome, Influenza Vaccination, and Antecedent Respiratory and Gastrointestinal Infections: A Case-Centered Analysis in the Vaccine Safety Datalink, 2009–2011. *PLoS One* 8: e67185.
27. Poland GA, Poland CM, Howe CL (2013) Influenza vaccine and Guillain-Barre syndrome: making informed decisions. *Lancet* 381: 1437–1439.
28. Stowe J, Andrews N, Wise L, Miller E (2009) Investigation of the temporal association of Guillain-Barre syndrome with influenza vaccine and influenzalike illness using the United Kingdom General Practice Research Database. *Am J Epidemiol* 169: 382–388.