

## Title

Guiding synovial inflammation by macrophage phenotype modulation: an *in vitro* study towards a therapy for osteoarthritis

## Authors and affiliations

Lizette Utomo (l.utomo@erasmusmc.nl)<sup>1</sup>, Gerjo J.V.M. van Osch (g.vanosch@erasmusmc.nl)<sup>1,2</sup>, Yves Bayon (yves.bayon@medtronic.com)<sup>3</sup>, Jan A.N. Verhaar (j.verhaar@erasmusmc.nl)<sup>1</sup>, Yvonne M. Bastiaansen-Jenniskens (y.bastiaansen@erasmusmc.nl)<sup>1,\*</sup>

<sup>1</sup>Department of Orthopaedics, <sup>2</sup>Department of Otorhinolaryngology, Erasmus MC, University Medical Center Rotterdam, Rotterdam the Netherlands

<sup>3</sup>Medtronic-Sofradim Production, Trévoux, France

## \*Corresponding author

Name: Yvonne M. Bastiaansen-Jenniskens, PhD

Address: Erasmus MC, University Medical Center Rotterdam  
Wytemaweg 80, room Ee 16.51b  
3015 CN Rotterdam  
the Netherlands

Telephone: +31 107044626

E-mail: y.bastiaansen@erasmusmc.nl

Running title: Synovial macrophage modulation

## Abstract

*Objective:* The aims of this study were to modulate inflammation in synovial explants with the compounds: dexamethasone, rapamycin, bone morphogenetic protein 7 (BMP-7) and pravastatin, and to investigate the modulatory capacity of the compounds on specific macrophage phenotypes.

*Design:* Synovial explants from osteoarthritis (OA) patients were treated with  $10^{-6}$ M dexamethasone, 100 ng/mL rapamycin, 500 ng/mL BMP-7 or 50  $\mu$ M pravastatin. Half of the explants were pre-stimulated with IFN $\gamma$ +TNF $\alpha$  to simulate acute inflammation. Inflammatory state of the synovium was assessed with gene expression analysis. Primary human monocytes were isolated and stimulated towards macrophage phenotypes M(IFN $\gamma$ +TNF $\alpha$ ), M(IL-4) and M(IL-10) with the respective cytokines, followed by treatment with the compounds.

*Results:* Dexamethasone had an anti-inflammatory effect on IFN $\gamma$ +TNF $\alpha$  stimulated and osteoarthritic synovium, likely due to suppression of pro-inflammatory M(IFN $\gamma$ +TNF $\alpha$ ) macrophages while enhancing anti-inflammatory M(IL4) and M(IL10) macrophages. Rapamycin and BMP-7 further enhanced inflammation in stimulated synovium, but rapamycin did not have a clear effect on non-stimulated synovium. Rapamycin suppressed M(IL-4) and M(IL-10) macrophages without affecting M(IFN $\gamma$ +TNF $\alpha$ ). BMP-7 suppressed M(IFN $\gamma$ +TNF $\alpha$ ) and enhanced M(IL-10) in the macrophage cultures. Pravastatin did not affect synovium, but enhanced M(IL-10).

*Conclusions:* These data indicate that macrophage phenotype modulation can be used to guide joint inflammation and thereby contribute to the development of new therapies to delay the progression of OA. The varying effects of the compounds on synovium of different degrees of inflammation, indicate that the modulatory capacity of the compounds depends on OA stage and underlines the importance of identifying this stadium for adequate treatment.

Keywords: synovium, inflammation, osteoarthritis, macrophage phenotypes, modulation

## Introduction

Osteoarthritis (OA), a chronic degenerative disease affecting the whole joint, is characterized by erosion of articular cartilage and synovial inflammation. Macrophages, along with fibroblasts, reside in the synovial lining of joints. These cells can become activated by stimuli from their microenvironment, a process that results in a spectrum of phenotypes [1]. This spectrum ranges from pro-inflammatory (M1) macrophages, induced by stimuli such as interferon (IFN), tumor necrosis factors (TNFs) [2] or lipopolysaccharides (LPS) [3], to anti-inflammatory macrophages (M2), which can be further divided into specific subtypes depending on their stimuli such as IL-4 or IL-10 [3]. Because of their importance in diseases and homeostasis, macrophages are frequently the focus of development of new interventions and treatment strategies [4]. Early *in vivo* studies have shown that depletion of phagocytic synovial lining cells, resulted in less influx of polymorphonuclear neutrophils, reduced proteoglycan degradation and reduced chondrocyte death of cartilage after induction of collagen induced arthritis (CIA) [5]. It was suggested that monocytes/macrophages are mainly responsible for this cartilage damage [6]. Moreover, osteophyte formation was also reduced after depletion of synovial macrophages in an OA mouse model [7]. The fact that high levels of pro-inflammatory cytokines were found in synovial fluid of OA joints and after trauma [8, 9], together with the role macrophages seem to play in the development of OA and other inflammatory joint diseases, led to the hypothesis that a potentially specific strategy to control inflammation would be modulating the polarization state of macrophages. Therefore, the aim of this study was to guide synovial inflammation with four compounds representing commonly used classes of drugs in the clinics, that may have the potential to modulate macrophage phenotype: dexamethasone, rapamycin, bone morphogenetic protein 7 (BMP-7) and pravastatin. Dexamethasone is a corticosteroid with well-known anti-inflammatory effects [10] and is used to treat numerous inflammatory-based diseases. Rapamycin, known by its non-proprietary name sirolimus, is an immunosuppressive commonly used in transplantation medicine [11]. It inhibits T-cell activation by inhibiting signal transduction via mammalian target of rapamycin (mTor) [12]. BMP-7 is a member of the transforming growth factor beta (TGF- $\beta$ ) superfamily and is clinically used in orthopedic studies and applications, such as long bone non-union fractures [13, 14]. Additionally, it has been shown that BMP-7 directed the polarization of THP-1 monocytes into an anti-inflammatory state [15]. Pravastatin is a member of the statins, a group of cholesterol synthesis inhibitors with a broad spectrum of effects, including anti-inflammatory effects [16]. Since all compounds are known to affect inflammatory processes, we investigated their effects on synovium of different degrees of inflammation. For this reason, OA synovium was

additionally stimulated with IFN $\gamma$ +TNF $\alpha$  to simulate acute inflammation. Furthermore, to examine the modulatory effects of the compounds on macrophage phenotypes, human monocyte-derived macrophages were polarized to specific phenotypes and treated with the same compounds.

## **Method**

### *Modulation of synovium*

To examine the effect of stimulation with IFN $\gamma$ +TNF $\alpha$ , synovial tissue was obtained from OA patients ( $n=4$ ,  $60\pm 13$ Y) undergoing total knee replacement surgery. The synovium was washed twice with 0.9% NaCl (Sigma-Aldrich, St. Louis, USA) and cut into pieces of 30-35 mg wet weight. To simulate acute inflammation [8, 9], the explants were cultured for 24h with 10 ng/mL Interferon- $\gamma$  (IFN $\gamma$ ; PeproTech, Rocky Hill, NJ, USA) and 10 ng/mL Tumor Necrosis Factor- $\alpha$  (TNF $\alpha$ , PeproTech) in medium (Dulbecco's Modified Eagle Medium, low glucose (DMEM; Gibco, Carlsbad, USA) supplemented with 1% Insulin-Transferrin-Selenium (ITS+ Premix, Corning, New York, USA), 50  $\mu$ g/mL gentamicin (Gibco), 1.5  $\mu$ g/mL amphotericin B (Fungizone; Gibco) and 25  $\mu$ g/mL L-ascorbic acid 2-phosphate (Sigma-Aldrich)). After 24h of stimulation, the synovial explants were harvested and stored at  $-80^{\circ}\text{C}$  until assessment for their inflammatory state after stimulation using gene expression analysis. To examine the effect of the compounds, synovial explants of other patients ( $n=4$ ;  $63\pm 3$ Y) were pre-cultured for 24h with or without IFN $\gamma$ +TNF $\alpha$  as described above to obtain acute inflamed synovium. Then,  $10^{-6}$  M dexamethasone (Sigma-Aldrich), 100 ng/mL rapamycin (R&D Systems, Minneapolis, USA), 500 ng/mL BMP-7 (PeproTech), or 50  $\mu$ M pravastatin (Sigma-Aldrich) were added to the medium and the explants were cultured for an additional 3 days. Doses were chosen based on literature [15, 17-22]. Dimethyl sulfoxide (DMSO; Sigma-Aldrich) was used as vehicle for dexamethasone and rapamycin and the final DMSO concentration in the cultures was 0.01%. Donor demographics and culture conditions are shown in Table 1. The medium including compounds and stimuli was refreshed 24h prior to harvest and the explants were stored at  $-80^{\circ}\text{C}$  until further processing for gene expression analysis.

### *Monocyte isolation, stimulation and macrophage modulation*

Primary human monocytes were polarized to specific macrophage phenotypes and cultured with the compounds. Monocytes were isolated by Ficoll density gradient separation and CD14<sup>+</sup> selection as described previously [23] from human buffy coats of male donors, 52±14Y (Sanquin Blood bank, Amsterdam, the Netherlands). For every experiment, monocytes were pooled from at least two donors and plated in monolayers in 48-well plates (Corning Costar, NY, USA) at a density of 500,000 monocytes/cm<sup>2</sup> in X-VIVO™ 15 medium (Lonza, Verviers, Belgium) supplemented with 20% heat-inactivated fetal calf serum (FCS; Lonza), 50 µg/mL gentamicin and 1.5 µg/mL amphotericin B. To model a range of phenotypes of activated macrophages, the monocytes were stimulated 1 h after plating with 10 ng/mL IFN $\gamma$  and 10 ng/mL TNF $\alpha$  (from now on referred to as M(IFN $\gamma$ +TNF $\alpha$ )), 10 ng/mL Interleukin-4 (IL-4; PeproTech), to obtain M(IL-4) or 10 ng/mL IL-10 (PeproTech) to obtain M(IL-10) [2, 3]. The macrophages were cultured and stimulated for 3 days at 37°C and 5% CO<sub>2</sub>. After the stimulation period, the polarized macrophages were treated either with 10<sup>-6</sup> M dexamethasone, 100 ng/mL rapamycin, 500 ng/mL BMP-7 or 50 µM pravastatin and cultured for an additional 3 days. The final DMSO concentration used in the cultures was 0.01%. The medium including compounds and stimuli was refreshed 24h prior to harvest. After culture, the medium was collected, centrifuged at 200 x g to remove detached cells, and the supernatants were stored at -80°C until cytokine measurements. The cells were harvested in PBS/0.1% Triton X-100 (Sigma-Aldrich) for DNA quantification or in RLT lysis buffer/1%  $\beta$ -mercaptoethanol (Qiagen, Hilden, Germany/Sigma-Aldrich) for mRNA isolation.

#### *Gene expression analysis*

mRNA isolation, cDNA synthesis and PCR analysis were performed as described previously. [24] Gene expression of Interleukin-6 (*IL6*), Interleukin-1 $\beta$  (*IL1B*), Tumor Necrosis Factor- $\alpha$  (*TNFA*), chemokine (C-C motif) ligand 18 (*CCL18*), Interleukin-1 Receptor Antagonist (*IL1RA*), Mannose receptor, C type 1/CD206 (*MRC1/CD206*), cluster of differentiation 163 (*CD163*), Toll-Like-Receptor 4 (*TLR4*), and Transforming Growth Factor  $\beta$ 1 (*TGFB1*) was evaluated. Glyceraldehyde-3-phosphate dehydrogenase (*GAPDH*), hypoxanthine phosphoribosyltransferase 1 (*HPRT1*) and ubiquitin C (*UBC*) were all tested as housekeepers, where *GAPDH* was found the most stable (data not shown) and was therefore further used as normalization for the genes of interest. The amplification efficiency of all primers (Table S1) was between 0.90 and 1.05 and the relative expression was determined by the 2<sup>- $\Delta$ CT</sup> formula.

### *Quantification of cytokine production*

Production of IL-6, CCL18 and soluble CD163 (sCD163) by the macrophages was quantified by enzyme-linked immunosorbent assays (ELISAs) according to the manufacturer's instructions of the human IL-6 ELISA Development Kit (PeproTech), human CCL18 DuoSet Development Kit (R&D Systems) and human soluble CD163 DuoSet Development Kit (R&D Systems). As an indication of the number of cells in the monolayers and to normalize cytokine production, the DNA content of the monolayers was measured using a modified CyQUANT assay (Invitrogen, Carlsbad, USA) as described previously [24].

### *M1/M2-index calculation*

An M1/M2-index was calculated based on the expression of the pro-inflammatory (M1) genes (*IL6*, *IL1B* and *TNFA*) and the anti-inflammatory (M2) genes (*CCL18*, *IL1RA*, *CD206* and *CD163*). The mean of the relative expression of M1 genes per sample to the overall M1 gene expression of all samples, was divided over the mean of the relative expression of M2 genes per sample to the overall M2 gene expression of all samples [23].

### *Statistics*

MS Excel 2010 and PASW Statistics 21.0 (SPSS Inc., Chicago, USA) were used for statistical evaluation. To take into account the variability between donors, a mixed linear model after log transformation was used, followed by a Bonferroni's *post hoc* comparisons test. For the synovial explant experiments, inflammation state of the synovium (i.e., stimulation with IFN $\gamma$ +TNF $\alpha$ ) and donor were defined as random factors, while compound treatment was defined as a fixed factor. For the macrophages cultured in monolayers, polarization together with compound treatment were defined as fixed factors, while individual experiments were considered as random factors in the model. Differences were considered to be statistically significant for  $P < 0.05$ .

## **Results**

### *Stimulation of osteoarthritic synovium leads to a difference in gene expression profiles*

To investigate the modulatory capacity of the compounds on synovium of different degrees of inflammation, a culture model with or without IFN $\gamma$ +TNF $\alpha$  stimulation was set up. Gene expression of *IL6*, *IL1B*, *TNFA*, *CCL18* and *IL1RA* was higher in OA synovium stimulated with IFN $\gamma$ +TNF $\alpha$  than of synovium without stimulation when cultured in DMEM only [Fig. 1(A)]. The expression profile of IFN $\gamma$ +TNF $\alpha$  stimulated synovium resulted in a higher M1/M2-index [Fig. 1(B)], indicating that acute inflammation was indeed induced in the culture model.

#### *Dexamethasone, rapamycin, BMP-7 and pravastatin can modulate the gene expression profile of synovium*

In IFN $\gamma$ +TNF $\alpha$  stimulated OA synovium treated with dexamethasone, *IL1B*, *IL6*, *TNFA*, *IL1RA* and *CCL18* were lower than in the stimulated synovium cultured in DMSO control medium, while *CD206* and *CD163* were higher. This expression profile resulted in a lower M1/M2-index than in the controls. Treatment with rapamycin increased *IL6*, while lowering *CCL18* in stimulated synovium. This resulted in a higher M1/M2-index than in the DMSO controls. BMP-7 lowered expression of *IL1RA* compared to the DMEM controls, while pravastatin did not affect any of the genes of interest. For both BMP-7 and pravastatin, this expression profile resulted in a higher M1/M2-index than in the controls [Fig. 2(A)].

In non-stimulated OA synovium treated with dexamethasone, *IL1B* was lower than in non-stimulated synovium cultured in DMSO control medium, while *CD163* was higher, eventually resulting in a lower M1/M2-index. Rapamycin did not affect the genes of interest, but the overall inflammation was reduced as indicated by a lower M1/M2-index. BMP-7 lowered *IL1B* and increased *CCL18*, resulting in a lower M1/M2-index than in the DMEM controls. Pravastatin did not affect the genes of interest and did not alter the overall inflammatory state of non-stimulated synovium [Fig. 2(B)].

#### *Characterization and modulation of primary polarized macrophages*

To assess the modulatory capacity of the compounds on specific macrophage phenotypes, primary monocyte-derived macrophages were polarized towards specific macrophage phenotypes and treated with the compounds. Without treatment, M(IFN $\gamma$ +TNF $\alpha$ ) had high expression levels of *IL6*, *IL1B*, *TNFA*, *IL10* and

*TFGB1* and high protein production of IL-6. M(IL-4) expressed high levels of *CC18*, *IL1RA* and *CD206* and had high CCL18 protein production. M(IL-10) had higher expression of *CD206* than M(IFN $\gamma$ +TNF $\alpha$ ), but was lower than in M(IL-4). *CD163* was mainly expressed in anti-inflammatory macrophages and expression levels were higher in M(IL-10) than in M(IL-4). sCD163 protein production was high in M(IL-10) [Fig. 3(A) and (B)]. This confirmed that stimulation with IFN $\gamma$ +TNF $\alpha$  resulted in pro-inflammatory macrophages and IL-4 or IL-10 stimulation resulted in the development of anti-inflammatory macrophages of distinct phenotypes. As the IL-6, CCL18 and sCD163 protein production patterns were also specific for our generated phenotypes, we decided to use these three proteins as read-out parameters for detailed examination of the effects of the compounds on macrophages in monolayer.

Interestingly, after treatment with dexamethasone, IL-6 production by the M(IFN $\gamma$ +TNF $\alpha$ ) was decreased, while production levels were maintained in M(IL-4) and M(IL-10). Production of CCL18 by M(IFN $\gamma$ +TNF $\alpha$ ) was maintained as well, whereas CCL18 production was increased in M(IL-4) and M(IL-10). sCD163 production was highly increased in all three macrophage phenotypes after treatment with dexamethasone [Fig. 4(A)]. Treatment with rapamycin lowered the IL-6 and CCL18 production of M(IL-4), CCL18 production of M(IL-10) and sCD163 production of M(IL-4) [Fig. 4(B)]. BMP-7 treatment did not have an effect on the IL-6, CCL18 and sCD163 protein production of M(IFN $\gamma$ +TNF $\alpha$ ) and M(IL-4), while CCL18 and sCD163 production was increased in M(IL-10) [Fig. 4(C)]. Pravastatin did not have an effect on the IL-6 and CCL18 production of either macrophage phenotypes, yet sCD163 was higher in M(IL-10) than in the untreated controls [Fig. 4(D)].

## **Discussion**

In this study, we have shown that the overall inflammatory state of synovial explants can be modulated by dexamethasone, rapamycin, and BMP-7. Because macrophages play a crucial role in inflammatory processes that contribute to inflammation, phenotype modulation can be used to control synovial inflammation, and subsequently may also inhibit the progression of post-traumatic OA.

Nowadays, many compounds are being tested for their anti-inflammatory effects *in vitro* as potential new interventions for OA. However, completely suppressing inflammation, either via macrophages or by directly inhibiting cytokines, may not be the most successful method to treat OA, as inflammation to a certain extend is required for proper wound healing [25]. Since the composition of synovial tissue varies at different



stages of OA [26] and even between patients, we can assume that the composition of macrophage phenotypes in the synovium also differs, which was indicated in our previous work [27]. To our knowledge, we are the first to show modulation of synovium of different degrees of inflammation using compounds that are already clinically applied. In addition, we showed that these compounds target specific macrophage phenotypes by either enhancing or suppressing their function. Our culture model using human synovium allowed us to study *in vitro* the effects of the compounds on synovial macrophages in their natural microenvironment. We are aware that we cannot rule out the possibility that by treating the explants with the compounds, fibroblasts or other cells present in the synovium were also affected by the compounds. Therefore, we compared the modulatory effects of the compounds on the entire synovial explants in terms of change in overall synovial inflammation, with their effect on different phenotypes of macrophages cultured in monolayers. Macrophages, once migrated into tissue, are in an activated state. With the polarization of M(IFN $\gamma$ +TNF $\alpha$ ), M(IL-4) and M(IL-10), we aim to model this range of phenotypes. Our aim was to modulate already polarized macrophages in synovial tissue and not to intervene with the process of polarization from monocyte to macrophage. Therefore, assessment of the effects of the compounds on unstimulated macrophages is beyond the scope of this study. Additionally, we have shown that compounds that were initially known for their anti-inflammatory effect, can either behave in a pro-inflammatory or anti-inflammatory manner and that the behavior of the compounds differs and depends on the macrophage phenotypes present in the tissue that is treated. The specific modulatory effects of the compounds on synovium and macrophages could provide valuable insights for the development of specialized therapies aiming at modulation of specific macrophage phenotypes.

In our study, dexamethasone had an anti-inflammatory effect on synovium and suppressed the pro-inflammatory macrophage phenotype, while enhancing the anti-inflammatory macrophages. Since *CD163* expression can be assumed as a marker for M(IL-10), these data combined suggests that dexamethasone has the capacity to modulate synovial tissue and to specifically enhance the macrophage phenotypes resembling M(IL-10). The effect of dexamethasone on macrophages was shown earlier to vary depending on macrophage origin, as phagocytic activity in alveolar macrophages was induced more abundantly than in peritoneal macrophages [28]. The difference in anti-inflammatory capacity per macrophage source suggests that treatment with dexamethasone may result in responses that differ according to macrophage type or origin. Intra-articular delivery of dexamethasone was shown to have chondroprotective effects and reduced inflammation in a post-traumatic OA animal model [29]. This could be explained by the fact that dexamethasone enhanced M(IL-10), which was seen in our present study, as well as of others [30], on gene expression level and by

increased levels of sCD163 in the macrophage cultures. Although this subtype expressing CD163 has been described as the tissue repair phenotype [31], it has also been associated with (chronic) inflammatory diseases [32, 33]. Additionally, CD163 expressing macrophages induced by intra-articular injection of triamcinolone acetonide, also a corticosteroid, have been linked as well with the prevention of osteophyte formation in an OA rat model [34].

Rapamycin exerted an overall pro-inflammatory effect on IFN $\gamma$ +TNF $\alpha$  stimulated synovium in this study, while the effect on non-stimulated OA synovium was less clear as no differences in expression levels were seen of any of the genes of interest. Our data indicated that the increase of inflammation of stimulated synovium in response to rapamycin was likely due to suppression of anti-inflammatory macrophages, without affecting the pro-inflammatory macrophages, as indicated by reduced IL-6, CCL18 and sCD163 protein production in the anti-inflammatory macrophages, while these levels were maintained in M(IFN $\gamma$ +TNF $\alpha$ ) after treatment with rapamycin. This was in line with an earlier study, where inhibition of the mTOR signaling pathway was shown to regulate macrophage polarization. Murine macrophages in which *Tsc1* was specifically deleted, and therefore mTOR complex 1 constitutively activated, were unable to polarize towards M2, while the pro-inflammatory response to LPS was enhanced. Treatment of these macrophages with rapamycin rescued this M2 polarization deficiency [17]. Furthermore, rapamycin seemed to shift the polarization of human macrophages towards a pro-inflammatory phenotype *in vitro* [35]. BMP-7 had an overall pro-inflammatory effect on IFN $\gamma$ +TNF $\alpha$  stimulated synovium, while an anti-inflammatory effect on non-stimulated OA synovium. The effect on non-stimulated OA synovium can partially be explained by the effects seen on polarized macrophages, where BMP-7 increased the CCL18 and sCD163 production by M(IL-10), suggesting an anti-inflammatory effect. These data were in line with another study [15], where it was shown that BMP-7 directed the polarization of THP-1 monocytes into an anti-inflammatory state. The anti-inflammatory effect of BMP-7 indicates that the modulatory capacity of BMP-7 may depend on the phenotype of the cells that are present in the tissue, assuming that the majority of the macrophages in OA synovium have a similar phenotype to IL-10 stimulated primary macrophages. On the other hand, the reason for the overall pro-inflammatory effect of BMP-7 seen in IFN $\gamma$ +TNF $\alpha$  stimulated synovial tissue remains unclear, although it is likely due to a decrease of *IL1RA* expression which led to a reduced M1/M2-index. However, no effect was seen on pro-inflammatory macrophages in the monolayer cultures. These data combined suggest that BMP-7 may also have an effect on other cells than macrophages that are present in synovium that are involved in inflammatory processes as well. Other studies have shown that intra-articular administration of BMP-7 [18, 36-38] and rapamycin [39-41] protected OA

progression after induction of experimental OA. These studies focused mainly on articular cartilage quality in end-stage OA and have not taken into account the direct effect of BMP-7 and rapamycin on the synovial membrane. The varying effects of these two compounds on IFN $\gamma$ +TNF $\alpha$  stimulated and non-stimulated synovium, underlines the importance of identifying OA stadium in order to clinically use BMP-7 or rapamycin as a therapy for OA.

Although statins are known for their pleiotropic effects, pravastatin did not have a clear modulatory effect on OA synovial tissue in our study. On polarized macrophages however, pravastatin increased sCD163 production in both anti-inflammatory phenotypes, which is in line with other studies that have shown anti-inflammatory effects of statins on macrophage polarization [42, 43], chondrocytes [44, 45], and progression of OA and arthritis *in vivo* [46, 47]. Additionally, systemic statin use has also been shown to be associated with reduced progression of knee OA [48] and seemed to reduce the activity of rheumatoid arthritis in humans [49]. We chose to assess the effects of pravastatin over other statins such as simvastatin, since the latter needs to be bio-activated by the liver, or manually if it is not administered orally [50]. The fact that pravastatin was unable to modulate synovial tissue, suggests that it may not be suitable to be used intra-articular for guiding synovial inflammation by macrophage phenotype modulation, unless specifically targeting M(IL-10) like macrophages.

To conclude, dexamethasone, rapamycin and BMP-7 can modulate the overall inflammatory state of OA synovium by altering their expression profile. Future research could include large population patient studies to specifically correlate initial synovitis to the response after treatment. Directing synovial inflammation to delay the progression of OA may therefore be a suitable personalized medicine approach for which the optimal compound can be selected when the inflammatory state of the synovium has been determined.

### **Author contributions**

LU designed the study, performed the experiments, analyzed and interpreted the data, and wrote the manuscript. GJVMvO contributed to data interpretation and edited the manuscript. YB and JANV edited the manuscript. YMBJ designed the study, interpreted the data and edited the manuscript. All authors approved the final version of the manuscript.

### **Funding**

This study was financially supported by the Dutch Arthritis Foundation (#13-3-302). YMBJ is also supported by a Veni grant from Technology Foundation STW and by The Dutch Arthritis Foundation (#LLP11).

### **Conflict of interests**

YB is an employee of Medtronic-Sofradim Production. None of the financial supporters were involved in the design, conduct or analysis of this study.

### **Ethics approval**

The experiments involving human synovium have been conducted with the approval of the medical ethical committee of the Erasmus Medical Center (#MEC2004-322). The synovium was obtained with implicit consent as waste material from patients undergoing knee replacement surgery. The patients had the right to refuse as stated by the guidelines of the Dutch Federation of Biomedical Scientific Societies ([www.federa.org](http://www.federa.org)).

### **References**

1. Murray PJ, Allen JE, Biswas SK, Fisher EA, Gilroy DW, Goerdts S, et al. Macrophage activation and polarization: nomenclature and experimental guidelines. *Immunity* 2014; 41: 14-20.
2. Cassol E, Cassetta L, Rizzi C, Alfano M, Poli G. M1 and M2a polarization of human monocyte-derived macrophages inhibits HIV-1 replication by distinct mechanisms. *J Immunol* 2009; 182: 6237-6246.
3. Spiller KL, Anfang RR, Spiller KJ, Ng J, Nakazawa KR, Daulton JW, et al. The role of macrophage phenotype in vascularization of tissue engineering scaffolds. *Biomaterials* 2014; 35: 4477-4488.
4. Dey A, Allen J, Hankey-Giblin PA. Ontogeny and polarization of macrophages in inflammation: blood monocytes versus tissue macrophages. *Front Immunol* 2014; 5: 683.
5. Van Lent PL, Holthuysen AE, Van Rooijen N, Van De Putte LB, Van Den Berg WB. Local removal of phagocytic synovial lining cells by clodronate-liposomes decreases cartilage destruction during collagen type II arthritis. *Ann Rheum Dis* 1998; 57: 408-413.

6. Van Lent PL, Blom A, Holthuysen AE, Jacobs CW, Van De Putte LB, Van Den Berg WB. Monocytes/macrophages rather than PMN are involved in early cartilage degradation in cationic immune complex arthritis in mice. *J Leukoc Biol* 1997; 61: 267-278.
7. Blom AB, van Lent PL, Holthuysen AE, van der Kraan PM, Roth J, van Rooijen N, et al. Synovial lining macrophages mediate osteophyte formation during experimental osteoarthritis. *Osteoarthritis Cartilage* 2004; 12: 627-635.
8. Bigoni M, Sacerdote P, Turati M, Franchi S, Gandolla M, Gaddi D, et al. Acute and late changes in intraarticular cytokine levels following anterior cruciate ligament injury. *J Orthop Res* 2013; 31: 315-321.
9. Catterall JB, Stabler TV, Flannery CR, Kraus VB. Changes in serum and synovial fluid biomarkers after acute injury (NCT00332254). *Arthritis Res Ther* 2010; 12: R229.
10. Barnes PJ. How corticosteroids control inflammation: Quintiles Prize Lecture 2005. *Br J Pharmacol* 2006; 148: 245-254.
11. Halloran PF. Immunosuppressive drugs for kidney transplantation. *N Engl J Med* 2004; 351: 2715-2729.
12. Ballou LM, Lin RZ. Rapamycin and mTOR kinase inhibitors. *J Chem Biol* 2008; 1: 27-36.
13. Pecina M, Giltaij LR, Vukicevic S. Orthopaedic applications of osteogenic protein-1 (BMP-7). *Int Orthop* 2001; 25: 203-208.
14. Bishop GB, Einhorn TA. Current and future clinical applications of bone morphogenetic proteins in orthopaedic trauma surgery. *Int Orthop* 2007; 31: 721-727.
15. Rocher C, Singla DK. SMAD-PI3K-Akt-mTOR pathway mediates BMP-7 polarization of monocytes into M2 macrophages. *PLoS One* 2013; 8: e84009.
16. Mihos CG, Pineda AM, Santana O. Cardiovascular effects of statins, beyond lipid-lowering properties. *Pharmacol Res* 2014; 88: 12-19.
17. Byles V, Covarrubias AJ, Ben-Sahra I, Lamming DW, Sabatini DM, Manning BD, et al. The TSC-mTOR pathway regulates macrophage polarization. *Nat Commun* 2013; 4: 2834.
18. Hayashi M, Muneta T, Ju YJ, Mochizuki T, Sekiya I. Weekly intra-articular injections of bone morphogenetic protein-7 inhibits osteoarthritis progression. *Arthritis Res Ther* 2008; 10: R118.

19. Lee DK, Park EJ, Kim EK, Jin J, Kim JS, Shin JJ, et al. Atorvastatin and simvastatin, but not pravastatin, up-regulate LPS-induced MMP-9 expression in macrophages by regulating phosphorylation of ERK and CREB. *Cell Physiol Biochem* 2012; 30: 499-511.
20. Mlambo G, Sigola LB. Rifampicin and dexamethasone have similar effects on macrophage phagocytosis of zymosan, but differ in their effects on nitrite and TNF-alpha production. *Int Immunopharmacol* 2003; 3: 513-522.
21. Shappley RK, Spentzas T. Differential Role of Rapamycin and Torin/KU63794 in Inflammatory Response of 264.7 RAW Macrophages Stimulated by CA-MRSA. *Int J Inflam* 2014; 2014: 560790.
22. Takahashi T, Muneta T, Tsuji K, Sekiya I. BMP-7 inhibits cartilage degeneration through suppression of inflammation in rat zymosan-induced arthritis. *Cell Tissue Res* 2011; 344: 321-332.
23. Grotenhuis N, Bayon Y, Lange JF, Van Osch GJ, Bastiaansen-Jenniskens YM. A culture model to analyze the acute biomaterial-dependent reaction of human primary macrophages. *Biochem Biophys Res Commun* 2013; 433: 115-120.
24. Utomo L, Pleumeekers MM, Nimeskern L, Nurnberger S, Stok KS, Hildner F, et al. Preparation and characterization of a decellularized cartilage scaffold for ear cartilage reconstruction. *Biomed Mater* 2015; 10: 015010.
25. Koh TJ, DiPietro LA. Inflammation and wound healing: the role of the macrophage. *Expert Rev Mol Med* 2011; 13: e23.
26. Benito MJ, Veale DJ, FitzGerald O, van den Berg WB, Bresnihan B. Synovial tissue inflammation in early and late osteoarthritis. *Ann Rheum Dis* 2005; 64: 1263-1267.
27. Fahy N, de Vries-van Melle ML, Lehmann J, Wei W, Grotenhuis N, Farrell E, et al. Human osteoarthritic synovium impacts chondrogenic differentiation of mesenchymal stem cells via macrophage polarisation state. *Osteoarthritis Cartilage* 2014; 22: 1167-1175.
28. Nakamura Y, Murai T, Ogawa Y. Effect of in vitro and in vivo administration of dexamethasone on rat macrophage functions: comparison between alveolar and peritoneal macrophages. *Eur Respir J* 1996; 9: 301-306.
29. Huebner KD, Shrive NG, Frank CB. Dexamethasone inhibits inflammation and cartilage damage in a new model of post-traumatic osteoarthritis. *J Orthop Res* 2014; 32: 566-572.


30. Buechler C, Ritter M, Orso E, Langmann T, Klucken J, Schmitz G. Regulation of scavenger receptor CD163 expression in human monocytes and macrophages by pro- and antiinflammatory stimuli. *J Leukoc Biol* 2000; 67: 97-103.
31. Gordon S, Martinez FO. Alternative activation of macrophages: mechanism and functions. *Immunity* 2010; 32: 593-604.
32. Fonseca JE, Edwards JC, Blades S, Goulding NJ. Macrophage subpopulations in rheumatoid synovium: reduced CD163 expression in CD4+ T lymphocyte-rich microenvironments. *Arthritis Rheum* 2002; 46: 1210-1216.
33. Stilund M, Reuschlein AK, Christensen T, Moller HJ, Rasmussen PV, Petersen T. Soluble CD163 as a marker of macrophage activity in newly diagnosed patients with multiple sclerosis. *PLoS One* 2014; 9: e98588.
34. Siebelt M, Korthagen N, Wei W, Groen H, Bastiaansen-Jenniskens Y, Muller C, et al. Triamcinolone acetonide activates an anti-inflammatory and folate receptor-positive macrophage that prevents osteophytosis in vivo. *Arthritis Res Ther* 2015; 17: 352.
35. Mercalli A, Calavita I, Dugnani E, Citro A, Cantarelli E, Nano R, et al. Rapamycin unbalances the polarization of human macrophages to M1. *Immunology* 2013; 140: 179-190.
36. Badlani N, Oshima Y, Healey R, Coutts R, Amiel D. Use of bone morphogenic protein-7 as a treatment for osteoarthritis. *Clin Orthop Relat Res* 2009; 467: 3221-3229.
37. Sekiya I, Tang T, Hayashi M, Morito T, Ju YJ, Mochizuki T, et al. Periodic knee injections of BMP-7 delay cartilage degeneration induced by excessive running in rats. *J Orthop Res* 2009; 27: 1088-1092.
38. Hurtig M, Chubinskaya S, Dickey J, Rueger D. BMP-7 protects against progression of cartilage degeneration after impact injury. *J Orthop Res* 2009; 27: 602-611.
39. Carames B, Hasegawa A, Taniguchi N, Miyaki S, Blanco FJ, Lotz M. Autophagy activation by rapamycin reduces severity of experimental osteoarthritis. *Ann Rheum Dis* 2012; 71: 575-581.
40. Takayama K, Kawakami Y, Kobayashi M, Greco N, Cummins JH, Matsushita T, et al. Local intra-articular injection of rapamycin delays articular cartilage degeneration in a murine model of osteoarthritis. *Arthritis Res Ther* 2014; 16: 482.
41. Matsuzaki T, Matsushita T, Tabata Y, Saito T, Matsumoto T, Nagai K, et al. Intra-articular administration of gelatin hydrogels incorporating rapamycin-micelles reduces the development of experimental osteoarthritis in a murine model. *Biomaterials* 2014; 35: 9904-9911.

42. Zhang O, Zhang J. Atorvastatin promotes human monocyte differentiation toward alternative M2 macrophages through p38 mitogen-activated protein kinase-dependent peroxisome proliferator-activated receptor gamma activation. *Int Immunopharmacol* 2015; 26: 58-64.
43. Ma W, Liu Y, Wang C, Zhang L, Crocker L, Shen J. Atorvastatin inhibits CXCR7 induction to reduce macrophage migration. *Biochem Pharmacol* 2014; 89: 99-108.
44. Dombrecht EJ, Van Offel JF, Bridts CH, Ebo DG, Seynhaeve V, Schuerwegh AJ, et al. Influence of simvastatin on the production of pro-inflammatory cytokines and nitric oxide by activated human chondrocytes. *Clin Exp Rheumatol* 2007; 25: 534-539.
45. Chang CH, Hsu YM, Chen YC, Lin FH, Sadhasivam S, Loo ST, et al. Anti-inflammatory effects of hydrophilic and lipophilic statins with hyaluronic acid against LPS-induced inflammation in porcine articular chondrocytes. *J Orthop Res* 2014; 32: 557-565.
46. Aktas E, Sener E, Gocun PU. Mechanically induced experimental knee osteoarthritis benefits from anti-inflammatory and immunomodulatory properties of simvastatin via inhibition of matrix metalloproteinase-3. *J Orthop Traumatol* 2011; 12: 145-151.
47. Barbosa CP, Ritter AM, da Silva LG, Grespan R, Cuman RK, Hernandes L, et al. Effects of simvastatin, ezetimibe, and their combination on histopathologic alterations caused by adjuvant-induced arthritis. *Inflammation* 2014; 37: 1035-1043.
48. Clockaerts S, Van Osch GJ, Bastiaansen-Jenniskens YM, Verhaar JA, Van Glabbeek F, Van Meurs JB, et al. Statin use is associated with reduced incidence and progression of knee osteoarthritis in the Rotterdam study. *Ann Rheum Dis* 2012; 71: 642-647.
49. Lv S, Liu Y, Zou Z, Li F, Zhao S, Shi R, et al. The impact of statins therapy on disease activity and inflammatory factor in patients with rheumatoid arthritis: a meta-analysis. *Clin Exp Rheumatol* 2015; 33: 69-76.
50. Dong W, Vuletic S, Albers JJ. Differential effects of simvastatin and pravastatin on expression of Alzheimer's disease-related genes in human astrocytes and neuronal cells. *J Lipid Res* 2009; 50: 2095-2102.




**Table 1: Donor demographics and culture conditions of synovial explants**

<b>Donor</b>	<b>Gender</b>	<b>Age</b>	<b>Non-stimulated/IFN<math>\gamma</math>+TNF<math>\alpha</math> stimulated</b>	<b>Compound treatment</b>
1	female	42	Non-stimulated IFN $\gamma$ +TNF $\alpha$ stimulation	None (analyzed after 24h)
2	female	74	Non-stimulated IFN $\gamma$ +TNF $\alpha$ stimulation	None (analyzed after 24h)
3	male	65	Non-stimulated IFN $\gamma$ +TNF $\alpha$ stimulation	None (analyzed after 24h)
4	male	65	Non-stimulated IFN $\gamma$ +TNF $\alpha$ stimulation	None (analyzed after 24h) DMEM (vehicle control) DMSO (vehicle control) Dexamethasone Rapamycin BMP-7 Pravastatin
5	male	65	Non-stimulated IFN $\gamma$ +TNF $\alpha$ stimulation	DMEM (vehicle control) DMSO (vehicle control) Dexamethasone Rapamycin BMP-7 Pravastatin
6	male	65	Non-stimulated IFN $\gamma$ +TNF $\alpha$ stimulation	DMEM (vehicle control) DMSO (vehicle control) Dexamethasone Rapamycin BMP-7 Pravastatin
7	male	64	Non-stimulated IFN $\gamma$ +TNF $\alpha$ stimulation	DMEM (vehicle control) DMSO (vehicle control) Dexamethasone Rapamycin BMP-7 Pravastatin




**Figure 1: Gene expression profile of osteoarthritic synovium with and without IFN $\gamma$ +TNF $\alpha$  stimulation.**


**A)** Expression of genes encoding for pro-inflammatory proteins (*IL1B*, *IL6* and *TNFA*) and genes encoding for anti-inflammatory proteins (*IL1RA*, *CCL18*, *CD206* and *CD163*) relative to the expression of Glyceraldehyde-3-phosphate dehydrogenase (*GAPDH*). **B)** An M1/M2-index based on expression of all measured pro-inflammatory genes and anti-inflammatory genes to provide an overall assessment of the inflammatory state of the synovium with or without IFN $\gamma$ +TNF $\alpha$  stimulation. Data is presented as boxplots with whiskers from minimum to maximum for  $n=4$  donors in triplicate.



**Figure 2: Modulation of stimulated and non-stimulated osteoarthritic (OA) synovial tissue. (A)** Gene expression profiles of IFN $\gamma$ +TNF $\alpha$  stimulated OA synovium, and **(B)** non-stimulated OA synovium after treatment with dexamethasone, rapamycin, BMP-7 or pravastatin relative to the expression levels of the vehicle controls as represented by the dotted line. 0.01% DMSO was used as vehicle control for dexamethasone and rapamycin and X-VIVO medium was used as vehicle control for BMP-7 and pravastatin. The M1/M2-index provides an overview of the overall inflammatory state of the synovial tissue after treatment with the compounds based on expression of all the measured pro-inflammatory genes and anti-inflammatory genes. Data is presented in boxplots with whiskers from minimum to maximum for  $n=4$  donors in triplicate.



**Figure 3: Characterization of primary human macrophages stimulated with IFN $\gamma$  and TNF $\alpha$  (M(IFN $\gamma$ +TNF $\alpha$ )), IL-4 (M(IL-4)) and IL-10 (M(IL-10)). (A) Gene expression profile relative to *GAPDH* expression and (B) protein production of IL-6, CCL18 and sCD163 corrected for amount of DNA. Data is shown as mean  $\pm$  SD for  $n=3$  donors in 5-fold. The error bars represent the variation between donors.**



**Figure 4: Modulation of primary polarized macrophages.** IL-6, CCL18 and sCD163 protein production of M(IFN $\gamma$ +TNF $\alpha$ ), M(IL-4) and M(IL-10) macrophages after modulation with **A**) dexamethasone, **B**) rapamycin, **C**) BMP-7, and **D**) pravastatin. Data is presented for  $n=3$  independent experiments in triplicate as boxplots with whiskers from minimum to maximum and relative to the compound vehicle controls represented as a dotted line. 0.01% DMSO was used as vehicle control for dexamethasone and rapamycin and X-VIVO medium was used as vehicle control for BMP-7 and pravastatin.

**Supplementary Table S1: Primers and probes used for qRT-PCR analysis**

<b>Gene</b>	<b>Primer sequences</b>
<i>IL6</i>	Fw: TCGAGCCCACCGGGAACGAA Rev: GCAGGGAAGGCAGCAGGCAA
<i>IL1B</i>	Fw: CCCTAAACAGATGAAGTGCTCCTT Rev: GTAGCTGGATGCCGCCAT
<i>TNFA</i>	Fw: GCCGCATCGCCGTCTCCTAC Rev: AGCGCTGAGTCGGTCACCCT
<i>CCL18</i>	Fw: GCACCATGGCCCTCTGCTCC Rev: GGGCACTGGGGGCTGGTTTC
<i>IL1RA</i>	Fw: AACAGAAAGCAGGACAAGCG Rev: CCTTCGTCAGGCATATTGGT
<i>CD206</i>	Fw: TGGCCGTATGCCGGTCACTGTTA Rev: ACTTGTGAGGTCACCGCCTTCCT
<i>CD163</i>	Fw: GCGGGAGAGTGGAAGTGAAAG Rev: GTTACAAATCACAGAGACCGCT
<i>TLR4</i>	Fw: GGCATGCCTGTGCTGAGTT Rev: CTGCTACAACAGATACTACAAGCACACT
<i>IL10</i>	Fw: CCTGGAGGAGGTGATGCCCCA Rev: GACAGCGCCGTAGCCTCAGC
<i>TGFB1</i>	Fw: GTGACAGCAGGGATAACACACTG Rev: CATGAATGGTGGCCAGGTC Probe: ACATCAACGGGTTCACTACCGGC
<i>GAPDH</i>	Fw: CAACGGATTTGGTCGTATTGGG Rev: TGCCATGGGTGGAATCATATTGG Probe: GGCGCCCAACCAGCC