



Regulation of transferrin receptor synthesis by human cytotrophoblast cells in culture

M.J. Kroos, J.S. Starreveld, C.E.H. Verrijt, H.G. van Eijk, J.P. van Dijk*

Department of Chemical Pathology, Erasmus University Rotterdam, P.O. Box 1738, 3000 DR Rotterdam, The Netherlands

Received 27 September 1995; revision received 9 November 1995; accepted 28 November 1995

Abstract

The aim of this study was to examine the capacity of the syncytiotrophoblast to regulate transferrin receptor (TfR) synthesis in response to modulations in maternal iron supply. The model used was the primary trophoblast cell culture. Trophoblast cells isolated from term human placentas were cultured in iron-poor (Medium 199), iron-depleted (desferrioxamine (DFO)) and iron-supplemented (diferric transferrin (hTf-2Fe), ferric ammonium citrate (FAC)) medium. TfR synthesis was reduced in response to hTf-2Fe supplementation. FAC did not modulate TfR synthesis. Iron deprivation by DFO resulted in clear stimulation of TfR synthesis. These results show that the differentiating trophoblast cells respond to perturbations in the (transferrin-mediated) iron supply by adjustments in the rate of TfR synthesis. Taking syncytiotrophoblast in culture as model for the maternal/fetal interface *in vivo*, our results would suggest that the placenta is able to make short term adjustments of the capacity for iron uptake.

Keywords: Human; Syncytiotrophoblast; Transferrin; Receptor; Synthesis; Iron

1. Introduction

The growing fetus needs increasing amounts of iron. Maternal diferric transferrin (Tf-2Fe) is taken up by the placental syncytiotrophoblast by means of (transferrin) receptor mediated endocytosis [1]. After release of iron from the so formed endosomes to the cytosol, the endosomes recycle, fuse with the microvillous membrane and apo-Tf is set free into the maternal circulation. The processes involved in the subsequent transfer of iron through the cytosol and across the basal plasma membrane are unknown [1–3].

It was shown previously, that the increase of placental iron transfer during pregnancy is related to an increase of the number of transferrin receptors (TfR) on the expanding maternal/fetal interface. The Tf/TfR affinity is not affected [4].

What factors influence the number of TfRs? Placental

growth follows a developmental programme. This of course does not exclude the possibility that parameters like TfR density can, to some degree, be adjusted to the momentary needs. It is to be expected that the growing placenta responds to reductions in the maternal iron supply with an increase of the microvillous TfR number or density. An increased maternal iron supply will probably induce a reduction of TfR number or density. These adjustments can in principle be achieved by TfR redistribution between cell interior and cell surface [5], and/or by adjustments in the rate of TfR synthesis.

Using the primary culture of trophoblast cells from term human placentas [6,7], we investigated the influences of iron supplementation and withdrawal on denovo TfR synthesis. It is shown that, conforming with the Iron Responsive Element (IRE) hypothesis [8–10], supplementation of the culture media with hTf-2Fe reduces, and iron withdrawal with DFO stimulates synthesis. Supplementation with anorganic ferric ammonium citrate (FAC) has no effect. The results are discussed in the context of the regulation of placental iron transfer.

* Corresponding author, Fax: +31 10 4360615; email: vandijk@chepa.fgg.eur.nl.

2. Materials and methods

2.1. Chemicals

Calcium- and magnesium-free solutions of Eagle's balanced salts (EBBS), fetal calf serum (FCS), Dulbecco's modification of Eagle's medium with 20 mM HEPES (DMEM-H), medium 199 (M199), penicillin, streptomycin and amphotericin were obtained from Flow Labs (ICN Biomedicals, Zoetermeer, Netherlands). Modified Eagle's medium without L-methionine and L-glutamine (MEM) was obtained from Gibco, UK; gentamicin was from Schering Corp., USA; trypsin (1:250 t.c.) was from Sigma Chemical Company, St Louis, USA and DNase grade II was from Boehringer Mannheim, Germany; aprotinin, L-methionine and PMSF were from Sigma, USA; percoll was from Pharmacia, Upsala, Sweden and ^{35}S -translabelled methionine was purchased from ICN Biomedicals, USA. Mouse anti-CD9 and goat anti-mouse IgG magnetic beads were purchased from DAKO (ITK Diagnostics BV, Uithoorn, Netherlands). All other chemicals and reagents (highest purity) were from commercial sources. Human (apo)transferrin (hTf) was obtained from Behringwerke (Marburg a.d. Lahn, Germany).

2.2. Cytotrophoblast-cell isolation

Human term placentas from pregnancies without complications were obtained from the Department of Obstetrics, University Hospital Rotterdam/Dijkzigt, within half an hour after spontaneous delivery. These placentas were processed according to [2] with slight modifications. Approximately 30 g villous tissue was preincubated for 10 min at 37°C with 50 ml of the enzyme solution as used by [2]. The supernatant was discarded and the remaining villous tissue was digested following the original procedure [2]. Cells obtained were counted (using a Bürker counting-chamber) and diluted to 6×10^5 cells/ml in culture medium (see further). This cell-suspension was plated out in 35-mm diameter Falcon culture dishes (Greiner and Söhne, Germany) at a density of 1.5×10^6 cells/dish. Culture medium consisted of 80% v/v M199; 20% v/v FCS; 4 mM L-glutamine; 0.3 mg/ml gentamicin; 50 IU/ml penicillin; 50 µg/ml streptomycin and 2.5 µg/ml amphotericin. Osmolarity was between 280 and 300 mosmol/kg, pH 7.4. The medium was sterilized by filtration on 0.22 µm Millipore-GS filter (Millipore SA Molsheim, France). Cultures were incubated at 37°C in humidified 5% $\text{CO}_2/95\%$ air.

2.3. Cell culture conditions

The cells in culture were allowed to recover from the

isolation procedure for 24 h. In order to remove non-adherent cells, the dishes were washed four times with 2 ml phosphate buffered saline (PBS). Cell culture was continued another 24 h in 1.5 ml fresh culture medium before denovo synthesis of TfR was assessed. These fresh media were fortified with iron (hTf-2Fe (1.25 mM) or FAC (10 µg/l)) or iron depleted by means of 50 µM DFO. Control series: M199 without additions.

2.4. Transferrin receptor synthesis

Cells were washed twice with PBS. Methionine-free medium was added and the cells were incubated for 1 h at 37°C, followed by 2 h incubation with 60–80 µCi translabelled ^{35}S -methionine (1089 Ci/mmol) per 1.5×10^6 cells, also in methionine-free medium. The medium was removed and the cells were washed twice with 2 ml PBS. The cells were lysed for 30 min at 4°C in 300 µl lysate buffer consisting of 1% Triton X-100, 1 mM phenyl-methyl-sulfon-fluoride (PMSF) and 0.5 U/ml aprotinin in PBS pH 7.4. To recover all cell material the dishes were washed twice with 100 µl lysate buffer. Lysis was checked using an inverted microscope. The cell-lysates were centrifuged for 2 min at 10 000 g (r_{av} , 5 cm). Bovine serum albumin (4 mg/ml) and L-methionine (10^4 times the original ^{35}S -methionine concentration) were added to 300 µl supernatant of all the samples. TfR precipitation was achieved with 50 µl [hTf-2Fe]-Sephacrose-4B (Tf 100% saturated with iron), followed by gentle rotation of the samples during 18 h at 4°C. The samples were spun down (5 min, 1000 g) and the precipitates were washed four times with PBS containing 0.2% sodium-desoxycholate, 0.2% Triton X-100, 1 mM PMSF, and 1 mg/ml BSA. The final precipitates were boiled for 5 min with electrophoresis sample buffer (4 ml distilled water, 1 ml 0.5 M Tris-HCl pH 6.8, 0.8 ml glycerol, 1.6 ml 10% SDS, 0.4 ml 2β-mercaptoethanol, 0.2 ml bromophenol blue), and centrifuged for 2 min at 10 000 g (r_{av} , 5 cm). The supernatants were electrophoresed under reducing conditions, (Mini-Protean II Cell, Bio-Rad) using 10% homogenous ready gels (Bio-Rad). After drying, the gels were exposed to Kodak X-OMAT AR film.

2.5. Total protein synthesis

To 0.02 ml of the cell lysate, 0.1 ml distilled water and 0.1 ml ice-cold 20% trichloroacetic acid (TCA) was added. After 10 min on ice, the samples were centrifuged for 15 min at 1500 g and 4°C. The precipitate was washed with 0.05 ml 10% TCA and subsequently vigorously mixed with 0.5 ml Soluene-350 (Packard). A sample of 0.02 ml was carefully transferred to liquid-scintillation bottles, and 10 ml Instagel (Packard) was added. After vigorous mixing and clearing up, the radiochemical concentration was measured in an Isocap 300 counter.

Counting efficiency was always >85%. The amount of radiation (dpm) present in the total TCA precipitate was taken as a measure for total protein synthesis.

2.6. Statistics

The significance of the results was tested by using the Student-Newman-Keuls test ($\alpha_T < 0.05$). This test corrects the outcomes for multiple comparisons.

3. Results

A typical fluorogram is shown in Fig. 1. It demonstrates the influence of manipulation of the iron content of the culture media on denovo TfR synthesis. Notice that enrichment of the medium with hTf-2Fe gives inhibition of the receptor synthesis, whereas supplementation with FAC is ineffective. Iron deprivation with DFO stimulates TfR synthesis.

The semiquantitative data obtained by laser-densitometry of the fluorograms are given in Fig. 2. Again, compared to the time controls, hTf-2Fe acts inhibitory, DFO stimulatory, and FAC does not seem to influence TfR synthesis. The data are expressed relative to total protein synthesis. Compared to both the control and the FAC series, TfR synthesis was significantly lower in the




Fig. 2. Influence of culture time, iron supplementation and iron depletion on TfR synthesis. Quantification was achieved by densitometry of the developed films. The data are expressed relatively to total protein synthesis. Mean \pm S.D. of four independent experiments including one experiment with cells additionally immunopurified with anti-CD9 [14] are given.

hTf-2Fe series ($\alpha_T < 0.02$), and significantly increased in the DFO series ($\alpha_T < 0.001$). Culture time did not affect TfR synthesis.

4. Discussion

It was previously shown that cytotrophoblast cells in culture express TfRs and that the total number of TfRs as well as the number of surface TfRs increased with culture time [2,5,7].

Expression of TfRs is part of the biochemical differentiation of the trophoblast cells towards syncytiotrophoblast or syncytiotrophoblast-like structures [7,12]. The total number of TfRs (as well as the surface TfR population) could, to some degree, be up- or down-regulated, depending on the culture conditions with respect to iron depletion or iron supplementation [5,13,14]. This suggests that the differentiating trophoblast cell is able to adjust the rate of TfR synthesis to the momentary needs.

The results obtained in this investigation clearly prove that TfR synthesis can be adaptively modified. Iron fortification of the culture media with hTf-2Fe reduced denovo TfR synthesis, whereas iron deprivation with desferrioxamine (DFO) induced a stimulation. Supplementation with ferric ammonium citrate (FAC) appeared to be ineffective.

For many cell types it was previously shown that adaptive change in the rate of both ferritin and TfR synthesis is regulated at the post-transcriptional level by iron. Increase of low molecular weight (lmw) iron in the cell stimulates the synthesis of ferritin and — at the same time — decreases TfR synthesis. Decrease of lmw-iron leads to the opposite events. These opposite effects of




Fig. 1. Trophoblast transferrin receptor synthesis. Cells were isolated and cultured as described in materials and methods. TfR synthesis was assessed by ³⁵S-methionine pulse labelling and precipitation with [hTf-2Fe]-Sephrose-4B. After SDS-electrophoresis (reducing conditions) the gels were dried and exposed to X-ray film. The monomeric form of the homodimeric TfR (mol-weight homomere: 90.000 kDal) is indicated by an arrow. Lane 1: 42 h hTf-2Fe supplementation; lane 2: 42 h DFO supplementation; lane 3: 42 h FAC supplementation; lane 4: 42 h control. The results of one out of four representative experiments are shown.

iron are explained by the Iron Responsive Element (IRE) hypothesis [8–10]. The effect of iron is mediated by an Iron Responsive Element Binding Protein (IRE-BP). This IRE-BP has 50% homology with mitochondrial aconitase [11]. In the presence of iron this protein has aconitase activity due to the 4FeS clusters formed. In the absence of iron no FeS-clusters can be formed and the protein loses its aconitase activity. At the same time binding of IRE-BP to the IREs on the mRNA of the ferritin (5'-untranslated region) and TfR (3'-untranslated region) is enabled. This causes a reduction of ferritin synthesis and — at the same time — an increase of the TfR synthesis, due to improved stability of TfR mRNA. Reduction of intracellular low-molecular-weight-iron leads to the reversed cascade.

In terms of the IRE hypothesis we would expect that hTf-2Fe is taken up easily, resulting in an increased low-molecular-weight-iron concentration and an IRE sensitive reduction of TfR mRNA translation. The negative results obtained with FAC could be explained by a restricted availability of anorganic chelated iron. However, this explanation of the results fails. As previously shown the trophoblast cells take up FAC very rapidly, resulting in a clear increase in ferritin iron load and stimulation of *de novo* ferritin synthesis [14]. In contrast, supplementation of M199 with hTf-2Fe only resulted in a marginal rise of the cellular iron content [14,15], despite the fact that the rate of iron uptake from hTf-2Fe was about 80–110 pmol/h/mg cell protein [2,14,15]. The absence of iron accumulation upon hTf-2Fe fortification, implicates iron release from the cells, without storage above the direct cellular needs [14,15]. In case of FAC supplementation, the syncytiotrophoblast is unable to keep intracellular iron low, and ferritin synthesis is stimulated.

The results suggest that the term syncytiotrophoblast can balance Tf mediated iron uptake and iron release. Reduction of surface TfRs by inhibition of *de novo* TfR synthesis may be one of the mechanisms involved in this regulatory process.

References

- [1] van Dijk JP. Review article: regulatory aspects of placental iron transfer — a comparative study. *Placenta* 1988; 8: 347–354.
- [2] Douglas GC, King BF. Uptake and processing of ¹²⁵I-labelled transferrin and ⁵⁹Fe-labelled transferrin by isolated human trophoblast cells. *Placenta* 1990; 11: 41–57.
- [3] Harris ED. Brief critical review: new insights into placental iron transport. *Nutr Rev* 1992; 50: 329–337.
- [4] van Dijk JP, van der Zande FGM, Kroos MJ, Starreveld JS, van Eijk HG. Number and affinity of transferrin-receptors at the placental microvillous plasma membrane of the guinea pig: influence of gestational age and degree of transferrin glycan chain complexity. *J Dev Physiol* 1993; 19: 221–226.
- [5] Starreveld JS, van Dijk JP, Kroos MJ, van Eijk HG. Regulation of transferrin receptors expression and distribution in *in vitro* cultured human cytotrophoblasts. *Clin Chim Acta* 1993; 220: 47–60.
- [6] Kliman HJ, Nestler JE, Sernasi E, Sanger JM, Strauss III JF. Purification, characterisation and *in vitro* differentiation of cytotrophoblasts from human term placentae. *Endocrinology* 1986; 118: 1567–1582.
- [7] Bierings MB, Adriaansen J, van Dijk JP. The appearance of transferrin receptors on cultures human cytotrophoblast and *in vitro* formed syncytiotrophoblast. *Placenta* 1988; 9: 387–396.
- [8] Kuhn LC, Hentze MW. Coordination of cellular iron metabolism by post-transcriptional gene regulation. *J Inorg Biochem* 1992; 47: 183–195.
- [9] Meleforts O, Hentze MW. Iron regulatory factor — the conductor of cellular iron regulation. *Blood* 1993; 7: 251–258.
- [10] Mattia E, Renswoude J. The pivotal role of ferritin in cellular iron homeostasis. *Bioassays* 1988; 8: 107–111.
- [11] Haile DJ, Raoult TA, Tang Chin J, Haford JB, Klausner RD. Reciprocal control of RNA-binding and aconitase activity in regulation of the iron responsive element binding protein: role of the iron-sulfur cluster. *Proc Natl Acad Sci USA* 1992; 89: 7536–7540.
- [12] Bierings MB, Baert MRM, van Eijk HG, van Dijk JP. Transferrin receptor expression and the regulation of placental iron uptake. *Mol Cell Biochem* 1991; 100: 31–38.
- [13] Bierings MB, van Eijk HG, Baert MRM, van Dijk JP. Regulation of transferrin receptor expression in term human cytotrophoblasts. *Trophoblast Res* 1992; 6: 237–248.
- [14] Starreveld JS, Kroos MJ, van Suijlen JDE, Verrijt CEH, van Eijk HG, van Dijk JP. Ferritin in cultured human cytotrophoblasts: synthesis and subunit distribution. *Placenta* 1995; 16: 383–395.
- [15] Starreveld JS, Kroos MJ, van Dijk JP, van Eijk HG. The effects of iron supplementation on iron uptake by differentiating cytotrophoblasts. *Reprod Fertil Dev* 1996. Accepted for publication.