

LETTER

Response to 'T-helper 17 cell cytokines and interferon type I: partners in crime in systemic lupus erythematosus?' – Authors' reply

Zana Brkic¹, Odilia BJ Corneth^{1,2}, Marjan A Versnel^{1†} and Erik Lubberts^{1,2*†}

See related research by Brkic et al., <http://arthritis-research.com/content/16/2/R62> and related letter by Dolff et al., <http://arthritis-research.com/content/16/3/409>

We would like to reply to the letter by Dolff and colleagues [1] regarding our article in a recent issue of *Arthritis Research & Therapy* [2]. We thank the authors for their interest in our work and the critical reading of our article. The aim of our study was to investigate a possible association between the IFN type I signature and memory T helper 17 (Th17) cells and their cytokines in systemic lupus erythematosus (SLE). Since CCR6 can be expressed by regulatory T (Treg) cells, we have excluded CD25^{high} cells to discriminate between CD4⁺CD45RO⁺CCR6⁺CD25⁻ (primary Th17 cells) and Treg cells. Concerning the observation on co-expression of IL-17A with IFN- γ , we would like to remark that IFN- γ is an IFN type II, and not an IFN type I, cytokine and does not bind to the IFN type I receptor. Detection of IFN type I activity is hampered by the difficulty to assess the different subtypes of this cytokine. Therefore, analysis of IFN type I-induced gene expression in RNA from peripheral blood cells, the so-called IFN type I signature, is used as a measure for IFN type I activity. Although we fully agree with the authors that the relationship between Th cells and IFN type I deserves further study, their remark on the 'genetic' signature is confusing and probably refers to the IFN type I-induced gene expression signature, which is detected at the RNA level.

The relationship between CD25⁺ Tregs and IFN type I is certainly of interest for further study as these Tregs are carrying the IFN type I receptor and thus respond to increased systemic IFN type I activity in SLE. Also, follow-up studies with a focus on adaptive and innate cells producing

IL-17, including the relation with IL-21 and IL-22 and plasticity, will be of interest [3, 4]. In our article, we presented data supporting a potential co-activity between IFN type I and CD4⁺memory CCR6⁺ Th17 cytokines. Further studies are needed to confirm this co-activity with a focus on revealing the mechanism of this dangerous link in SLE.

Abbreviations

IFN: Interferon; IL: Interleukin; SLE: Systemic lupus erythematosus; Th17: T helper 17; Treg: Regulatory T.

Competing interests

The authors declare that they have no competing interests.

Published: 12 June 2014

References

1. Dolff S, Abdulahad WH, Kallenberg CGM: Response to 'T-helper 17 cell cytokines and interferon type I: partners in crime in systemic lupus erythematosus?'. *Arthritis Res Ther* 2014, **16**:409.
2. Brkic Z, Corneth OB, van Helden-Meeuwsew CG, Dolhain RJ, Maria NI, Paulissen SM, Davelaar N, van Hamburg JP, van Daele PL, Dalm VA, van Hagen PM, Hazes JM, Versnel MA, Lubberts E: T helper 17 cell cytokines and interferon type I: partners in crime in systemic lupus erythematosus? *Arthritis Res Ther* 2014, **16**:R62.
3. Acosta-Rodriguez EV, Rivino L, Geginat J, Jarrossay D, Gattorno M, Lanzavecchia A, Sallusto F, Napolitani G: Surface phenotype and antigenic specificity of human interleukin 17-producing T helper memory cells. *Nat Immunol* 2007, **8**:639–646.
4. Hirota K, Duarte JH, Veldhoen M, Hornsby E, Li Y, Cua DJ, Ahlfors H, Wilhelm C, Tolaini M, Menzel U, Garefalaki A, Potocnik AJ, Stockinger B: Fate mapping of IL-17-producing T cells in inflammatory responses. *Nat Immunol* 2011, **12**:255–263.

doi:10.1186/ar4578

Cite this article as: Brkic et al.: Response to 'T-helper 17 cell cytokines and interferon type I: partners in crime in systemic lupus erythematosus?' – Authors' reply. *Arthritis Research & Therapy* 2014 **16**:410.

* Correspondence: e.lubberts@erasmusmc.nl

[†]Equal contributors

¹Department of Immunology, Erasmus MC, University Medical Center, Wytemaweg 80, 3015 CN Rotterdam, The Netherlands

²Department of Rheumatology, Erasmus MC, University Medical Center, Wytemaweg 80, 3015 CN Rotterdam, The Netherlands