



Perspective

National survey data for zoonotic schistosomiasis in the Philippines grossly underestimates the true burden of disease within endemic zones: implications for future control



Remigio M. Olveda^a, Veronica Tallo^a, David U. Olveda^b, Marianne T. Inobaya^{a,b},
Thao N. Chau^c, Allen G. Ross^{b,*}

^a Research Institute for Tropical Medicine, Department of Health, Manila, The Philippines

^b Menzies Health Institute Queensland, Griffith University, Gold Coast, Australia

^c Discipline of Public Health, Flinders University, Adelaide, Australia.

ARTICLE INFO

Article history:

Received 4 November 2015

Received in revised form 13 January 2016

Accepted 19 January 2016

Corresponding Editor: Eskild Petersen, Aarhus, Denmark.

Keywords:

Schistosomiasis

Control

Retrospective national data

GIS risk maps

Policy

SUMMARY

Zoonotic schistosomiasis has a long endemic history in the Philippines. Human mass drug administration has been the cornerstone of schistosomiasis control in the country for the past three decades. Recent publications utilizing retrospective national survey data have indicated that the national human prevalence of the disease is <1%, hence the disease is now close to elimination. However, the evidence for such a claim is weak, given that less than a third of the human population is currently being treated annually within endemic zones and only a third of those treated actually swallow the tablets. For those who consume the drug at the single oral dose of 40 mg/kg, the estimated cure rate is 52% based on a recent meta-analysis. Thus, approximately 5% of the endemic human population is in reality receiving the appropriate treatment. To compound this public health problem, most of the bovines in the endemic communities are concurrently infected but are not treated under the current national control programme. Given this evidence, it is believed that the human prevalence of schistosomiasis within endemic regions has been grossly underestimated. Inherent flaws in the reporting of national schistosomiasis prevalence data are reported here, and the problems of utilizing national retrospective data in making geographic information system (GIS) risk maps and advising policy makers of the outcomes are highlighted.

© 2016 The Authors. Published by Elsevier Ltd on behalf of International Society for Infectious Diseases. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

1. Introduction

Schistosomiasis, or Bilharzia, is a neglected tropical parasitic disease caused by blood flukes of the genus *Schistosoma*. Globally, it ranks third among the most devastating tropical diseases, and is a major cause of morbidity in the tropics. Schistosomiasis was first reported in the Philippines in 1906.¹ Approximately 12 million people, residing in 28 endemic provinces located in 12 different geographical zones, are at risk of infection in the country.² Overall, a total of 190 municipalities and 1212 barangays (villages) are currently endemic for the disease based on surveys conducted over the past decade. Two new endemic foci reported in the northern (Gonzaga, Cagayan) and central (Calatrava, Negros Occidental)

parts of the country were confirmed in 2004 and 2006, respectively.³

Considerable optimism surrounds mass drug administration (MDA) for the control of schistosomiasis globally, for which praziquantel has served as the cornerstone since its development in 1979.⁴ Numerous studies have claimed that preventive chemotherapy (i.e., 40 mg/kg of praziquantel), given annually or biannually, can significantly reduce the prevalence and intensity of infection, and control morbidity in the long term.⁴ In the last decade, close to one billion US dollars has been raised for MDA campaigns against neglected tropical diseases, largely from international donors, and delivered vertically to local endemic communities through national health care services.⁴

There is now strong evidence that large mammals (e.g., water buffaloes, cattle, and dogs) are contributing significantly to disease transmission and complicating control.⁴ Given the zoonotic nature of the disease in the Philippines it is evident that the incidence,

* Corresponding author: Griffith University, Logan Campus, University Drive, Meadowbrook QLD 4131, Australia. Tel.: +61 733821098.
E-mail address: a.ross@griffith.edu.au (A.G. Ross).

prevalence, and morbidity of this disease will not be controlled by MDA alone. There is the need for innovative cost-effective strategies to control schistosomiasis japonica in the long term. The inherent flaws in the reporting of the national schistosomiasis prevalence data are described herein, and the problems of utilizing national retrospective data in making geographic information system (GIS) risk maps and advising policy-makers of the outcomes is highlighted.

2. National schistosomiasis surveillance

The control programme for schistosomiasis in the Philippines has been under the supervision of the Schistosomiasis Control Service (SCS). The human prevalence of *Schistosoma japonicum* infection was defined as the proportion of individuals who showed at least one parasite egg on two Kato–Katz (KK) thick smears versus the number of people examined in the endemic population. The estimated exposed population for schistosomiasis in the Philippines is approximately 1.8 million.² The annual results of case finding are shown in Table 1. From 1985 to 1990, approximately 30% (range 26–37%) of the target population was examined. In 1990, the SCS received additional funds from the World Bank through the Philippine Health Development Programme (PHDP), and from 1991 to 1993 it managed to examine approximately 75% of the target population (range 75–79%) with the increased funding. However, funds through the PHDP were subsequently reduced in 1994 and funding was eventually terminated in 1995. As a direct result, the proportion of individuals examined annually from 1994 to 1999 decreased to approximately 25% (range 22–44%). The percentage of individuals examined plummeted further from 2000 to the present.^{5–7} In sum, due to limited financial resources, active surveillance for schistosomiasis in the Philippines now comprises approximately 10% of the known endemic population in the country. This has resulted in a gross underreporting of the disease given its highly focal nature and clumped distribution.

With support from World Health Organization (WHO), a stratified two-step systematic cluster sampling survey was conducted from 2005 to 2008 in order to determine the national prevalence of schistosomiasis.⁸ Provinces in both endemic and non-endemic regions were identified for inclusion in the survey. Five municipalities were selected randomly from the selected provinces, and one village per municipality was selected for follow-up. Two hundred and seventy-four eligible subjects were

chosen randomly from the selected households in each village, and eligible subjects were requested to submit two stool samples.⁸ From each stool sample, two KK thick smears were examined for the presence of *S. japonicum* eggs. This sampling strategy selected 115 villages out of a total 1212 known endemic villages. Compliance for the first stool submission of eligible subjects was 45% in Mindanao, 60% in the Visayas, and 76% in Luzon.⁸ It is noteworthy that this national survey strategy examined less than 10% of the known endemic population and many individuals refused to provide stool for examination. Thus the generalizability of these findings for 'endemic zones' is questionable and should be viewed with considerable caution. It is well known that the distribution of the disease is highly focal (clumped) in nature, thus the random selection of a small segment of the endemic population (<10%) could lead to a gross underestimation of the true burden. This combined with the fact that only one or two slides were read for each individual raises concerns in the reporting of national prevalence in the Philippines.

3. Problems with the national control programme

Population-based chemotherapy as the sole control strategy for zoonotic schistosomiasis in the Philippines was initiated in 1980. The initial approach was the treatment of positive cases with the drug praziquantel, administered at an oral dose of 60 mg/kg, divided into two equal doses taken 4 h apart. The compliance rate from 1985 to 1999 for annual free treatment was reportedly high at approximately 85% in those who were found positive by case finding. However, it is important to note that only 30% of the endemic population was covered by yearly case finding and treatment. This suggests that 70% of the endemic population was left untreated.

In 2000, the SCS implemented a policy of annual MDA for residents 5–65 years of age residing in endemic villages with a human prevalence of infection greater than 15%. Case finding and treatment was continued in villages with a prevalence of <15%.⁹ In 2009 the MDA policy was changed to provide treatment to all individuals aged 5–65 years residing in endemic villages regardless of the prevalence.¹⁰ In 2011, following the recommendation of the WHO, the SCS reduced the dosage of praziquantel in the MDA programme to a single oral dose of 40 mg/kg.¹¹

Despite the fact that only a third of the infected population was treated annually (except from 1991 to 1993) through the case finding and treatment scheme, and that less than a third complied with the free treatment, the yearly national prevalence of *S. japonicum* in the country was reported to have declined dramatically from a high of 10% in 1985 to a low of 0.49% in 2008. Table 1 illustrates the continuous drop in national human prevalence of schistosomiasis from more than 10% in 1985 to less than 4% in 1999. The prevalence was reportedly maintained at less than 5% until 2005.^{5,6,12} Subsequent prevalence surveys (2005–2008) using the above-mentioned stratified two-step systematic cluster sampling approach, recorded a very low national level of infection of less than 1%.⁸ The remarkably low national prevalence reported for *S. japonicum* infection in the last 10 years has now created the impression that the disease is no longer a serious public health problem by the government and that the continuous implementation of the existing national MDA programme for humans could lead to disease elimination. However, given the poor drug coverage (less than one third of the endemic population covered) and poor compliance with the free annual treatment (only one third of the population actually take the drug), it is difficult to believe that the current national human prevalence has dropped to <1%. Moreover, given the zoonotic nature of the disease, where over 80% of bovines are infected, the likelihood of a

Table 1
Reported human prevalence of schistosomiasis based on national survey data^a

Year	Number of individuals examined	Percentage of target population of 1.8 million individuals	Number positive	Prevalence (%)
1985	505 851	28.6	39 046	10.4
1986	459 291	26	34 150	7.4
1987	683 918	37	44 925	6.6
1988	423 708	24	26 953	6.4
1989	468 355	27	35 197	7.5
1990	528 359	30	34 611	6.6
1991	1 317 933	75	79 500	6.0
1992	1 326 283	75	55 896	4.2
1993	1 402 216	79	54 648	3.9
1994	771 970	44	40 446	5.2
1995	267 346	15	12 091	4.5
1996	374 262	21	15 293	4.1
1997	404 811	23	18 301	4.5
1999	392 133	22	12 871	3.3

^a Note: From 1985 to 1999, case finding was done by examining two Kato–Katz thick smears from a single stool sample. Source of data: Department of Health Schistosomiasis Control Service annual reports, 1997 and 1999.

reduction in the national prevalence is highly unlikely.^{13–15} So, what is the true burden of this disease within endemic zones?

4. National burden of disease

Figure 1 summarizes the national schistosomiasis prevalence for humans reported by Leonardo et al. in 2012, Magalhães et al. in 2014, and by various other authors during 2002–2014 in known endemic zones.^{8,16} The national survey conducted in 2008, which used the stratified two-step systematic cluster sampling scheme, recorded very low prevalence rates in the known highly endemic provinces of Western Samar, Northern Samar, and Leyte, with reported prevalence rates of 0.8%, 2.4%, and 0.9%, respectively.⁸ In the newly identified areas of transmission in Cagayan Province, the recorded prevalence was 0.4%.⁸ The prevalence rates generated by Magalhães et al. in 2014 for the same endemic provinces were 0–3%, 0–3%, 0–5%, and 0.4%.¹⁶ In contrast, other research teams performing studies from 2002 to 2014 reported prevalence rates ranging from 0.7% to 47% for Western Samar, 5% to 46% for Northern Samar, and 25% to 60% for Leyte (Table 2). In Cagayan, two endemic villages were surveyed and the prevalence rates were found to be 10% and 31%, respectively.

Table 2 illustrates the key outcomes of various studies that have investigated the prevalence of schistosomiasis in various endemic regions throughout the country from 2002 to 2014. In 2002, a longitudinal study completed in Leyte, investigating immune correlates of resistance to infection, showed that the human prevalence in those aged 15–30 years was 60% (1262/1699).¹⁷ Another study, which examined *S. japonicum* infection among pregnant women, conducted in Leyte between 2008 and 2012, showed that 25.4% (373/1468) of the subjects were positive for schistosomiasis.¹⁸ A cross-sectional survey of 50 endemic villages in Western Samar Province in 2006, as part of the Schistosomiasis Transmission Ecology Project (STEP), reported prevalence rates ranging from 0.7% to 47% (mean 15%).¹⁹ Another cross-sectional

survey of schistosomiasis was conducted in 2012 on 18 221 individuals residing in 22 schistosomiasis endemic villages in the province of Northern Samar.²⁰ Despite an active schistosomiasis control programme in Northern Samar for over 30 years, which included an MDA campaign in the last 5 years, the mean prevalence of schistosomiasis among 10 435 subjects evaluated was 27.1% (95% confidence interval 26.3–28.0%).²⁰ Moreover, a cross-sectional ultrasound study in the same study location showed high levels of schistosomiasis-induced morbidity in the schistosomiasis endemic communities. Left lobe liver enlargement (≥ 70 mm) was evident in 89.3% of subjects. Twenty-five percent of the study population had grade II/III liver parenchyma fibrosis and 13.3% had splenomegaly (≥ 100 mm).^{20,21} In 2012, two communities in the province of Cagayan (Magrafil and Santa Maria) were screened for the prevalence of schistosome infection and this was found to be 10% and 31%, respectively (personal correspondence).²²

5. Discussion

The recently reported national human prevalence (and GIS risk maps) of schistosomiasis in the Philippines has been estimated to be less than 3% in the endemic regions of the country.^{8,16} As a result, the disease is no longer considered to be a major public health problem by the government. Moreover, it is now recommended that the limited national control funds be spent on more targeted MDA, leading to disease elimination.¹⁶ However, when one examines more closely how the national surveys were conducted and how the current national programme is being deployed, it is difficult to ascertain by what means that national prevalence could have dropped to such a low rate over the past decade. Moreover, other studies over the past decade (2002–2014) have shown evidence that the disease remains highly endemic throughout the country in both human and bovine populations, with reported human prevalence rates as high as 50% in some




Figure 1. Comparison of the human schistosomiasis prevalence rates in highly endemic zones of the Philippines: Leonardo et al.⁸ (left panel), Magalhães et al.¹⁶ (centre panel), and of various other authors for the period 2002–2014 (right panel).

Table 2
Key infection outcomes of studies investigating *Schistosoma japonicum* infection in humans in the Philippines

Year	Study description	Key outcomes	Reference
2000	Immune correlate study on human <i>S. japonicum</i> in a well-defined population in Leyte, Philippines; 1262 out of 1699 individuals aged 15–30 years from three study villages were examined for <i>S. japonicum</i> infection.	Prevalence of infection among 15–30-year-olds was 60%	Acosta et al. ¹⁷
2005	A cross-sectional survey of <i>S. japonicum</i> infection was carried out in 50 irrigated and rain-fed villages in Western Samar Province, Philippines, as part of the Schistosomiasis Transmission Ecology Project (STEP). The survey was performed using a household cluster sampling technique.	Prevalence ranged from 0.7% to 47% (mean 15%)	Tarafder et al. ¹⁹
2008	Between 2008 and 2012 in Leyte, pregnant women were examined for <i>S. japonicum</i> infection as part of the screening process for enrolment in the study entitled, "The impact of praziquantel given at 12–16 weeks of gestation on pregnancy outcomes: a double blind placebo controlled trial".	373 out of 1468 (25.4%) pregnant women were positive for schistosomiasis	Olveda et al. ¹⁸
2012	In Northern Samar Province, cross-sectional surveys were performed to determine the prevalence of <i>S. japonicum</i> infection in 22 endemic villages in the municipalities of Palapag and Laoang, as part of the baseline study for the project entitled, "Towards sustainable control and elimination of schistosomiasis in the Philippines".	Prevalence of <i>S. japonicum</i> infection ranged from 5% to 46% (mean 27%)	Ross et al. ²⁰
2013	A cross-sectional ultrasound survey in schistosomiasis endemic villages in the municipality of Palapag, Northern Samar Province, was performed to determine the prevalence of the different degrees of hepatic fibrosis in the residents under study.	Severe hepato-splenic morbidities including grades II and III schistosomiasis hepatic fibrosis was seen in 18% of the population	Olveda et al. ²¹
2013	In Cagayan valley, in the northern part of the Philippines, a cross-sectional survey of <i>S. japonicum</i> was conducted in two villages (Magrafil and Santa Maria) as part of a baseline study for the project called, "Echohealth approach towards the control of schistosomiasis in the Philippines".	The prevalence of infection in these two villages was 10% and 31%, respectively	Olveda et al. ²²
2014	A diagnostic study of <i>S. japonicum</i> endemic villages in Northern Samar Province was conducted in 2012. Faecal samples were collected from 560 humans and examined by KK method and real-time PCR (qPCR) assay.	A high human prevalence rate (90.2%) was recorded by qPCR, while the KK method indicated a prevalence of 22.9%	Gordon et al. ¹⁵

KK, Kato–Katz thick smear.

localities, and over 80% in bovine populations.^{13–15,20} Given the zoonotic nature of the disease, it is difficult to ignore such evidence. Clearly, this discrepancy in reported findings warrants closer scrutiny given the implications for national control and future policy.

It is noteworthy that the recent national surveys were based on only one or two KK thick smear stool slides for those who provided stool, thus this will lead to a gross underestimation of the true prevalence. It has been estimated that a single KK misses approximately 55% of light infections and that five KK smears would be needed to detect 95% of such infections.²³ 'Intra-individual' variation has been observed in repeated egg counts.²³ Additionally, it has been shown that a small number of individuals in endemic communities excrete a large proportion of eggs. This 'inter-individual' variation in egg counts is thought to reflect the variation in worm loads.²³ Quality control in reading such slides is paramount if one wants to truly determine the underlying prevalence in an endemic population. In the Philippines, based on the personal research experience of the present investigators, the ability of microscopists to correctly identify schistosomiasis eggs varies considerably (0–60%) among trained personnel; thus, if quality measures are not in place, this again will lead to a gross underestimation of the human prevalence.

The national schistosomiasis control programme, using MDA of humans as the sole strategy, has stated that the human prevalence has dropped dramatically over the past decade. However, the evidence for such a claim is weak given that less than a third of the human population residing within endemic zones are currently being treated annually and only a third of those treated actually take the drug. For those who take the drug at the single oral dose of 40 mg/kg, the estimated cure rate is 52% based on a recent meta-analysis.⁴ In sum, approximately 5% of the endemic human population is in reality receiving the appropriate treatment. Moreover, in these same endemic communities, most of the bovines are currently infected (>75%) but are not being treated under the current control programme. Given this evidence, it is believed that the prevalence of schistosomiasis has been grossly

underestimated within endemic areas. Through the present authors' own research, it has been found that the disease is moderately endemic (5–25%) in most endemic regions.^{14,15,17–20}

6. Conclusions

Schistosomiasis remains a major public health problem in the Philippines despite recent published reports. Given the zoonotic nature of the disease, mass drug administration in humans as a sole national control strategy will not lead to disease elimination. An integrated national control programme that carefully considers the role of bovines in human transmission, along with irrigation, focal mollusciciding, annual mass drug administration of both humans (60 mg/kg) and bovines (25 mg/kg), and health education is advocated. Although this intersectoral approach will require a substantial effort, it is the only strategy that will lead to sustainability.

Acknowledgements

We thank the Australian National Health and Medical Research Council and the International Development Research Centre, Canada for providing financial support for our research.

Financial support: Our work is supported by the Australian National Health and Medical Research Council and the International Development Research Centre, Canada.

Conflict of interest: All authors: No reported conflicts. All authors have submitted the ICMJE Form. Conflicts that the editors consider relevant to the content of the manuscript have been disclosed.

References

1. Wooley P. The occurrence of *Schistosoma japonicum* Vel Cattol in the Philippines. *Philippine Journal of Science* 1906;1:83–90.
2. Blas BL, Rosales MI, Lipayon IL, Yasuraoka K, Matsuda H, Hayashi M. The schistosomiasis problem in the Philippines: a review. *Parasitol Int* 2004;53:127–34.

3. Leonardo L, Rivera P, Saniel O, Antonio Solon J, Chigusa Y, Villacorte E, et al. New endemic foci of schistosomiasis infections in the Philippines. *Acta Trop* 2015;**141**(Pt B):354–60.
4. Ross AG, Olveda RM, Li Y. An audacious goal: the elimination of schistosomiasis in our lifetime through mass drug administration. *Lancet* 2015;**30**:2220–1. [http://dx.doi.org/10.1016/S0140-6736\(14\)61417-3](http://dx.doi.org/10.1016/S0140-6736(14)61417-3)
5. Philippine Department of Health, Schistosomiasis Control Service Annual 1997 Report. Department of Health, Manila, The Philippines; 1998.
6. Philippine Department of Health, Schistosomiasis Control Service Annual 1999 Report. Department of Health, Manila, The Philippines; 2000.
7. Philippine Department of Health, Schistosomiasis Control Service Annual 2005 Report. Department of Health, Manila, The Philippines; 2006.
8. Leonardo L, Rivera P, Saniel O, Villacorte E, Lebanan MA, Crisostomo B, et al. A national baseline prevalence survey of schistosomiasis in the Philippines using stratified two-step systematic cluster sampling design. *J Trop Med* 2012;**2012**: 936128.
9. Mass treatment strategy for the control and elimination of schistosomiasis: field operational guide. Manila: Department of Health; 2000.
10. Philippine Department of Health, Schistosomiasis Control Service Annual 2009 Report. Department of Health, Manila, The Philippines; 2010.
11. Philippine Department of Health, Schistosomiasis Control Service Annual 2011 Report. Department of Health, Manila, The Philippines; 2012.
12. Philippine Department of Health, Schistosomiasis Control Service Annual 2004 Report. Department of Health, Manila, The Philippines; 2005.
13. Wu HW, Qin YF, Chu K, Meng R, Liu Y, McGarvey ST, et al. High prevalence of *Schistosoma japonicum* infection in water buffaloes in the Philippines assessed by real-time polymerase chain reaction. *Am J Trop Med Hyg* 2010;**82**:646–52.
14. Gordon CA, Acosta LP, Gray DJ, Olveda RM, Jarilla B, Gobert GN, et al. High prevalence of *Schistosoma japonicum* infection in Carabao from Samar Province, the Philippines: implications for transmission and control. *PLoS Negl Trop Dis* 2012;**6**:e1778.
15. Gordon CA, Acosta LP, Gobert GN, Olveda RM, Ross AG, Williams GM, et al. Real-time PCR demonstrates high prevalence of *Schistosoma japonicum* in the Philippines: implications for surveillance and control. *PLoS Negl Trop Dis* 2015;**9**:e0003483.
16. Soares Magalhães RJ, Salamat MS, Leonardo L, Gray DJ, Carabin H, Halton K, et al. Geographical distribution of human *Schistosoma japonicum* infection in The Philippines: tools to support disease control and further elimination. *Int J Parasitol* 2014;**44**:977–84.
17. Acosta LP, Waive G, Aligui GD, Tiu WU, Olveda RM, McManus DP. Immune correlate study on human *Schistosoma japonicum* in a well-defined population in Leyte, Philippines: II. Cellular immune responses to *S. japonicum* recombinant and native antigens. *Acta Trop* 2002;**84**:137–49.
18. Olveda RM, Acosta LP, Tallo V, Baltazar PI, Lesiguez JL, Estanislao GG, et al. Efficacy and safety of praziquantel for the treatment of human schistosomiasis during pregnancy: a phase 2, randomised, double-blind, placebo-controlled trial. *Lancet Infect Dis* 2015 Nov 2. [http://dx.doi.org/10.1016/S1473-3099\(15\)00345-X](http://dx.doi.org/10.1016/S1473-3099(15)00345-X)
19. Tarafder MR, Balolong Jr E, Carabin H, Bélisle P, Tallo V, Joseph L, et al. A cross-sectional study of the prevalence of intensity of infection with *Schistosoma japonicum* in 50 irrigated and rain-fed villages in Samar Province, the Philippines. *BMC Public Health* 2006;**6**:61.
20. Ross AG, Olveda RM, Chy D, Olveda DU, Li Y, Harn DA, et al. Can mass drug administration lead to the sustainable control schistosomiasis in the Philippines? *J Infect Dis* 2015;**211**:283–9. <http://dx.doi.org/10.1093/infdis/jiu416>
21. Olveda DU, Olveda RM, Lam AK, Chau TN, Li Y, Gisparil 2nd AD, Ross AG. Utility of diagnostic imaging in the diagnosis and management of schistosomiasis. *Clin Microbiol* 2014;**3**(2). <http://dx.doi.org/10.4172/2327-5073.1000142>. pii: 142.
22. Olveda R. Research update on neglected tropical diseases in the Philippines. Presented at the Regional Network for Asian Schistosomiasis and Other Zoonoses (RNAS+), Bogor, Indonesia, October 13–15, 2014. Unpublished results.
23. Ross AG, Li Y, Sleigh AC, Williams GM, McManus DP. Faecal egg aggregation in humans infected with *Schistosoma japonicum* in China. *Acta Trop* 1998;**70**:205–10.