



REVIEW

Open Access

Complicated intra-abdominal infections in a worldwide context: an observational prospective study (CIAOW Study)

Massimo Sartelli^{1*}, Fausto Catena², Luca Ansaloni³, Ernest Moore⁴, Mark Malangoni⁵, George Velmahos⁶, Raul Coimbra⁷, Kaoru Koike⁸, Ari Leppaniemi⁹, Walter Biffi⁴, Zsolt Balogh¹⁰, Cino Bendinelli¹⁰, Sanjay Gupta¹¹, Yoram Kluger¹², Ferdinando Agresta¹³, Salomone Di Saverio¹⁴, Gregorio Tugnoli¹⁴, Elio Jovine¹⁵, Carlos Ordonez¹⁶, Carlos Augusto Gomes¹⁷, Gerson Alves Pereira Junior¹⁸, Kuo-Ching Yuan¹⁹, Miklosh Bala²⁰, Miroslav P Peev⁶, Yunfeng Cui²¹, Sanjay Marwah²², Sanoop Zachariah²³, Boris Sakakushev^{24,25}, Victor Kong²⁶, Adamu Ahmed²⁷, Ashraf Abbas²⁸, Ricardo Alessandro Teixeira Gonsaga²⁹, Gianluca Guercioni³⁰, Nereo Vettoretto³¹, Elia Poiasina³, Offir Ben-Ishay¹², Rafael Díaz-Nieto³², Damien Massalou³³, Matej Skrovina³⁴, Ihor Gerych³⁵, Goran Augustin³⁶, Jakub Kenig³⁷, Vladimir Khokha³⁸, Cristian Tranà³⁹, Kenneth Yuh Yen Kok⁴⁰, Alain Chichom Mefire⁴¹, Jae Gil Lee⁴², Suk-Kyung Hong⁴³, Helmut Alfredo Segovia Lohse⁴⁴, Wagih Ghnam⁴⁵, Alfredo Verni⁴⁶, Varut Lohsiriwat⁴⁷, Boonying Siribumrungwong⁴⁸, Alberto Tavares⁴⁹, Gianluca Baiocchi⁵⁰, Koray Das⁵¹, Julien Jarry⁵², Maurice Zida⁵³, Norio Sato⁸, Kiyoshi Murata⁵⁴, Tomohisa Shoko⁵⁵, Takayuki Irahara⁵⁶, Ahmed O Hamedelneel⁵⁷, Noel Naidoo⁵⁸, Abdul Rashid Kayode Adesunkanmi⁵⁹, Yoshiro Kobe⁶⁰, AK Attri¹¹, Rajeev Sharma¹¹, Federico Coccolini³, Tamer El Zalabany⁶¹, Khalid Al Khalifa⁶¹, Juan Sanjuan¹⁶, Rita Barnabé¹⁵ and Wataru Ishii⁶²

Abstract

Despite advances in diagnosis, surgery, and antimicrobial therapy, mortality rates associated with complicated intra-abdominal infections remain exceedingly high. The World Society of Emergency Surgery (WSES) has designed the CIAOW study in order to describe the clinical, microbiological, and management-related profiles of both community- and healthcare-acquired complicated intra-abdominal infections in a worldwide context. The CIAOW study (Complicated Intra-Abdominal infection Observational Worldwide Study) is a multicenter observational study currently underway in 57 medical institutions worldwide. The study includes patients undergoing surgery or interventional drainage to address complicated intra-abdominal infections. This preliminary report includes all data from almost the first two months of the six-month study period. Patients who met inclusion criteria with either community-acquired or healthcare-associated complicated intra-abdominal infections (IAIs) were included in the study. 702 patients with a mean age of 49.2 years (range 18–98) were enrolled in the study. 272 patients (38.7%) were women and 430 (62.3%) were men. Among these patients, 615 (87.6%) were affected by community-acquired IAIs while the remaining 87 (12.4%) suffered from healthcare-associated infections. Generalized peritonitis was observed in 304 patients (43.3%), whereas localized peritonitis or abscesses was registered in 398 (57.7%) patients. The overall mortality rate was 10.1% (71/702). The final results of the CIAOW Study will be published following the conclusion of the study period in March 2013.

* Correspondence: m.sartelli@virgilio.it

¹Department of Surgery, Macerata Hospital, Macerata, Italy

Full list of author information is available at the end of the article

Introduction

Intra-abdominal infections (IAIs) include a wide spectrum of pathological conditions, ranging from uncomplicated appendicitis to fecal peritonitis [1].

From a clinical perspective, IAIs are classified in two major categories: complicated and uncomplicated.

In uncomplicated IAIs, the infectious process only involves a single organ and does not spread to the peritoneum. Patients with such infections can be managed with either surgical resection or antibiotics. When the focus of infection is treated effectively by surgical excision, 24-hour perioperative prophylaxis is typically sufficient. Patients with less severe intra-abdominal infections, including acute diverticulitis and certain forms of acute appendicitis, may be treated non-operatively.

In complicated IAIs, the infectious process extends beyond a singularly affected organ, and causes either localized peritonitis or diffuse peritonitis. The treatment of patients with complicated intra-abdominal infections involves both source control and antibiotic therapy.

Intra-abdominal infections are further classified in two groups: community-acquired intra-abdominal infections (CA-IAIs) and healthcare-associated intra-abdominal infections (HA-IAIs). CA-IAIs are acquired directly in the community while HA-IAIs develop in hospitalized patients or residents of long-term healthcare facilities. HA-IAIs are associated with higher rates of mortality due to the patients' poorer underlying health and an increased likelihood of infection by multi-drug resistant microorganisms.

Source control encompasses all measures undertaken to eliminate the source of infection and to control ongoing contamination.

The most common source of infection in community-acquired intra-abdominal infections is the appendix, followed by the colon, and then the stomach. Dehiscence complicates 5-10% of intra-abdominal bowel anastomoses and is associated with high rates of mortality [2].

Ultrasound- and CT-guided percutaneous drainage of abdominal and extra-peritoneal abscesses have proven to be safe and effective in select patients [3-10].

Surgery is the most important therapeutic recourse for controlling intra-abdominal infections.

Generally, the choice of the procedure depends on the anatomical source of infection, on the degree of peritoneal inflammation, on the generalized septic response and on the patient's general conditions.

Patients suffering from severe peritonitis are prone to persisting intra-abdominal sepsis, even when the source of infection has been neutralized. Timely re-laparotomy is the only possible known surgical recourse, capable to significantly improve patient outcome in these cases.

In the event of secondary peritonitis, the decision and timing of re-laparotomy is largely subjective and is often based on a surgeon's professional experience. Factors

indicative of progressive or persistent organ failure during early postoperative follow-up analysis are the strongest indicators of ongoing infection and suggest positive findings upon re-laparotomy [11-13].

Three methods of localized, mechanical management of abdominal sepsis following the initial laparotomy, which was performed for purposes of source control, are currently debated within the medical community: open-abdomen, planned re-laparotomy and on-demand re-laparotomy

Antimicrobial therapy plays an integral role in the management of intra-abdominal infections, especially in critically ill patients requiring immediate empiric antibiotic therapy.

Empiric antibiotic therapy accounts for the most frequently isolated microorganisms as well as any local trends of antibiotic resistance.

The major pathogens involved in community-acquired intra-abdominal infections are Enterobacteriaceae and anaerobic microbes (especially *B. fragilis*).

An antimicrobial-based approach to treating intra-abdominal infections involves a delicate balance between the optimization of empirical therapy, which has been shown to improve clinical outcomes, and the reduction of excessive antimicrobial use, which has been proven to increase the rate of emergence of antimicrobial-resistant strains.

The threat of antimicrobial resistance is one of the major challenges associated with the antimicrobial management of complicated intra-abdominal infections.

The recent and rapid spread of serine carbapenemases in *Klebsiella pneumoniae* (KPC) has become an important concern when administering antimicrobial therapy in hospitals worldwide [14].

The growing emergence of multidrug-resistant bacteria and the limited availability of new antibiotics to counteract them has brought about an impending crisis with alarming implications (especially regarding gram-negative microorganisms).

Methods

Aim

The purpose of the study is to describe the clinical, microbiological, and treatment profiles of both community-associated and healthcare-acquired complicated intra-abdominal infections (IAIs) in a worldwide context.

Patients older than 18 years with both community-acquired and healthcare-associated intra-abdominal infections will be included in the database.

In Europe, the CIAO Study has recently ended, concluding a six-month, multicenter observational study across twenty European countries. The study's findings have recently been published [15].

Given the promising results of the CIAO Study, the World Society of Emergency Surgery (WSES) has designed a prospective observational study investigating the management of complicated intra-abdominal infections in a worldwide context.

Study population

The CIAOW study (Complicated Intra-Abdominal Infection Observational Worldwide Study) is a multicenter observational study currently underway in 57 medical institutions worldwide. The study includes patients undergoing surgery or interventional drainage to address complicated IAIs.

Medical institutions from each continent participate in the study. The geographical distribution of the participating centers is represented in Figure 1.

Study design

The study does not attempt to change or modify the laboratory or clinical practices of the participating physicians, and neither informed consent nor formal approval by an Ethics Committee has been required.

The study meets the standards outlined in the Declaration of Helsinki and Good Epidemiological Practices.

The study is monitored by the coordination center, which investigates and verifies missing or unclear data submitted to the central database. It is performed under the direct supervision of the board of directors of WSES.

Data collection

In each center, the coordinator collects and compiles data in an online case report system. These data include the following: (i) patient and disease characteristics, i.e., demographic data, type of infection (community- or healthcare-acquired), severity criteria, previous curative antibiotic therapy administered in the 7 days preceding surgery; (ii) origin of infection and surgical procedures performed; and (iii) microbiological data, i.e., identification of bacteria and microbial pathogens within the peritoneal fluid, the presence of yeasts (if applicable), and the antibiotic susceptibilities of bacterial isolates.

The primary endpoints include the following:

- Clinical profiles of intra-abdominal infections
- Epidemiological profiles (epidemiology of the microorganisms isolated from intra-abdominal samples and these organisms' resistance to antibiotics)
- Management profiles

Statistical analysis

At the end of the six-month study period statistical comparisons will be performed using the Student's t-test, χ^2 analysis, or the Kruskal-Wallis/Wilcoxon tests, as

dictated by the natural parameters of the data. Statistical significance will be defined as a P-value less than 0.05 ($P < 0.05$). Multivariate analysis will be carried out by means of stepwise logistic regressions in order to assess the predictive factors of mortality during hospitalization. Adjusted odds ratios (OR) and their 95% confidence intervals (CI) will also be included.

Inclusion criteria

- Patients older than 18 years
- Community- and healthcare-acquired complicated intra-abdominal infections

Exclusion criteria

- Age under 18 years old
- Pancreatitis
- Primary peritonitis.

Preliminary results

Patients

This preliminary report includes all data from the first two months of the six-month study period.

702 patients with a mean age of 49.2 years (range 18–98) were enrolled in the study. 272 patients (38.7%) were women and 430 (62.3%) were men. Among these patients, 615 (87.6%) were affected by community-acquired IAIs while the remaining 87 (12.4%) suffered from healthcare-associated infections.


304 patients (43.3%) were affected by generalized peritonitis while 398 (57.7%) suffered from localized peritonitis or abscesses.

112 patients (15.9%) were admitted in critical condition (severe sepsis, septic shock).

Source control

The various sources of infection are outlined in Table 1. The most frequent source of infection was acute appendicitis. 243 cases were attributable to this condition.

The most frequently performed procedure employed to address complicated appendicitis was the open appendectomy. 136 patients (55.9%) admitted for complicated appendicitis underwent open appendectomies: 95 patients (69.8%) for localized infection or abscesses and 41 patients (31.2%) for generalized peritonitis. A laparoscopic appendectomy was performed on 93 patients (39.4%) presenting with complicated acute appendicitis, 82 (88.2%) and 11 (11.8%) of whom underwent the procedure for localized peritonitis/abscesses and generalized peritonitis, respectively. Open colonic resection was performed on 1 patient to address complicated appendicitis. In the other cases of complicated appendicitis, conservative treatment (percutaneous drainage, surgical drainage,



and non-operative treatment) was performed. 7 (3%) patients underwent percutaneous drainage to address appendicular abscesses.

For patients with complicated acute cholecystitis (104 cases), the most frequently performed procedure to address cholecystitis was the open cholecystectomy. 53 cholecystitis patients (51%) underwent this procedure. A laparoscopic cholecystectomy was performed on 27 patients (26%). In the remaining cases, conservative treatment methods (percutaneous drainage, non-operative treatment) were alternatively employed.

Among the patients with complicated diverticulitis (40) the Hartmann resection was the most frequently performed procedure. 12 patients (30%) underwent a Hartmann resection. All these resections were open procedures. 8 of these patients underwent a Hartmann resection for generalized peritonitis, while the remaining 4 underwent the same procedure for localized peritonitis or abscesses.

Colo-rectal resection was performed in 11 cases (27.5%) (4 with and 7 without stoma protection).

The other patients received conservative treatment (percutaneous drainage, non-operative treatment, surgical drainage and stoma). Only two (5%) underwent laparoscopic lavage and drainage.

Of the 100 patients with gastro-duodenal perforations, the most frequent surgical procedure was gastro-duodenal suture. It was performed in 91 patients (91%): 85 patients underwent open gastro-duodenal suture and 6 patients underwent laparoscopic gastro-duodenal suture. Four (4%) patients underwent gastro-duodenal resection. The

remaining patients (5%) received conservative treatment (non-operative treatment, surgical drainage).

Among the 53 patients with small bowel perforations, 35 underwent open small bowel resection (79.5%) and one (4.5%) underwent laparoscopic small bowel resection. Fourteen patients were treated by stoma. Two patients were treated by open drainage

Among the 38 patients with colonic non-diverticular perforation, 15 patients (66%) underwent open Hartmann resection, 1 patient (2.6%) underwent laparoscopic Hartmann resection, 9 (25%) underwent open resection with anastomosis and without stoma protection, and 4 underwent open resection with stoma protection (10.5%).

Microbiology

Intraperitoneal specimens were collected from 415 (59.1%) patients.

Intraperitoneal specimens were isolated from 336 of the 615 patients with community-acquired intra-abdominal infections (54.6%). Among the remaining 87 patients with healthcare-associated intra-abdominal infections, intraperitoneal specimens were collected from 79 patients (90.9%).

The major pathogens involved in intra-abdominal infections were found to be Enterobacteriaceae.

The aerobic bacteria identified in samples of peritoneal fluid are reported in Table 2.

According to CIAOW Study data, ESBL producers were the most commonly identified drug-resistant microorganism involved in IAIs.

Table 1 Source of infection

Source of infection	Patients N 702 (100%)
Appendicitis	243 (34.6%)
Cholecystitis	104 (14.8%)
Post-operative	53 (7.5%)
Colonic non diverticular perforation	38 (5.4%)
Gastroduodenal perforations	100 (14.2%)
Diverticulitis	40 (5.7%)
Small bowel perforation	53 (7.5%)
Others	52 (7.4%)
PID	8 (1.1%)
Post traumatic perforation	11 (1.6%)

1 identified isolate of *Klebsiella pneumoniae* proved resistant to Carbapenems.

Among the identified aerobic gram-negative isolates, there were 25 isolates of *Pseudomonas aeruginosa*, comprising 5.5% of all identified aerobic bacteria isolates.

Among the identified aerobic gram-positive bacteria, Enterococci (*E. faecalis* and *E. faecium*) were the most prevalent, representing 10.5% of all aerobic isolates, 3 glycopeptide-resistant Enterococci were identified; 2 were glycopeptide-resistant *Enterococcus faecalis* isolates and 1 was glycopeptide-resistant *Enterococcus faecium* isolates.

Tests for anaerobes were conducted for 168 patients.

52 anaerobes were observed. The most frequently identified anaerobic pathogen was *Bacteroides*. 39 *Bacteroides* isolates were observed during the course of the study.

Additionally, 36 *Candida* isolates were collectively identified. 30 were *Candida albicans* and 6 were non-*albicans Candida*.

Outcome

The overall mortality rate was 10.1% (71/702).

68 patients (9.7%) were admitted to the intensive care unit in the early recovery phase immediately following surgery.

90 patients (12.8%) ultimately required additional surgeries; 54% of these underwent relaparotomies “on-demand”, 28.9% underwent open abdomen procedures.

According to univariate statistical analysis, a critical clinical condition (severe sepsis and septic shock) upon hospital admission was the most significant risk factor for death; indeed, the rate of patient mortality was 36.6% (41/112) among critically ill patients (patients presenting with septic shock and severe sepsis upon admission), but the mortality rate was only 5.1% (30/590) for clinically stable patients ($p < 0.0001$).

For patients with generalized peritonitis, the mortality rate was 18% (55/304) while patients with localized peritonitis or abscesses demonstrated a mortality rate of only 4% (16/398) ($p < 0,001$).

The immediate post-operative clinical course was a significant parameter for predicting mortality: the rate of patient mortality was 54.9% (51/93) among critically ill patients (patients presenting with septic shock and severe sepsis upon the immediate post-operative course), but the mortality rate was only 3.3% (20/609) for clinically stable patients ($p < 0.0001$).

Table 2 Aerobic bacteria identified in peritoneal fluid

Total	455 (100%)
Aerobic gram-negative bacteria	352
<i>Escherichia coli</i>	226(49.7%)
(<i>Escherichia coli</i> resistant to third generation cephalosporins)	37 (8.1%)
<i>Klebsiella pneumoniae</i>	53 (11.6%)
(<i>Klebsiella pneumoniae</i> resistant to third generation cephalosporins)	13 (2.9%)
<i>Klebsiella oxytoca</i>	3 (0.7%)
<i>Enterobacter</i>	10 (2.2%)
<i>Proteus</i>	13 (2.9%)
<i>Pseudomonas</i>	25 (5.5%)
Others	22 (4.8%)
Aerobic gram-positive bacteria	103
<i>Enterococcus faecalis</i>	27 (5.9%)
<i>Enterococcus faecium</i>	21 (4.6%)
<i>Staphylococcus Aureus</i>	11 (2.4%)
<i>Streptococcus spp.</i>	29 (6.5%)
Others	15 (3.3%)

Preliminary statistical analyses were performed using MedCalc[®] statistical software.

Conclusion

Complicated intra-abdominal infections remain an important cause of morbidity with poor clinical prognoses.

The purpose of the CIAOW Study is to describe the epidemiological, clinical, microbiological, and treatment profiles of both community-acquired and healthcare-acquired complicated intra-abdominal infections (IAIs) based on the data collected over a six-month period (October 2012 to March 2013) from 56 medical institutions Worldwide.

The final results of the CIAOW Study will be published following the conclusion of the study period in March 2013.

Competing interests

The authors declare that they have no competing interests.

Authors' contributions

MS designed the study and wrote the manuscript. All authors read and approved the final manuscript.

Author details

¹Department of Surgery, Macerata Hospital, Macerata, Italy. ²Emergency Surgery, Maggiore Parma Hospital, Parma, Italy. ³Department of General Surgery, Ospedali Riuniti, Bergamo, Italy. ⁴Department of Surgery, Denver Health Medical Center, Denver, CO, USA. ⁵American Board of Surgery, Philadelphia, PA, USA. ⁶Harvard Medical School, Division of Trauma, Emergency Surgery and Surgical Critical Care Massachusetts General Hospital, Boston, MA, USA. ⁷Department of Surgery, UC San Diego Health System, San Diego, CA, USA. ⁸Department of Primary Care & Emergency Medicine, Kyoto University Graduate School of Medicine, Kyoto, Japan. ⁹Department of Abdominal Surgery, University Hospital Meilahti, Helsinki, Finland. ¹⁰Department of Surgery, University of Newcastle, Newcastle, NSW, Australia. ¹¹Department of Surgery, Govt Medical College and Hospital, Chandigarh, India. ¹²Department of General Surgery, Rambam Health Care Campus, Haifa, Israel. ¹³Department of Surgery, Adria Hospital, Adria, Italy. ¹⁴Trauma Surgery Unit, Maggiore Hospital, Bologna, Italy. ¹⁵Department of Surgery, Maggiore Hospital, Bologna, Italy. ¹⁶Department of Surgery, Universidad del Valle, Fundacion Valle del Lili, Cali, Colombia. ¹⁷Department of Surgery, Monte Sinai Hospital, Juiz de Fora, Brazil. ¹⁸Emergency Unit, Department of Surgery, Ribeirão Preto, Brazil. ¹⁹Department of Surgery, Chang Gung Memorial Hospital, Taoyuan, Taiwan. ²⁰Department of General Surgery, Hadassah Medical Center, Jerusalem, Israel. ²¹Department of Surgery, Tianjin Nankai Hospital, Nankai Clinical School of Medicine, Tianjin Medical University, Tianjin, China. ²²Department of Surgery, Pt BDS Post-graduate Institute of Medical Sciences, Rohtak, India. ²³Department of Surgery, MOSC medical college, Cochin, India. ²⁴First Clinic of General Surgery, University Hospital /UMBAL/ St George Plovdiv, Plovdiv, Bulgaria. ²⁵General Surgery Clinic, Medical University/University Hospital St.George, Plovdiv, Bulgaria. ²⁶Department of Surgery, Edendale Hospital, Pietermaritzburg, Republic of South Africa. ²⁷Department of Surgery, Ahmadu Bello University Teaching Hospital Zaria, Kaduna, Nigeria. ²⁸Department of Surgery, Mansoura University Hospital, Mansoura, Egypt. ²⁹Department of Surgery, Faculdades Integradas Padre Albino, Catanduva, Brazil. ³⁰Department of Surgery, Mazzoni Hospital, Ascoli Piceno, Italy. ³¹Department of Surgery, Mellini Hospital, Chiari (BS), Italy. ³²Department of General and Digestive Surgery, Virgen de la Victoria, University Hospital, Malaga, Spain. ³³Department of General Surgery and Surgical Oncology, Université de Nice Sophia-Antipolis, University Hospital of Nice, Nice, France. ³⁴Department of Surgery, Hospital and Oncological Centre, Novy Jicin, Czech Republic. ³⁵Department of General Surgery, Lviv Emergency Hospital, Lviv, Ukraine. ³⁶Department of Surgery, University Hospital Center Zagreb, Zagreb, Croatia. ³⁷3rd Department of General Surgeon Jagiellonian University, Narutowicz Hospital, Krakow, Poland.

³⁸Department of Surgery, Mozyr City Hospital, Mozyr, Belarus. ³⁹Department of Surgery, Ancona University, Ancona, Italy. ⁴⁰Department of Surgery, Ripas Hospital, Bandar Seri Begawan, Brunei. ⁴¹Clinical Sciences, Regional Hospitals Limbe and Buea, Limbe, Cameroon. ⁴²Department of Surgery, Severance Hospital, Yonsei University College of Medicine, Seoul, Republic of Korea. ⁴³Division of Trauma and Surgical Critical Care, Department of Surgery, University of Ulsan, Seoul, Republic of Korea. ⁴⁴II Cátedra de Clínica Quirúrgica, Hospital de Clínicas, Asunción, Paraguay. ⁴⁵Department of Surgery, Khamis Mushayt General Hospital, Khamis Mushayt, Saudi Arabia. ⁴⁶Department of Surgery, Cutral Co Clinic, Neuquen, Argentina. ⁴⁷Department of Surgery, Faculty of Medicine Siriraj Hospital, Bangkok, Thailand. ⁴⁸Department of Surgery, Thammasat University Hospital, Pathumthani, Thailand. ⁴⁹Department of Surgery, Hospital Regional de Alta Especialidad del Bajío, Leon, Mexico. ⁵⁰Department of Clinical and Experimental Sciences, Brescia University, Brescia, Italy. ⁵¹General Surgery, Adana Numune Training and Research Hospital, Adana, Turkey. ⁵²Visceral Surgery, Military Hospital Desgenettes, Lyon, France. ⁵³Visceral Surgery, Teaching Hospital Yalgado Ouedraogo, Ouagadougou, Burkina Faso. ⁵⁴Department of Acute and Critical care medicine, Tokyo Medical and Dental University, Tokyo, Japan. ⁵⁵The Shock Trauma and Emergency Medical Center, Matsudo City Hospital, Chiba, Japan. ⁵⁶Emergency and Critical Care Center of Nippon Medical School, Tama-Nagayama Hospital, Tokyo, Japan. ⁵⁷Department of Surgery, Our Lady of Lourdes Hospital, Drogheda, Ireland. ⁵⁸Department of Surgery, Port Shepstone Hospital, Port Shepstone, South Africa. ⁵⁹Department of Surgery, College of Health Sciences, Obafemi Awolowo University Hospital, Ile-Ife, Nigeria. ⁶⁰Department of Emergency and Critical Care Medicine, Chiba University Hospital, Chiba, Japan. ⁶¹Department of Surgery, Bahrain Defence Force Hospital, Manama, Bahrain. ⁶²Department of Emergency Medicine, Kyoto Second Red Cross Hospital, Kyoto, Japan.

Received: 28 December 2012 Accepted: 2 January 2013

Published: 3 January 2013

References

- Sartelli M, Viale P, Koike K, Pea F, Tumietto F, van Goor H, Guercioni G, Nespoli A, Tranà C, Catena F, Ansaloni L, Leppaniemi A, Biffi W, Moore FA, Poggetti R, Pinna AD, Moore EE, WSES consensus conference: **Guidelines for first-line management of intra-abdominal infections.** *World J Emerg Surg* 2011, **6**:2.
- Sartelli M: **A focus on intra-abdominal infections.** *World J Emerg Surg*. 2010, **5**:9.
- Azzarello G, Lanteri R, Rapisarda C, Santangelo M, Racialbuto A, Minutolo V, Di Cataldo A, Licata A: **Ultrasound-guided percutaneous treatment of abdominal collections.** *Chir Ital* 2009, **61**:337-340.
- Gazelle GS, Mueller PR: **Abdominal abscess: Imaging and intervention.** *Radiol Clin North Am* 1994, **32**:913-932.
- VanSonnenberg E, Ferrucci JT, Mueller PR, Wittenberg J, Simeone JF: **Percutaneous drainage of abscesses and fluid collections: Technique, results, and applications.** *Radiology* 1982, **142**:1-10.
- Bouali K, Magotteaux P, Jadot A, Saive C, Lombard R, Weerts J, Dallemagne B, Jehaes C, Delforge M, Fontaine F: **Percutaneous catheter drainage of abdominal abscess after abdominal surgery: Results in 121 cases.** *J Belg Radiol* 1993, **76**:11-14.
- VanSonnenberg E, Wing VW, Casola G, Coons HG, Nakamoto SK, Mueller PR, Ferrucci JT Jr, Halasz NA, Simeone JF: **Temporizing effect of percutaneous drainage of complicated abscesses in critically ill patients.** *Am J Roentgenol* 1984, **142**:821-826.
- Bufalari A, Giustozzi G, Moggi L: **Postoperative intra-abdominal abscesses: Percutaneous versus surgical treatment.** *Acta Chir Belg* 1996, **96**(5):197-200.
- VanSonnenberg E, Mueller PR, Ferrucci JT Jr: **Percutaneous drainage of 250 abdominal abscesses and fluid collections. I. Results, failures, and complications.** *Radiology* 1984, **151**:337-341.
- Jaffe TA, Nelson RC, DeLong D, Paulson EK: **Practice Patterns in Percutaneous Image-guided Intra-abdominal Abscess Drainage: Survey of Academic and Private Practice Centres.** *Radiology* 2004, **233**:750-756.
- van Ruler O, Lamme B, Gouma DJ, Reitsma JB, Boermeester MA: **Variables associated with positive findings at relaparotomy in patients with secondary peritonitis.** *Crit Care Med* 2007, **35**(2):468-476.

12. Hutchins RR, Gunning MP, Lucas DN, Allen-Mersh TG, Soni NC: **Relaparotomy for suspected intraperitoneal sepsis after abdominal surgery.** *World J Surg* 2004, **28**(2):137–141.
13. Lamme B, Mahler CW, van Ruler O, Gouma DJ, Reitsma JB, Boermeester MA: **Clinical predictors of ongoing infection in secondary peritonitis: systematic review.** *World J Surg* 2006, **30**(12):2170–2181.
14. Hawser SP, Bouchillon SK, Lascols C, Hackel M, Hoban DJ, Badal RE, Woodford N, Livermore DM: **Susceptibility of *Klebsiella pneumoniae* isolates from intra-abdominal infections and molecular characterization of ertapenem-resistant isolates.** *Antimicrob Agents Chemother* 2011, **55**(8):3917–3921.
15. Sartelli M, Catena F, Ansaloni L, Leppaniemi A, Taviloglu K, Goor H, Viale P, Lazzareschi DV, Coccolini F, Corbella D, Werra C, Marrelli D, Colizza S, Scibè R, Alis H, Torer N, Navarro S, Sakakushev B, Massalou D, Augustin G, Catani M, Kauhanen S, Pletinckx P, Kenig J, Saverio S, Jovine E, Guercioni G, Skrovina M, Diaz-Nieto R, Ferrero A, et al: **Complicated intra-abdominal infections in Europe: a comprehensive review of the CIAO study.** *World J Emerg Surg* 2012, **7**(1):36.

doi:10.1186/1749-7922-8-1

Cite this article as: Sartelli et al.: Complicated intra-abdominal infections in a worldwide context: an observational prospective study (CIAOW Study). *World Journal of Emergency Surgery* 2013 **8**:1.

Submit your next manuscript to BioMed Central and take full advantage of:

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit

