

RESEARCH ARTICLE

Open Access

Population-level effects of the national diabetes prevention programme (FIN-D2D) on the body weight, the waist circumference, and the prevalence of obesity

Titta M Salopuro^{1,2*}, Timo Saaristo^{1,3}, Heikki Oksa¹, Hannu Puolijoki⁴, Mauno Vanhala^{5,6,7}, Tapani Ebeling⁸, Leo Niskanen⁹, Jaakko Tuomilehto^{2,10}, Matti Uusitupa^{6,11} and Markku Peltonen²

Abstract

Background: The implementation project of the national diabetes prevention programme in Finland, FIN-D2D, was carried out in primary health care in the area of five hospital districts during 2003-2007.

Methods: The population strategy of FIN-D2D was primarily aimed at increasing the awareness of type 2 diabetes and preventing obesity. To investigate the effects of this strategy, we studied the changes in the prevalence of obesity, overweight, and central obesity among a random independent sample of individuals aged 45-74 years in the FIN-D2D area; and assessed whether they differed from a sample of individuals in the control area, which consisted of four geographical areas not participating in FIN-D2D (FINRISK study). Data was obtained for 5850/ 6406 (in the beginning/ in the end) individuals. The duration of the observation period varied from three to five years.

Results: The mean body weight decreased from 78.7 to 78.1 kg ($p = 0.041$) in the FIN-D2D area, and from 78.7 to 78.0 kg ($p = \text{NS}$) in the control area. The prevalence of obesity ($\text{BMI} \geq 30 \text{ kg/m}^2$) decreased in the FIN-D2D area (26.5% vs. 24.4%, $p = 0.015$), and in the control area (28.4% vs. 25.2%, $p = 0.005$). The prevalence of morbid obesity ($\text{BMI} \geq 40 \text{ kg/m}^2$) remained unchanged in the FIN-D2D area, but increased in the control area (1.2% vs. 2.3%, $p = 0.007$). The mean waist circumference remained unchanged in the FIN-D2D area, but increased in the control area (92.8 vs. 94.0 cm, $p = 0.005$).

Conclusions: The prevalence of obesity may be decreasing among 45-74 year old Finns. We still need a longer time perspective and future studies to see whether this favourable trend can be sustained in Finland. The actions of this implementation project can at least partly explain the differences in the mean waist circumference and the prevalence of morbid obesity between the intervention and control areas.

Background

During the last decades, the prevalence of obesity has doubled in adults in the United States [1] and in Finland [2]. However, recent reports have shown that the obesity epidemic might be slowing down [3,4]. Obesity is strongly associated with a reduction in life expectancy, as well as with an increased risk of type 2 diabetes (T2DM), coronary heart disease, and certain cancers [5]. Furthermore, obesity increases the risk of musculoskeletal

disorders [6] and early retirement [7]. Obesity increases the risk of T2DM to a much greater extent than it increases the risk of other diseases, independently of age, race, and physical activity [8]. Furthermore, numerous large-scale studies have shown that overall mortality rises steadily as a function of increasing body weight [9], and even small increases in weight across a population can have a devastating impact on public health.

According to a large meta-analysis, more than a half of the Europeans will suffer from hyperglycemia and T2DM during their lifetime [10,11]. Thus, the economic burden is already big and will be huge in the forthcoming years

* Correspondence: titta.salopuro@thl.fi

¹Pirkanmaa Hospital District, Tampere, Finland

Full list of author information is available at the end of the article

[12]. The most efficient way to manage T2DM and its complications is to prevent the onset of diabetes and pre-diabetic states. As several intervention trials have demonstrated, prevention of T2DM and its complications is possible and efficient by modest changes in lifestyle, primarily in diet and physical activity [13-17]. Moreover, it has been shown that the achieved benefit from lifestyle interventions can be sustained for years after the active period [18-20]. A recent meta-analysis [21] has shown that lifestyle interventions for patients at high risk of diabetes, delivered by a variety of healthcare providers in routine clinical settings, are feasible but appear to be of limited clinical benefit one year after intervention. Although weight and waist circumference reductions were achieved, translation into routine practice had less effect on diabetes risk reduction [21].

To translate the scientific evidence regarding prevention of T2DM into daily practice and public health, the first large-scale national type 2 diabetes prevention programme was carried out in Finland, covering three concurrent strategies: the population strategy, the high-risk strategy, and the strategy of early diagnosis and treatment [22,23]. The baseline results of the implementation project, FIN-D2D, have shown that only 58% of men and 67% of women have normal glucose tolerance, and the disturbances of glucose metabolism are closely associated with obesity and central obesity [24]. FIN-D2D supported the population strategy that aimed to improve the health of the entire population through a healthy diet and regular physical activity [22]. Different manners, such as media communication, training, lifestyle counselling and an extensive network supporting these activities, were used to prevent obesity, metabolic syndrome and other risk factors of T2DM [25]. The project was carried out in five hospital districts in Finland in 2003-2007.

In this study based on two independent surveys with a follow-up of 3 to 5 years, we examined the effects of FIN-D2D on the prevalence of obesity, overweight, and central obesity at the population-level, among 45-74 years old Finnish men and women, and assessed whether these effects differ from the respective changes among Finnish population living in four geographical areas not participating in FIN-D2D. The results of this study reflect the achievements of all three concurrent strategies at the population-level, whereas the 1-year results of the high-risk strategy have been published elsewhere [26].


Methods

FIN-D2D was carried out in five hospital districts, covering altogether 1.5 million people (South Ostrobothnia, Central Finland, Pirkanmaa, Northern Ostrobothnia, and Northern Savo) during 2003-2007. As a part of the evaluation, population-based health examination surveys with independent random samples from the population

were conducted in the beginning and at the end of the project period. Individuals aged 45-74 years, stratified according to sex, 10-year age groups (45-54, 55-64, and 65-74 years) and five geographical areas, were selected from the National Population Register at the beginning of the project (year 2002 for Northern Ostrobothnia and Northern Savo, and year 2004 for South Ostrobothnia, Central Finland, and Pirkanmaa) and at the end of the project (year 2007 for all five areas). Data on anthropometric measurements were obtained for 3812 and 4156 individuals (at the beginning and at the end, respectively).

The FINRISK study [27,28] is a chronic disease risk factor survey of the middle-aged Finnish population that has been carried out every five years since 1972. The geographical FINRISK study areas which were not participating in FIN-D2D project were used as a control area in this study (Figure 1). In this study we used the data from FINRISK-2002 and FINRISK-2007 studies covering the following geographical areas in Finland: Northern Savo, Northern Karelia, Northern Ostrobothnia, Kainuu, Turku and Loimaa region, and the cities of Helsinki and Vantaa. Northern Savo and Northern Ostrobothnia belong to the FIN-D2D area (hospital districts of Northern Savo and Northern Ostrobothnia, respectively), thus the data from these areas were combined with the FIN-D2D data from the hospital districts of South Ostrobothnia, Central Finland, and Pirkanmaa, to constitute the FIN-D2D area [25]. Study participants aged 45-74 years were included in this study. Data on anthropometric measurements were obtained for 2038 and 2250 (years 2002 and 2007, respectively) individuals from the areas of Northern Karelia, Kainuu, Turku-Loimaa, and Helsinki-Vantaa [25]. This study population will be referred to as the control area in this study.

The primary strategy of the FIN-D2D was a high-risk strategy, the aim of which was to include prevention of diabetes and reduction of cardiovascular risk factor levels among high-risk individuals in daily routines in primary health care centers and occupational health care outpatient clinics [26]. The population strategy focused on raising awareness of diabetes and its risk factors in the overall population, and the early treatment strategy focused on care of diabetes in individuals who had screening-detected diabetes [26]. The different strategies have been described in detail elsewhere [23,26]. Subjects were invited by mail to a clinical examination. Together with the invitation, they also received a self-administered questionnaire on socioeconomic background, medical history, and health behaviour. They were asked to complete the questionnaire at home, and bring it with them to the health examination, which was carried out according to the WHO MONICA project protocol [29]. At the study site, height, weight and waist circumference were measured using a standardized



protocol. Height was measured to the nearest 0.1 cm. Body weight of the participants wearing usual light indoor clothing without shoes was measured with a 0.1 kg precision on a balanced beam scale (FINRISK, FIN-D2D 2004) or on an electronic scale (FIN-D2D 2007).

BMI was calculated as weight (kg) divided by height squared (m²). Overweight and obesity were defined as BMI 25-29.9 kg/m² and ≥30 kg/m², respectively. Obesity was further divided into class 1 (BMI 30-34.9), class 2 (BMI 35-39.9), and class 3 (BMI ≥40) obesity [25]. Central obesity was defined according to the IDF criteria [30]: waist circumference ≥94 cm in men, and ≥80 cm in women, and according to the criteria used by WHO [31]: waist circumference ≥102 cm in men, and ≥88 cm in women.

Statistical analysis was done by the SPSS program version 17 (SPSS, Chigago, IL, USA). The univariate ANOVA was used to test the difference in the continuous variables and in the group-specific changes in them. Logistic regression analysis was performed to evaluate the differences in various prevalences between the study areas. Interaction term 'study area × time point' was used to assess differences in changes between the areas. Adjustment for age and gender was done, when appropriate. In the stratified analysis, genders and three age groups (45-54, 55-64, and 65-74 years) were analysed separately. Continuous variables are presented as mean (standard deviation; SD) and categorical variables as percentages.

The study protocols for FIN-D2D population survey study as well as the FINRISK study were approved by the Ethical Committee of the Hospital District of Helsinki and Uusimaa. All the surveys were conducted according to the ethical rules of the National Public Health Institute, and the investigations were performed in accordance with the Declaration of Helsinki. All participants gave their written informed consent prior to participation in the study.

Results

The overall response rate was 60% and 63% (beginning and end, respectively) in the FIN-D2D area, and 49% and 58% (beginning and end, respectively) in the control area (Table 1). The mean age of the study participants was higher at the end than at the beginning in both study populations, due to the fact that the oldest age group (65-74 years) was not included in the following FINRISK-2002 study areas: Turku and Loimaa region, Northern Savo, Northern Ostrobothnia and the Kainuu area. No differences in age were seen between the FIN-D2D and control areas at either time point (Table 1).

Changes in body weight and BMI

The mean body weight decreased significantly only in men in the FIN-D2D area (from 85.6 ± 14.2 to 84.4 ± 14.3 kg, *p* = 0.030) (Table 1). When stratified into age groups, no significant changes were seen, although

Table 1 Characteristics of the study participants according to sex, study area and timepoint

	FIN-D2D area		Control area	
	Beginning	End	Beginning	End
Men				
Number of subjects, invited	3175	3329	2060	1921
Participated (%)	1789 (56.3)	1982 (59.5)	984 (47.8)	1066 (55.5)
Age, years	59.1 (8.1)	60.1 (8.5)*	59.3 (7.8)	60.2 (8.2)*
Weight, kg	85.6 (14.2)	84.4 (14.3)*	85.1 (13.2)	84.5 (14.4)
BMI, kg/m ²	27.8 (4.1)	27.4 (4.2)*	27.9 (3.9)	27.6 (4.3)
Waist, cm	99.2 (11.6)	99.2 (11.8)	98.5 (11.1)	99.1 (12.1)
Women				
Number of subjects, invited	3173	3319	2067	1931
Participated (%)	2023 (63.8)	2174 (65.5)	1054 (51.0)	1184 (61.3)
Age, years	58.5 (8.2)	59.3 (8.3)*	58.3 (7.8)	59.7 (8.4)*
Weight, kg	72.6 (13.7)	72.3 (14.2)	72.7 (13.7)	72.1 (14.2)
BMI, kg/m ²	27.7 (5.1)	27.6 (5.3)	28.0 (5.0)	27.6 (5.4)*
Waist, cm	89.2 (13.2)	89.7 (13.3)	87.5 (12.6)	89.4 (13.5)*
Total				
Number of subjects, invited	6348	6648	4127	3852
Participated (%)	3812 (60.1)	4156 (62.5)	2038 (49.4)	2250 (58.4)
Age, years	58.8 (8.1)	59.7 (8.4)*	58.8 (7.8)	60.0 (8.3)*
Weight, kg	78.7 (15.4)	78.1 (15.5)*	78.7 (14.8)	78.0 (15.6)
BMI, kg/m ²	27.8 (4.7)	27.5 (4.8)*	27.9 (4.5)	27.6 (4.9)*
Waist, cm	93.9 (13.4)	94.3 (13.5)	92.8 (13.1)	94.0 (13.7)*

Data are presented as means (standard deviation) except where noted otherwise.

* $p < 0.05$ (adjusted for age) for the change from the beginning to the end of the study.

similar decreasing trend was observed in nearly all age groups in both genders. Among men, weight seemed to decrease with increasing age. The mean BMI decreased significantly only in men in the FIN-D2D area and women in the control area (Table 1). When stratified into age groups, BMI showed a decreasing trend in both genders, areas, and all age groups across the studies, but most so among the youngest men in the FIN-D2D area (27.6 ± 4.2 vs. 27.0 ± 4.2 kg/m², $p = 0.015$). Among women, BMI increased with increasing age.

Changes in the prevalence of obesity

The prevalence of obesity was significantly lower in the 2007 survey among men in both study areas (Table 2). Respectively, the prevalence of normal weight was higher in the 2007 survey, especially among the youngest age group in men. The prevalence of normal weight was higher and the prevalence of overweight was lower in the 2007 survey among women in the control area (Table 3), whereas no changes among women in the FIN-D2D area were observed. The prevalence of obesity increased with increasing age among women (Table 3). Altogether, normal weight was more prevalent among women than among men (34.2% vs. 26.6%), whereas overweight was more prevalent among men than among women (49.8% vs. 37.8%) (Figure 2).

To examine these findings further, obesity was categorized into subclasses 1, 2 and 3. The prevalence of class 1 obesity decreased in both areas, especially among men in the youngest age group (Table 2). Class 3 (morbid) obesity increased in the control area (from 1.2% to 2.3%, OR 1.97, 95% CI 1.21-3.22, $p = 0.007$) and remained unchanged in the FIN-D2D area (from 2.0% to 1.9%, $p = \text{NS}$) (Figure 2). The interaction for study area \times time point was significant for the genders combined ($p = 0.020$), and for the women in the age group of 55-64 years ($p = 0.044$). The prevalence of morbid obesity was highest among 55-64 year old women in the control area at the end of the study (4.0%, 95% CI 2.0%-5.8%), and lowest among 65-74 year old men in the control area in the beginning of the study (0.4%, 95% CI -0.4%-1.1%). Altogether, class 2 and 3 obesity were more prevalent among women than among men (6.7% vs. 4.0%, and 2.5% vs. 1.2%, for classes 2 and 3, respectively) (Figure 2).

Changes in waist circumference and prevalence of central obesity

Baseline waist circumference was larger in women in the FIN-D2D area (89.2 ± 13.2 cm) than in the control area (87.5 ± 12.6 cm) ($p = 0.001$). The mean waist circumference increased among women in the control area, but remained unchanged in the FIN-D2D area (Table 1).

Table 2 Age-adjusted prevalence (95% confidence interval, CI) of normal weight, overweight, obesity and central obesity in the beginning of the study and the changes in them in men according to the study area and age group

	FIN-D2D area				p*	Control area				p*	p**
	%	(95% CI)	Change	(95% CI)		%	(95% CI)	Change	(95% CI)		
45-54 years											
Number of subjects	587		581			288		302			
Normal weight	29.1	(25.4 - 32.8)	6.2	(0.8 - 11.5)	.027	28.1	(22.9 - 33.3)	5.0	(-2.5 - 12.4)	.185	.830
Overweight	47.5	(43.5 - 51.6)	-1.1	(-6.8 - 4.7)	.667	46.2	(40.4 - 52.0)	1.8	(-6.3 - 9.9)	.769	.561
Obesity	23.3	(19.9 - 26.8)	-5.1	(-9.8 - -0.4)	.043	25.7	(20.6 - 30.8)	-6.8	(-13.5 - -0.1)	.066	.720
Class 1 obesity	18.1	(14.9 - 21.2)	-4.8	(-9.0 - -0.6)	.033	22.9	(18.0 - 27.8)	-8.3	(-14.6 - -2.1)	.014	.474
Class 2 obesity	3.6	(2.1 - 5.1)	-0.1	(-2.3 - 2.0)	.934	2.1	(0.4 - 3.7)	0.9	(-1.7 - 3.4)	.453	.514
Class 3 obesity	1.7	(0.7 - 2.8)	-0.2	(-1.6 - 1.3)	.813	0.7	(-0.3 - 1.7)	0.6	(-1.0 - 2.3)	.457	.447
Central obesity IDF	60.5	(56.5 - 64.4)	-3.2	(-8.8 - 2.5)	.310	56.6	(50.8 - 62.4)	-1.0	(-9.0 - 7.1)	.899	.659
Central obesity WHO	32.0	(28.2 - 35.8)	-2.8	(-8.1 - 2.5)	.379	29.9	(24.5 - 35.2)	-3.4	(-10.7 - 3.9)	.431	.862
55-64 years											
Number of subjects	716		657			422		364			
Normal weight	22.2	(19.2 - 25.3)	4.6	(0.0 - 9.1)	.050	21.8	(17.8 - 25.8)	4.0	(-1.9 - 10.0)	.174	.930
Overweight	52.0	(48.3 - 55.6)	-0.5	(-5.8 - 4.8)	.780	50.0	(45.2 - 54.8)	-0.8	(-7.9 - 6.2)	.782	.960
Obesity	25.8	(22.6 - 29.1)	-4.1	(-8.6 - 0.4)	.099	28.2	(23.9 - 32.5)	-3.2	(-9.4 - 3.0)	.320	.814
Class 1 obesity	19.7	(16.8 - 22.6)	-3.9	(-7.9 - 0.2)	.086	22.5	(18.5 - 26.5)	-3.8	(-9.5 - 1.9)	.191	.959
Class 2 obesity	4.9	(3.3 - 6.5)	-0.9	(-3.1 - 1.3)	.363	5.0	(2.9 - 7.1)	-0.9	(-3.8 - 2.1)	.569	.920
Class 3 obesity	1.3	(0.4 - 2.1)	0.7	(-0.6 - 2.1)	.258	0.7	(-0.1 - 1.5)	1.5	(-0.2 - 3.1)	.074	.414
Central obesity IDF	70.6	(67.3 - 74.0)	-0.9	(-5.7 - 4.0)	.749	67.5	(63.0 - 72.0)	4.1	(-2.4 - 10.6)	.231	.251
Central obesity WHO	37.9	(34.3 - 41.5)	3.0	(-2.2 - 8.2)	.217	37.9	(33.3 - 42.6)	2.6	(-4.3 - 9.4)	.473	.863
65-74 years											
Number of subjects	486		744			274		400			
Normal weight	24.1	(20.3 - 27.9)	1.2	(-3.8 - 6.1)	.588	21.9	(17.0 - 26.8)	4.1	(-2.5 - 10.7)	.227	.489
Overweight	50.8	(46.4 - 55.3)	0.8	(-4.9 - 6.5)	.828	49.3	(43.3 - 55.2)	2.5	(-5.2 - 10.2)	.515	.726
Obesity	25.1	(21.2 - 29.0)	-2.0	(-6.9 - 2.9)	.422	28.8	(23.4 - 34.2)	-6.6	(-13.2 - 0.1)	.051	.291
Class 1 obesity	20.8	(17.2 - 24.4)	-2.6	(-7.1 - 1.9)	.276	23.7	(18.7 - 28.8)	-7.0	(-13.1 - -0.9)	.023	.273
Class 2 obesity	3.3	(1.7 - 4.9)	1.1	(-1.1 - 3.4)	.370	4.7	(2.2 - 7.3)	0.0	(-3.3 - 3.3)	.969	.529
Class 3 obesity	1.0	(0.1 - 1.9)	-0.5	(-1.5 - 0.5)	.297	0.4	(-0.4 - 1.1)	0.4	(-0.8 - 1.6)	.538	.296
Central obesity IDF	74.1	(70.2 - 78.0)	-2.1	(-7.2 - 3.0)	.385	69.3	(63.8 - 74.8)	-0.5	(-7.6 - 6.6)	.887	.690
Central obesity WHO	39.9	(35.5 - 44.3)	0.1	(-5.5 - 5.7)	.976	38.3	(32.5 - 44.1)	1.4	(-6.2 - 8.9)	.730	.787
Total											
Number of subjects	1789		1982			984		1066			
Normal weight	25.0	(23.0 - 27.0)	4.1	(1.2 - 6.9)	.005	23.7	(21.0 - 26.3)	4.6	(0.8 - 8.4)	.018	.791
Overweight	50.2	(47.9 - 52.5)	-0.3	(-3.5 - 2.9)	.866	48.7	(45.6 - 51.8)	1.0	(-3.3 - 5.3)	.654	.640
Obesity	24.8	(22.8 - 26.8)	-3.8	(-6.5 - -1.1)	.006	27.6	(24.8 - 30.4)	-5.6	(-9.3 - -1.8)	.004	.496
Class 1 obesity	19.5	(17.6 - 21.3)	-3.8	(-6.2 - -1.3)	.003	23.0	(20.3 - 25.6)	-6.3	(-9.7 - -2.8)	.000	.291
Class 2 obesity	4.0	(3.1 - 4.9)	0.0	(-1.3 - 1.2)	.949	4.1	(2.8 - 5.3)	-0.1	(-1.9 - 1.6)	.867	.994
Class 3 obesity	1.3	(0.8 - 1.9)	0.0	(-0.7 - 0.7)	.976	0.6	(0.1 - 1.1)	0.8	(0.0 - 1.7)	.063	.124
Central obesity IDF	68.2	(66.1 - 70.4)	-1.9	(-4.9 - 1.0)	.204	64.8	(61.8 - 67.8)	0.6	(-3.5 - 4.7)	.776	.325
Central obesity WHO	36.5	(34.3 - 38.8)	0.2	(-2.9 - 3.2)	.919	35.7	(32.7 - 38.7)	0.0	(-4.1 - 4.2)	.986	.982

*p-value for change, adjusted for age.

**Interaction for study area × timepoint, adjusted for age.

Here, the interaction for study area × time point for both genders combined was also significant ($p = 0.040$). Among men, no significant changes in waist circumference were seen in the control area or in the FIN-D2D area (Table 1). Since the changes in the waist circumference were rather modest, we saw no changes in the

prevalence of central obesity in either gender. Central obesity was more prevalent in women than in men (49.3% vs. 36.5%, and 74.5% to 66.8%, WHO and IDF criteria, respectively). The prevalence of central obesity increased with increasing age in both genders (Tables 2 and 3).

Table 3 Age-adjusted prevalence (95% confidence interval, CI) of normal weight, overweight, obesity and central obesity in the beginning of the study and the changes in them in women according to the study area and age group

	FIN-D2D area				p*	Control area				p*	p**
	%	(95% CI)	Change	(95% CI)		%	(95% CI)	Change	(95% CI)		
45-54 years											
Number of subjects	725		713			369		358			
Normal weight	41.5	(37.9 - 45.1)	2.1	(-3.0 - 7.2)	.489	40.1	(35.1 - 45.1)	7.4	(0.2 - 14.6)	.067	.270
Overweight	35.2	(31.7 - 38.7)	-1.4	(-6.3 - 3.5)	.662	37.4	(32.4 - 42.4)	-7.0	(-13.8 - -0.1)	.062	.212
Obesity	23.3	(20.2 - 26.4)	-0.7	(-5.1 - 3.6)	.750	22.5	(18.2 - 26.8)	-0.4	(-6.5 - 5.6)	.958	.928
Class 1 obesity	16.4	(13.7 - 19.1)	-1.0	(-4.8 - 2.8)	.615	13.3	(9.8 - 16.8)	0.7	(-4.3 - 5.7)	.764	.607
Class 2 obesity	5.4	(3.7 - 7.0)	-0.5	(-2.8 - 1.8)	.694	7.6	(4.9 - 10.3)	-2.8	(-6.3 - 0.7)	.114	.312
Class 3 obesity	1.5	(0.6 - 2.4)	0.7	(-0.7 - 2.1)	.314	1.6	(0.3 - 2.9)	1.7	(-0.5 - 4.0)	.100	.581
Central obesity IDF	68.0	(64.6 - 71.4)	0.7	(-4.1 - 5.5)	.709	62.8	(57.8 - 67.7)	2.8	(-4.2 - 9.8)	.348	.602
Central obesity WHO	42.0	(38.4 - 45.6)	2.4	(-2.7 - 7.6)	.304	34.5	(29.6 - 39.4)	4.1	(-2.9 - 11.2)	.210	.641
55-64 years											
Number of subjects	812		746			436		407			
Normal weight	30.5	(27.4 - 33.7)	3.8	(-0.9 - 8.4)	.121	28.9	(24.6 - 33.2)	6.2	(-0.1 - 12.5)	.052	.513
Overweight	39.9	(36.5 - 43.3)	-0.9	(-5.8 - 4.0)	.734	40.4	(35.7 - 45.0)	-4.0	(-10.6 - 2.6)	.240	.454
Obesity	29.6	(26.4 - 32.7)	-2.9	(-7.4 - 1.6)	.214	30.7	(26.4 - 35.1)	-2.2	(-8.4 - 4.0)	.463	.876
Class 1 obesity	18.6	(15.9 - 21.3)	-1.3	(-5.1 - 2.5)	.509	20.9	(17.0 - 24.7)	-3.4	(-8.8 - 1.9)	.199	.530
Class 2 obesity	7.8	(5.9 - 9.6)	-0.5	(-3.1 - 2.1)	.722	7.8	(5.3 - 10.3)	-0.7	(-4.2 - 2.9)	.715	.933
Class 3 obesity	3.2	(2.0 - 4.4)	-1.1	(-2.7 - 0.6)	.192	2.1	(0.7 - 3.4)	1.9	(-0.4 - 4.2)	.116	.044
Central obesity IDF	76.5	(73.6 - 79.4)	2.9	(-1.3 - 7.0)	.166	73.3	(69.2 - 77.5)	2.1	(-3.8 - 8.0)	.496	.757
Central obesity WHO	52.7	(49.3 - 56.2)	-1.8	(-6.8 - 3.1)	.478	48.0	(43.3 - 52.8)	2.3	(-4.5 - 9.1)	.503	.337
65-74 years											
Number of subjects	486		715			249		419			
Normal weight	26.3	(22.4 - 30.3)	0.7	(-4.5 - 5.8)	.773	22.5	(17.3 - 27.7)	6.6	(-0.3 - 13.6)	.066	.163
Overweight	40.9	(36.6 - 45.3)	-0.8	(-6.5 - 4.9)	.665	41.4	(35.2 - 47.5)	-2.7	(-10.4 - 5.0)	.484	.744
Obesity	32.7	(28.5 - 36.9)	0.2	(-5.3 - 5.6)	.857	36.1	(30.1 - 42.2)	-3.9	(-11.4 - 3.5)	.319	.341
Class 1 obesity	23.0	(19.3 - 26.8)	0.0	(-4.8 - 4.9)	.880	26.1	(20.6 - 31.6)	-2.7	(-9.5 - 4.0)	.455	.463
Class 2 obesity	6.8	(4.5 - 9.0)	0.1	(-2.8 - 3.0)	.970	8.8	(5.3 - 12.4)	-1.9	(-6.1 - 2.3)	.375	.471
Class 3 obesity	2.9	(1.4 - 4.4)	0.1	(-1.9 - 2.0)	.854	1.2	(-0.2 - 2.6)	0.7	(-1.3 - 2.7)	.502	.573
Central obesity IDF	81.1	(77.6 - 84.6)	-0.8	(-5.4 - 3.7)	.808	77.5	(72.3 - 82.7)	5.1	(-1.1 - 11.3)	.099	.162
Central obesity WHO	58.8	(54.5 - 63.2)	-2.7	(-8.4 - 3.0)	.492	54.2	(48.0 - 60.4)	2.6	(-5.2 - 10.4)	.492	.341
Total											
Number of subjects	2023		2174			1054		1184			
Normal weight	33.5	(31.4 - 35.5)	2.1	(-0.7 - 5.0)	.147	31.3	(28.5 - 34.1)	6.7	(2.8 - 10.6)	.001	.071
Overweight	38.5	(36.3 - 40.6)	-1.0	(-4.0 - 1.9)	.491	39.6	(36.6 - 42.5)	-4.6	(-8.6 - -0.5)	.026	.155
Obesity	28.1	(26.1 - 30.0)	-1.1	(-3.8 - 1.6)	.436	29.1	(26.4 - 31.9)	-2.1	(-5.9 - 1.6)	.261	.703
Class 1 obesity	18.9	(17.2 - 20.6)	-0.6	(-2.9 - 1.8)	.642	19.4	(17.1 - 21.8)	-1.8	(-5.0 - 1.5)	.289	.645
Class 2 obesity	6.7	(5.6 - 7.8)	-0.4	(-1.9 - 1.1)	.604	8.0	(6.3 - 9.6)	-1.7	(-3.9 - 0.4)	.110	.326
Class 3 obesity	2.5	(1.8 - 3.2)	-0.1	(-1.1 - 0.8)	.805	1.7	(0.9 - 2.5)	1.4	(0.1 - 2.6)	.038	.080
Central obesity IDF	74.6	(72.7 - 76.5)	1.1	(-1.5 - 3.7)	.398	70.6	(67.9 - 73.4)	3.2	(-0.5 - 6.8)	.090	.342
Central obesity WHO	50.3	(48.2 - 52.5)	-0.4	(-3.4 - 2.6)	.802	44.8	(41.8 - 47.8)	3.0	(-1.1 - 7.1)	.149	.157


*p-value for change, adjusted for age.

**Interaction for study area × timepoint, adjusted for age.

Discussion

The Finnish Programme for the Prevention of Type 2 Diabetes [22] is the first large-scale national programme ever launched to combat the diabetes epidemic at the population-level. The implementation project of this program, FIN-D2D, focused mainly on the high-risk

strategy, and the strategy of early diagnosis and treatment [25]. The FIN-D2D also supported the population strategy and it gained a lot of publicity in the whole country because its main aim was to restrain the obesity epidemic by means of nutritional interventions and increased physical activity. As shown in the present



study, this goal was achieved, since the mean body weight, mean BMI, and the prevalence of obesity decreased, whereas the prevalence of normal weight increased. This suggests that the prevalence of obesity may be decreasing among 45-74 year old Finns. To our knowledge, this is the first report to show a decline in the prevalence of obesity at the population-level, both in

the FIN-D2D and the control area. Interestingly, the waist circumference increased in the control area, whereas it remained unchanged in the FIN-D2D area. However, the increase in waist circumference in the control area was more modest than reported earlier for years 1987-2002 [32], and the prevalence of central obesity did not increase significantly in either area.

There is large geographic variation in the prevalence of obesity among European countries, with central, eastern, and southern regions showing higher rates than the western and northern (including Finland) regions [33]. Studies from Spain, Canada, Denmark, Portugal, and South Australia [34-38] have shown that the prevalence of obesity has increased between the mid-1990s and the beginning of the 2000s. In Finland, national FINRISK studies have shown that the increase in BMI has slowed down in men, and has been quite stable in women between 1997 and 2002, whereas a remarkable shift towards higher waist circumference has been observed [32]. BMI did not increase significantly between 1997 and 2002 in the age groups of 35-64 years, but it increased significantly in the youngest age group, 25-34 years, where also the increase in waist circumference was greatest [32]. Among US adults the increases in the prevalence of obesity previously observed does not appear to be continuing at the same rate during the past 10 years, particularly for women [3]. In Sweden, the increase in the prevalence of obesity has levelled off during 2000-2005 [39] compared to the 10% increase seen during the years 1996-2001 [40]. In our study, a significant decrease in the prevalence of obesity at the population-level was seen for the first time.

In addition to the changes seen in the FIN-D2D area, most changes in body weight and the prevalence of obesity were similar in the control area, especially among men. Although the five hospital districts of FIN-D2D formed a pilot area, the information of the importance of healthy lifestyles in the prevention of diabetes had also spread outside the FIN-D2D area, which was the aim of population strategy, as well. Moreover, the control area is part of the FINRISK area, where the measurements of cardiovascular risk factors have been repeated every five years since 1972 with impressive results [27,28], and they have also received feedback of these results. Therefore, the control area can be considered better as a mini-intervention area rather than a pure control area. The latest FINRISK-survey showed that many cardiovascular risk factors were proceeding into the right direction among 25-64 year old individuals in Finland [4,41]. However, the use of alcohol was still high [42], and this could be a reason why the waist circumference was increasing [32], and why the beneficial trend in blood pressure had now levelled off [4].

There were also substantial differences among the studied areas on how they succeeded in the project, which can largely be explained by their different ways of action

in the prevention of chronic diseases, and also their different population structures. It should also be noted that the prevalence of diabetes in the FIN-D2D area has already for a long time been higher than the mean prevalence of diabetes in Finland [25]. The areas were thus establishing this project from different starting points, but they also reached the goals differently. The FIN-D2D was a national health promoting programme, and not an actual scientific study. However, from the very beginning it was decided to analyse its influence by comparing changes in adiposity markers in the FIN-D2D area and the preselected control area.

In this study based on the two cross-sectional surveys with 3 to 5 years of follow-up it was surprising that men in the FIN-D2D area (especially the youngest age group) were able to reduce their body weight more than women. Usually women are more likely to participate in lifestyle change programmes than men [43]. In FIN-D2D there were several projects focusing on men especially, which may have encouraged men to join these projects, and to also succeed in them. The threshold for men to participate is usually higher than that for women, but their commitment nevertheless is usually better [43-45]. Also, the interventions aimed at women may indirectly affect also men due to the fact that men usually eat meals prepared by women [46]. Women in the control area showed beneficial changes in the prevalence of normal weight and overweight, but on the other hand, adverse changes regarding waist circumference and the prevalence of morbid obesity. Of course, one could speculate, whether the result concerning morbid obesity is a real finding or just a statistical coincidence. However, both FIN-D2D and the control population are selected as random independent samples of individuals, specified by region, gender, and age group, being thus a representative sample of the population. Similar changes in the prevalence of morbid obesity were observed in all sub-regions, i.e. the prevalence of morbid obesity increased in all sub-regions of the control area, and remained unchanged, in all FIN-D2D hospital districts.

The prevalence of severe ($\text{BMI} \geq 35 \text{ kg/m}^2$) and morbid ($\text{BMI} \geq 40 \text{ kg/m}^2$) obesity has increased in Finland 2.7-fold and 5.6-fold, respectively, during 1978-2001 [47]. Although the number of massively obese individuals is still rather low in Finland, it would be of importance to restrain their number from increasing further. In USA the prevalence of morbid obesity is 5.4% in the age group of 40 or older [3], which is nearly three times the prevalence in Finland (1.9% with FIN-D2D and FINRISK studies combined). Although rather rare, this extreme group of people incur huge expenses along with increasing surgical treatment of obesity.

The present results indicate that the obesity epidemic might be stabilizing or even decreasing in Finland. If

this trend is sustainable, also the incidence of type 2 diabetes might follow the obesity trends, and ease off during the years to come. To achieve this goal, we must now be able to maintain and to continue the progression already made regarding body weight. Furthermore, it would also be very important to be able to spread healthier lifestyles into the younger age groups, since obesity appears to reduce life expectancy markedly, especially among younger adults [48].

Limitations of the study

There were some differences in the age group distribution, since the oldest age group was not included in certain baseline study areas. However, since Turku-Loimaa and Kainuu belong to the control area, and Northern Savo and Northern Ostrobothnia belong to the FIN-D2D area, the age group distribution remains similar between the study groups. These areas represent 39% of the control area population, and 31% of the FIN-D2D area population in the follow-up. In all statistical analysis, age was included as a cofactor. The participation rates were rather low, especially in the control area. We do not know the reason for this. Moreover, there were some differences between the areas concerning certain background factors. Larger urban areas were not evenly distributed between the FIN-D2D and the control area, with the control area including for example the big cities of Helsinki and Turku. Individuals in the FIN-D2D area had slightly shorter duration of education and their cigarette smoking was somewhat less frequent than in the control area, whereas no difference in leisure-time physical activity was seen. When these factors were taken into account, the observed interactions for study area \times time point were still significant for waist circumference ($p = 0.013$) and morbid obesity ($p = 0.028$), when both genders were combined. Furthermore, the interaction concerning waist circumference among women became significant with these adjustments ($p = 0.014$), whereas it was not significant without these adjustments ($p = 0.062$). Adding the variable of sub-region into the statistical models had no effect, and was thus left out. Thus, we can conclude that the differences between the areas were not due to these background factors.

Conclusions

In conclusion, the overall body weight and the prevalence of obesity have been decreasing among Finns aged 45 to 74 years, both at the FIN-D2D and the control area. The significant differences in the prevalence of morbid obesity and mean waist circumference observed between the FIN-D2D area and control area suggest that the implementation of a national diabetes prevention programme may have been successful in the five hospital districts where this programme was established.

However, since most of the findings were not significant, we can not be certain of the true extent of the change, and future surveys may give additional data on the direction of this trend in Finland.

Acknowledgements

FIN-D2D study has been financially supported by the hospital districts of Pirkanmaa, South Ostrobothnia, Northern Ostrobothnia, Central Finland and Northern Savo, the Finnish National Public Health Institute (current National Institute for Health and Welfare), the Finnish Diabetes Association, the Ministry of Social Affairs and Health in Finland, the Academy of Finland (grant number 129293), Commission of the European Communities, Directorate C-Public Health (grant agreement no. 2004310) and Finland's Slottery Machine Association in cooperation with the FIN-D2D Study Group, and the Steering Committee: Pihlajamäki J, Huttunen J, Kesäniemi A, Kiuru S, Niskanen L, Oksa H, Puolakka J, Puska P, Saaristo T, Vanhala M and Uusitupa M.

Author details

¹Pirkanmaa Hospital District, Tampere, Finland. ²Diabetes Prevention Unit, Department of Chronic Disease Prevention, National Institute for Health and Welfare, Helsinki, Finland. ³Finnish Diabetes Association, Tampere, Finland. ⁴South Ostrobothnia Hospital District, Seinäjoki, Finland. ⁵Unit of Family Practice, Central Finland Hospital District, Jyväskylä, Finland. ⁶Institute of Public Health and Clinical Nutrition, University of Eastern Finland, Kuopio, Finland. ⁷Unit of Primary Health Care, Kuopio University Hospital, Kuopio, Finland. ⁸Oulu University Hospital, Northern Ostrobothnia Hospital District, Oulu, Finland. ⁹Department of Medicine, Kuopio University Hospital, Hospital District of Northern Savo, Kuopio, Finland. ¹⁰Department of Public Health, University of Helsinki, Helsinki, Finland. ¹¹Research Unit, Kuopio University Hospital, Kuopio, Finland.

Authors' contributions

TMS, MP, TS, and MU made substantial contributions to conception, design, and interpretation of the data and they were involved in drafting the manuscript. All authors participated in the design and conduction of the study, acquisition of data and helped to draft the manuscript or revise it. TMS performed the statistical analysis. All authors read and approved the final manuscript.

Competing interests

The authors declare that they have no competing interests.

Received: 27 September 2010 Accepted: 19 May 2011

Published: 19 May 2011

References

- Ogden CL, Carroll MD, Curtin LR, McDowell MA, Tabak CJ, Flegal KM: Prevalence of overweight and obesity in the United States, 1999-2004. *JAMA* 2006, **295**(13):1549-1555.
- Kautiainen S, Rimpelä A, Vikat A, Virtanen SM: Secular trends in overweight and obesity among Finnish adolescents in 1977-1999. *Int J Obes Relat Metab Disord* 2002, **26**(4):544-552.
- Flegal KM, Carroll MD, Ogden CL, Curtin LR: Prevalence and trends in obesity among US adults, 1999-2008. *JAMA* 2010, **303**(3):235-241.
- Vartiainen E, Peltonen M, Laatikainen T, Sundvall J, Salomaa V, Jousilahti P, Puska P: FINRISKI-tutkimus: Sekä miesten että naisten sydän- ja verisuonisairauksien kokonaisriski pieneni viime vuosina. *Suom Lääkäril* 2008, **63**(15):1375-1381.
- Trayhurn P, Bing C, Wood IS: Adipose tissue and adipokines - energy regulation from the human perspective. *J Nutr* 2006, **136**(7 Suppl):1935S-1939S.
- Anandacoomarasamy A, Caterson I, Sambrook P, Fransen M, March L: The impact of obesity on the musculoskeletal system. *Int J Obes (Lond)* 2008, **32**(2):211-222.
- Popkin BM, Kim S, Rusev ER, Du S, Zizza C: Measuring the full economic costs of diet, physical activity and obesity-related chronic diseases. *Obes Rev* 2006, **7**(3):271-293.
- Adler A: Obesity and target organ damage: diabetes. *Int J Obes Relat Metab Disord* 2002, **26**(Suppl 4):S11-14.
- Cummings DE, Schwartz MW: Genetics and pathophysiology of human obesity. *Annu Rev Med* 2003, **54**:453-471.
- DECODE Study Group: Age- and sex-specific prevalences of diabetes and impaired glucose regulation in 13 European cohorts. *Diabetes Care* 2003, **26**(1):61-69.
- Glucose tolerance and mortality: comparison of WHO and American Diabetes Association diagnostic criteria. The DECODE study group. European Diabetes Epidemiology Group: **Diabetes Epidemiology: Collaborative analysis Of Diagnostic criteria in Europe.** *Lancet* 1999, **354**(9179):617-621.
- Wild S, Roglic G, Green A, Sicree R, King H: Global prevalence of diabetes: estimates for the year 2000 and projections for 2030. *Diabetes Care* 2004, **27**(5):1047-1053.
- Tuomilehto J, Lindström J, Eriksson JG, Valle TT, Hämäläinen H, Ilanne-Parikka P, Keinänen-Kiukaanniemi S, Laakso M, Louheranta A, Rastas M, Salminen V, Uusitupa M, Finnish Diabetes Prevention Study Group: Prevention of type 2 diabetes mellitus by changes in lifestyle among subjects with impaired glucose tolerance. *N Engl J Med* 2001, **344**(18):1343-1350.
- Knowler WC, Barrett-Connor E, Fowler SE, Hamman RF, Lachin JM, Walker EA, Nathan DM, Diabetes Prevention Program Research Group: Reduction in the incidence of type 2 diabetes with lifestyle intervention or metformin. *N Engl J Med* 2002, **346**(6):393-403.
- Pan XR, Li GW, Hu YH, Wang JX, Yang WY, An ZX, Hu ZX, Lin J, Xiao JZ, Cao HB, Liu PA, Jiang XG, Jiang YY, Wang JP, Zheng H, Zhang H, Bennett PH, Howard BV: Effects of diet and exercise in preventing NIDDM in people with impaired glucose tolerance. The Da Qing IGT and Diabetes Study. *Diabetes Care* 1997, **20**(4):537-544.
- Kosaka K, Noda M, Kuzuya T: Prevention of type 2 diabetes by lifestyle intervention: a Japanese trial in IGT males. *Diabetes Res Clin Pract* 2005, **67**(2):152-162.
- Ramachandran A, Snehalatha C, Mary S, Mukesh B, Bhaskar AD, Vijay V, Indian Diabetes Prevention Programme (IDPP): **The Indian Diabetes Prevention Programme shows that lifestyle modification and metformin prevent type 2 diabetes in Asian Indian subjects with impaired glucose tolerance (IDPP-1).** *Diabetologia* 2006, **49**(2):289-297.
- Li G, Zhang P, Wang J, Gregg EW, Yang W, Gong Q, Li H, Jiang Y, An Y, Shuai Y, Zhang B, Zhang J, Thompson TJ, Gerzoff RB, Roglic G, Hu Y, Bennett PH: **The long-term effect of lifestyle interventions to prevent diabetes in the China Da Qing Diabetes Prevention Study: a 20-year follow-up study.** *Lancet* 2008, **371**(9626):1783-1789.
- Lindström J, Ilanne-Parikka P, Peltonen M, Aunola S, Eriksson JG, Hemio K, Hämäläinen H, Härkönen P, Keinänen-Kiukaanniemi S, Laakso M, Louheranta A, Mannelin M, Paturi M, Sundvall J, Valle TT, Uusitupa M, Tuomilehto J, Finnish Diabetes Prevention Study Group: **Sustained reduction in the incidence of type 2 diabetes by lifestyle intervention: follow-up of the Finnish Diabetes Prevention Study.** *Lancet* 2006, **368**(9548):1673-1679.
- Diabetes Prevention Program Research Group, Knowler WC, Fowler SE, Hamman RF, Christophi CA, Hoffman HJ, Brenneman AT, Brown-Friday JO, Goldberg R, Venditti E, Nathan DM: **10-year follow-up of diabetes incidence and weight loss in the Diabetes Prevention Program Outcomes Study.** *Lancet* 2009, **374**(9702):1677-1686.
- Cardona-Morrell M, Rychetnik L, Morrell SL, Espinel PT, Bauman A: Reduction of diabetes risk in routine clinical practice: are physical activity and nutrition interventions feasible and are the outcomes from reference trials replicable? A systematic review and meta-analysis. *BMC Public Health* 2010, **10**:653.
- Programme for the Prevention of Type 2 Diabetes in Finland 2003-2010.** [http://www.diabetes.fi/files/1108/Programme_for_the_Prevention_of_Type_2_Diabetes_in_Finland_2003-2010.pdf].
- Saaristo T, Peltonen M, Keinänen-Kiukaanniemi S, Vanhala M, Saltevo J, Niskanen L, Oksa H, Korpi-Hyövälti E, Tuomilehto J, FIN-D2D Study Group: **National type 2 diabetes prevention programme in Finland: FIN-D2D.** *Int J Circumpolar Health* 2007, **66**(2):101-112.
- Saaristo TE, Barengo NC, Korpi-Hyövälti E, Oksa H, Puolijoki H, Saltevo JT, Vanhala M, Sundvall J, Saarikoski L, Peltonen M, Tuomilehto J: **High prevalence of obesity, central obesity and abnormal glucose tolerance in the middle-aged Finnish population.** *BMC Public Health* 2008, **8**:423.
- Implementation of Type 2 Diabetes Prevention. Project Plan 2003-2007.** [http://www.diabetes.fi/files/1107/Implementation_of_Type_2_Diabetes_Prevention_Project_Plan_2003-2007.pdf].

26. Saaristo T, Moilanen L, Korpi-Hyövälti E, Vanhala M, Saltevo J, Niiskanen L, Jokelainen J, Peltonen M, Oksa H, Tuomilehto J, Uusitupa M, Keinänen-Kiukaanniemi S: **Lifestyle intervention for prevention of type 2 diabetes in primary health care: one-year follow-up of the Finnish National Diabetes Prevention Program (FIN-D2D).** *Diabetes Care* 2010, **33**(10):2146-2151.
27. Puska P, Koskela K, Pakarinen H, Puumalainen P, Soininen V, Tuomilehto J: **The North Karelia Project: a programme for community control of cardiovascular diseases.** *Scand J Soc Med* 1976, **4**(2):57-60.
28. Vartiainen E, Jousilahti P, Alftan G, Sundvall J, Pietinen P, Puska P: **Cardiovascular risk factor changes in Finland, 1972-1997.** *Int J Epidemiol* 2000, **29**(1):49-56.
29. **The World Health Organization MONICA Project (monitoring trends and determinants in cardiovascular disease): a major international collaboration.** WHO MONICA Project Principal Investigators. *J Clin Epidemiol* 1988, **41**(2):105-114.
30. Alberti KG, Zimmet P, Shaw J, IDF Epidemiology Task Force Consensus Group: **The metabolic syndrome - a new worldwide definition.** *Lancet* 2005, **366**(9491):1059-1062.
31. Han TS, van Leer EM, Seidell JC, Lean ME: **Waist circumference action levels in the identification of cardiovascular risk factors: prevalence study in a random sample.** *BMJ* 1995, **311**(7017):1401-1405.
32. Lahti-Koski M, Harald K, Männistö S, Laatikainen T, Jousilahti P: **Fifteen-year changes in body mass index and waist circumference in Finnish adults.** *Eur J Cardiovasc Prev Rehabil* 2007, **14**(3):398-404.
33. Berghofer A, Pischon T, Reinhold T, Apovian CM, Sharma AM, Willich SN: **Obesity prevalence from a European perspective: a systematic review.** *BMC Public Health* 2008, **8**:200.
34. Aranceta J, Perez-Rodrigo C, Serra-Majem L, Bellido D, de la Torre ML, Formiguera X, Moreno B: **Prevention of overweight and obesity: a Spanish approach.** *Public Health Nutr* 2007, **10**(10A):1187-1193.
35. Katzmarzyk PT: **The Canadian obesity epidemic: an historical perspective.** *Obes Res* 2002, **10**(7):666-674.
36. Bendixen H, Holst C, Sorensen TI, Raben A, Bartels EM, Astrup A: **Major increase in prevalence of overweight and obesity between 1987 and 2001 among Danish adults.** *Obes Res* 2004, **12**(9):1464-1472.
37. Marques-Vidal P, Dias CM: **Trends in overweight and obesity in Portugal: the National Health Surveys 1995-6 and 1998-9.** *Obes Res* 2005, **13**(7):1141-1145.
38. Dal Grande E, Gill T, Taylor AW, Chittleborough C, Carter P: **Obesity in South Australian adults - prevalence, projections and generational assessment over 13 years.** *Aust N Z J Public Health* 2005, **29**(4):343-348.
39. Sundquist J, Johansson SE, Sundquist K: **Leveling off of prevalence of obesity in the adult population of Sweden between 2000/01 and 2004/05.** *BMC Public Health* 2010, **10**(1):119.
40. Sundquist K, Qvist J, Johansson SE, Sundquist J: **Increasing trends of obesity in Sweden between 1996/97 and 2000/01.** *Int J Obes Relat Metab Disord* 2004, **28**(2):254-261.
41. Borodulin K, Laatikainen T, Juolevi A, Jousilahti P: **Thirty-year trends of physical activity in relation to age, calendar time and birth cohort in Finnish adults.** *Eur J Public Health* 2008, **18**(3):339-344.
42. **Health Behaviour and Health among the Finnish Adult Population, Spring 2007.** [http://www.ktl.fi/attachments/suomi/julkaisut/julkaisusarja_b/2008/2008b06.pdf].
43. Assaf AR, Parker D, Lapane KL, Coccio E, Evangelou E, Carleton RA: **Does the Y chromosome make a difference? Gender differences in attempts to change cardiovascular disease risk factors.** *J Womens Health (Larchmt)* 2003, **12**(4):321-330.
44. Rejeski WJ, Brawley LR, Ambrosius WT, Brubaker PH, Focht BC, Foy CG, Fox LD: **Older adults with chronic disease: benefits of group-mediated counseling in the promotion of physically active lifestyles.** *Health Psychol* 2003, **22**(4):414-423.
45. Hankonen N, Absetz P, Ghisletta P, Renner B, Uutela A: **Gender differences in social cognitive determinants of exercise adoption.** *Psychology & Health* 2009, 1-15.
46. Wong M, Gucciardi E, Li L, Grace SL: **Gender and nutrition management in type 2 diabetes.** *Can J Diet Pract Res* 2005, **66**(4):215-220.
47. Lahti-Koski M, Seppänen-Nuijten E, Männistö S, Härkänen T, Rissanen H, Knekt P, Rissanen A, Heliövaara M: **Twenty-year changes in the prevalence of obesity among Finnish adults.** *Obes Rev* 2009, **11**:171-176.
48. Fontaine KR, Redden DT, Wang C, Westfall AO, Allison DB: **Years of life lost due to obesity.** *JAMA* 2003, **289**(2):187-193.

Pre-publication history

The pre-publication history for this paper can be accessed here:
<http://www.biomedcentral.com/1471-2458/11/350/prepub>

doi:10.1186/1471-2458-11-350

Cite this article as: Salopuro et al.: Population-level effects of the national diabetes prevention programme (FIN-D2D) on the body weight, the waist circumference, and the prevalence of obesity. *BMC Public Health* 2011 **11**:350.

Submit your next manuscript to BioMed Central and take full advantage of:

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit

