

ISSN: 1303 - 2968

 ISI 2013 IF 2-Year: 0.90 5-Year: 1.34 Average Citations per item: 4.2  
 SCOPUS 2013 SJR: 0.56 Cites per Doc. 2-Year: 1.05 3-Year: 1.38 4-Year: 1.57

[Home](#) [Mission](#) [Scope](#) [Editorial Board](#) [For Reviewers](#) [Submission](#) [Statistics](#) [Contact](#)

FIND ARTICLES

Search on JSSM

Search

Advanced Search &gt;&gt;&gt;

- [Current Issue](#)  [RSS](#)
- [In Press](#)  [RSS](#)

● [Back Issues](#)

## Back Issue

### June 2014 - Volume 13, Issue 2

#### Table of Contents

##### Research article

252 - 258

#### The Influence of Serial Carbohydrate Mouth Rinsing on Power Output during a Cycle Sprint

Shaun M. Phillips, Scott Findlay, Mykolas Kavaliauskas, Marie Clare Grant

[Abstract](#)[Full text](#)[PDF](#)

##### Research article

259 - 265

#### Effects and Sustainability of a 13-Day High-Intensity Shock Microcycle in Soccer

Patrick Wahl, Matthias Güldner, Joachim Mester

[Abstract](#)[Full text](#)[PDF](#)

##### Research article

266 - 270

#### An Enjoyable Distraction During Exercise Augments the Positive Effects of Exercise on Mood

Gregory J. Privitera, Danielle E. Antonelli, Abigail L. Szal

[Abstract](#)[Full text](#)[PDF](#)

##### Research article

271 - 279

#### Acute Impact of Inhaled Short Acting B-Agonists on 5 Km Running Performance

John Dickinson, Jiu Hu, Neil Chester, Mike Loosemore, Greg Whyte

[Abstract](#)[Full text](#)[PDF](#)

##### Research article

280 - 286

#### Acute Differences in Foot Strike and Spatiotemporal Variables for Shod, Barefoot or Minimalist Male Runners

Indexed in SCI Expanded, Focus on: Sports Science & Medicine, SciSearch, PubMed, PubMed Central, ProQuest (Physical Education Index), CAB Abstracts, SCOPUS, Index Copernicus, EBSCO, SPORTDiscus, DOAJ, J-Gate, GALE, GoogleScholar, ERA and SPONET

#### Most Readed Articles

#### Most Cited Articles

#### In Press

##### [The Effect of Fatigue on Electromyographic Characteristics during Obstacle Crossing of Different Heights in Young Adults](#)

Christos Antonopoulos, Dimitrios Patikas, Nikolaos Koutlianos, Sofia D. Papadopoulou, Dimitrios Chatzopoulos, Konstantinos Hatzikotoulas, Eleni Bassa, Christos Kotzamanidis

##### [Shuttle-Run Sprint Training in Hypoxia for Youth Elite Soccer Players: A Pilot Study](#)

Hannes Gatterer, Marc Philippe, Verena Menz, Florian Mosbach, Martin Faulhaber, Martin Burtcher

##### [Knowledge of Repetitions Range Affects Force Production in Trained Females](#)

Israel Halperin, Saied J. Aboodarda, Fabien A. Basset, David G. Behm

##### [Kinematic and Kinetic Profiles of Trunk and Lower Limbs during Baseball Pitching in Collegiate Pitchers](#)

Masahiro Kageyama, Takashi Sugiyama,

JSSM | Copyright 2001-2014 | All rights reserved.

It is forbidden the total or partial reproduction of this web site and the published materials, the treatment of its database, any kind of transition and for any means, either electronic, mechanic or other methods, without the previous written permission of the JSSM.

**Volume 13(2); May 2014****Case Reports**

---

[Could Low-Frequency Electromyostimulation Training be an Effective Alternative to Endurance Training? An Overview in One Adult](#)

Gaëlle Deley, Nicolas Babault

J Sports Sci Med. May 2014; 13(2): 444–450. Published online May 1, 2014.

PMCID: PMC3990903

[Article](#) [PubReader](#) [PDF-322K](#)[An Unusual Case of Traumatic Internal Carotid Artery Dissection during Snowboarding](#)

George Kalantzis, Ilias Georgalas, Bernard Y.P. Chang, Chin Ong, Nabil El-Hindy

J Sports Sci Med. May 2014; 13(2): 451–453. Published online May 1, 2014.

PMCID: PMC3990904

[Article](#) [PubReader](#) [PDF-187K](#)**Letters to Editors**

---

[Lack of Enjoyment Reduces the Motivation to Succeed in Sport](#)

Carra Johnson, Ronald J. Peters, Jr.

J Sports Sci Med. May 2014; 13(2): 454–455. Published online May 1, 2014.

PMCID: PMC3990905

[Article](#) [PubReader](#) [PDF-112K](#)[Stereotypes of Athletes' Use of Performance Enhancing Products](#)

Carra Johnson, Ronald J. Peters, Jr.

J Sports Sci Med. May 2014; 13(2): 456–457. Published online May 1, 2014.

PMCID: PMC3990906

[Article](#) [PubReader](#) [PDF-110K](#)[Reply to Letter to Editor: Stereotypes of Athletes' Use of Performance Enhancing Products](#)

Nkaku Kisaalita, Michael E. Robinson

J Sports Sci Med. May 2014; 13(2): 458–459. Published online May 1, 2014.

PMCID: PMC3990907

[Article](#) [PubReader](#) [PDF-238K](#)**Research Articles**

---

[The Paroxetine Effect on Exercise Performance Depends on the Aerobic Capacity of Exercising Individuals](#)

Francisco Teixeira-Coelho, João Paulo Uendele-Pinto, Ana Cláudia Alves Serafim, Samuel Penna Wanner, Márcio de Matos Coelho, Danusa Dias Soares

J Sports Sci Med. May 2014; 13(2): 232–243. Published online May 1, 2014.

PMCID: PMC3990874

[Article](#) [PubReader](#) [PDF-533K](#)[The Occurrence of Core Muscle Fatigue During High-Intensity Running Exercise and its Limitation to Performance: The Role of Respiratory Work](#)

Tomas K. Tong, Shing Wu, Jinlei Nie, Julien S. Baker, Hua Lin

J Sports Sci Med. May 2014; 13(2): 244–251. Published online May 1, 2014.

PMCID: PMC3990875

[Article](#) [PubReader](#) [PDF-384K](#)[The Influence of Serial Carbohydrate Mouth Rinsing on Power Output during a Cycle Sprint](#)

Shaun M. Phillips, Scott Findlay, Mykolas Kavaliauskas, Marie Clare Grant

J Sports Sci Med. May 2014; 13(2): 252–258. Published online May 1, 2014.

PMCID: PMC3990876

[Article](#) [PubReader](#) [PDF-290K](#)[Effects and Sustainability of a 13-Day High-Intensity Shock Microcycle in Soccer](#)

Patrick Wahl, Matthias Güldner, Joachim Mester

J Sports Sci Med. May 2014; 13(2): 259–265. Published online May 1, 2014.

PMCID: PMC3990877

[Article](#) [PubReader](#) [PDF-384K](#)

### [An Enjoyable Distraction During Exercise Augments the Positive Effects of Exercise on Mood](#)

Gregory J. Privitera, Danielle E. Antonelli, Abigail L. Szal

J Sports Sci Med. May 2014; 13(2): 266–270. Published online May 1, 2014.

PMCID: PMC3990878

[Article](#) [PubReader](#) [PDF-258K](#)

### [Acute Impact of Inhaled Short Acting B<sub>2</sub>-Agonists on 5 Km Running Performance](#)

John Dickinson, Jiu Hu, Neil Chester, Mike Loosemore, Greg Whyte

J Sports Sci Med. May 2014; 13(2): 271–279. Published online May 1, 2014.

PMCID: PMC3990879

[Article](#) [PubReader](#) [PDF-458K](#)

### [Acute Differences in Foot Strike and Spatiotemporal Variables for Shod, Barefoot or Minimalist Male Runners](#)

Ciara McCallion, Bernard Donne, Neil Fleming, Brian Blanksby

J Sports Sci Med. May 2014; 13(2): 280–286. Published online May 1, 2014.

PMCID: PMC3990880

[Article](#) [PubReader](#) [PDF-301K](#)

### [The Effects of Interset Rest on Adaptation to 7 Weeks of Explosive Training in Young Soccer Players](#)

Rodrigo Ramirez-Campillo, David C. Andrade, Cristian Álvarez, Carlos Henríquez-Olguín, Cristian Martínez, Eduardo Báez-SanMartín, Juan Silva-Urra, Carlos Burgos, Mikel Izquierdo

J Sports Sci Med. May 2014; 13(2): 287–296. Published online May 1, 2014.

PMCID: PMC3990881

[Article](#) [PubReader](#) [PDF-293K](#)

### [Effect of Court Dimensions on Players' External and Internal Load during Small-Sided Handball Games](#)

Matteo Corvino, Antonio Tessitore, Carlo Minganti, Marko Sibila

J Sports Sci Med. May 2014; 13(2): 297–303. Published online May 1, 2014.

PMCID: PMC3990882

[Article](#) [PubReader](#) [PDF-236K](#)

### [Ball Machine Usage in Tennis: Movement Initiation and Swing Timing While Returning Balls from a Ball Machine and from a Real Server](#)

Jan Carboch, Vladimir Süß, Tomas Kocib

J Sports Sci Med. May 2014; 13(2): 304–308. Published online May 1, 2014.

PMCID: PMC3990883

[Article](#) [PubReader](#) [PDF-371K](#)

### [A Simplified Approach for Estimating the Ventilatory and Respiratory Compensation Thresholds](#)

Giancarlo Condello, Ezekiel Reynolds, Carl Foster, Jos J. de Koning, Erika Casolino, Megan Knutson, John P. Porcari

J Sports Sci Med. May 2014; 13(2): 309–314. Published online May 1, 2014.

PMCID: PMC3990884

[Article](#) [PubReader](#) [PDF-791K](#)

### [Does Acute Vibration Exercise Enhance Horizontal Jump Performance?](#)

Darryl J. Cochrane, Hayden Booker

J Sports Sci Med. May 2014; 13(2): 315–320. Published online May 1, 2014.

PMCID: PMC3990885

[Article](#) [PubReader](#) [PDF-241K](#)

### [Efficacy of the FIFA 11+ Warm-Up Programme in Male Youth Football: A Cluster Randomised Controlled Trial](#)

Oluwatoyosi B. A. Owoeye, Sunday R. A. Akinbo, Bosede A. Tella, Olajide A. Olawale

J Sports Sci Med. May 2014; 13(2): 321–328. Published online May 1, 2014.

PMCID: PMC3990886

[Article](#) [PubReader](#) [PDF-226K](#)

### [The Effects of Cervical Muscle Fatigue on Balance – A Study with Elite Amateur Rugby League Players](#)

Guy Gosselin, Michael J. Fagan

J Sports Sci Med. May 2014; 13(2): 329–337. Published online May 1, 2014.

PMCID: PMC3990887

[Article](#) [PubReader](#) [PDF-405K](#)

### [Evaluation of the Physical Activity Biography: Sport and Transport](#)

Sandra Rogen, Peter Hofmann, Thomas Bauernhofer, Wolfram Müller  
J Sports Sci Med. May 2014; 13(2): 338–348. Published online May 1, 2014.  
PMCID: PMC3990888  
[Article](#) [PubReader](#) [PDF-295K](#)

[The Association of Flexibility, Balance, and Lumbar Strength with Balance Ability: Risk of Falls in Older Adults](#)

Emilio J. Martínez-López Emilio, Fidel Hita-Contreras, Pilar M. Jiménez-Lara, Pedro Latorre-Román, Antonio Martínez-Amat  
J Sports Sci Med. May 2014; 13(2): 349–357. Published online May 1, 2014.  
PMCID: PMC3990889  
[Article](#) [PubReader](#) [PDF-480K](#)

[Performance Indicators Related to Points Scoring and Winning in International Rugby Sevens](#)

Dean G. Higham, Will G. Hopkins, David B. Pyne, Judith M. Anson  
J Sports Sci Med. May 2014; 13(2): 358–364. Published online May 1, 2014.  
PMCID: PMC3990890  
[Article](#) [PubReader](#) [PDF-294K](#)

[Assessment of Isometric Trunk Strength – The Relevance of Body Position and Relationship between Planes of Movement](#)

Andrej Kocjan, Nejc Sarabon  
J Sports Sci Med. May 2014; 13(2): 365–370. Published online May 1, 2014.  
PMCID: PMC3990891  
[Article](#) [PubReader](#) [PDF-289K](#)

[Physical Activity and Sedentary Behavior in an Ethnically Diverse Group of South African School Children](#)

Joanne McVeigh, Rebecca Meiring  
J Sports Sci Med. May 2014; 13(2): 371–378. Published online May 1, 2014.  
PMCID: PMC3990892  
[Article](#) [PubReader](#) [PDF-283K](#)

[Towards Uniform Accelerometry Analysis: A Standardization Methodology to Minimize Measurement Bias Due to Systematic Accelerometer Wear-Time Variation](#)

Tarun R. Katapally, Nazeem Muhajarine  
J Sports Sci Med. May 2014; 13(2): 379–386. Published online May 1, 2014.  
PMCID: PMC3990893  
[Article](#) [PubReader](#) [PDF-776K](#)

[Learning Effects Associated With the Least Stable Level of the Biodex® Stability System During Dual and Single Limb Stance](#)

Mutlu Cug, Erik A. Wikstrom  
J Sports Sci Med. May 2014; 13(2): 387–392. Published online May 1, 2014.  
PMCID: PMC3990894  
[Article](#) [PubReader](#) [PDF-230K](#)

[Moderate Recovery Unnecessary to Sustain High Stroke Volume during Interval Training. A Brief Report](#)

Jamie Stanley, Martin Buchheit  
J Sports Sci Med. May 2014; 13(2): 393–396. Published online May 1, 2014.  
PMCID: PMC3990895  
[Article](#) [PubReader](#) [PDF-216K](#)

[Balance Training Exercises Decrease Lower-Limb Strength Asymmetry in Young Tennis Players](#)

Italo Sannicandro, Giacomo Cofano, Rosa A. Rosa, Andrea Piccinno  
J Sports Sci Med. May 2014; 13(2): 397–402. Published online May 1, 2014.  
PMCID: PMC3990896  
[Article](#) [PubReader](#) [PDF-243K](#)

[Acute Effects of Static and Dynamic Stretching on Balance, Agility, Reaction Time and Movement Time](#)

Dimitris Chatzopoulos, Christos Galazoulas, Dimitrios Patikas, Christos Kotzamanidis  
J Sports Sci Med. May 2014; 13(2): 403–409. Published online May 1, 2014.  
PMCID: PMC3990897  
[Article](#) [PubReader](#) [PDF-271K](#)

[Physiological Responses during Cycling With Oval Chainrings \(Q-Ring\) and Circular Chainrings](#)

Alfredo Cordova, Iban Latasa, Jesus Seco, Gerardo Villa, Javier Rodriguez-Falces  
J Sports Sci Med. May 2014; 13(2): 410–416. Published online May 1, 2014.  
PMCID: PMC3990898  
[Article](#) [PubReader](#) [PDF-452K](#)

[Comparative Kinematic Analysis of the Snatch Lifts in Elite Male Adolescent Weightlifters](#)

Erbil Harbili, Ahmet Alptekin

J Sports Sci Med. May 2014; 13(2): 417–422. Published online May 1, 2014.

PMCID: PMC3990899

[Article](#) [PubReader](#) [PDF-308K](#)[What is the Safest Sprint Starting Position for American Football Players?](#)

Bruno Bonnechere, Benoit Beyer, Marcel Rooze, Jan Serge Van Sint

J Sports Sci Med. May 2014; 13(2): 423–429. Published online May 1, 2014.

PMCID: PMC3990900

[Article](#) [PubReader](#) [PDF-2.3M](#)[Cross-Cultural Adaptation and Validation of the Spanish Version of the Performance Enhancement Attitude Scale \(Petróczy.\)](#)

Jaime Morente-Sánchez, Pedro Femia-Marzo, Mikel Zabala

J Sports Sci Med. May 2014; 13(2): 430–438. Published online May 1, 2014.

PMCID: PMC3990901

[Article](#) [PubReader](#) [PDF-280K](#)[Correct, Fake and Absent Pre-Information Does Not Affect the Occurrence and Magnitude of the Bilateral Force Deficit](#)

Lars Donath, Tobias Siebert, Oliver Faude, Christian Puta

J Sports Sci Med. May 2014; 13(2): 439–443. Published online May 1, 2014.

PMCID: PMC3990902

[Article](#) [PubReader](#) [PDF-259K](#)

## Review Article

---

[Biomechanical Analysis of the Swim-Start: A Review](#)

Julien Vantorre, Didier Chollet, Ludovic Seifert

J Sports Sci Med. May 2014; 13(2): 223–231. Published online May 1, 2014.

PMCID: PMC3990873

[Article](#) [PubReader](#) [PDF-416K](#)

---

Articles from Journal of Sports Science & Medicine are provided here courtesy of **Dept. of Sports Medicine, Medical Faculty of Uludag University**

Research article

## Moderate Recovery Unnecessary to Sustain High Stroke Volume during Interval Training. A Brief Report

Jamie Stanley<sup>1,2</sup>✉ and Martin Buchheit<sup>3</sup>

<sup>1</sup> Centre of Excellence for Applied Sport Science Research, Queensland Academy of Sport, Brisbane, Australia; <sup>2</sup> The University of Queensland, School of Human Movement Studies, Brisbane, Australia; <sup>3</sup> Sport Science Unit, Myorobie Association, Montvalezan, France

### Abstract

It has been suggested that the time spent at a high stroke volume (SV) is important for improving maximal cardiac function. The aim of this study was to examine the effect of recovery intensity on cardiovascular parameters during a typical high-intensity interval training (HIIT) session in fourteen well-trained cyclists. Oxygen consumption ( $\text{VO}_2$ ), heart rate (HR), SV, cardiac output (Qc), and oxygenation of *vastus lateralis* (TSI) were measured during a HIIT (3×3-min work period, 2 min of recovery) session on two occasions.  $\text{VO}_2$ , HR and Qc were largely higher during moderate-intensity (60%) compared with low-intensity (30%) ( $\text{VO}_2$ , effect size; ES = +2.6; HR, ES = +2.8; Qc, ES = +2.2) and passive (HR, ES = +2.2; Qc, ES = +1.7) recovery. By contrast, there was no clear difference in SV between the three recovery conditions, with the SV during the two active recovery periods not being substantially different than during exercise (60%, ES = -0.1; 30%, ES = -0.2). To conclude, moderate-intensity recovery may not be required to maintain a high SV during HIIT.

**Key words:** High-intensity interval training; cardiac output; cardiac function; arteriovenous oxygen difference.

### Introduction

High-intensity interval training (HIIT) is recognized as one of the most effective means of improving cardiorespiratory and metabolic function in athletes (Buchheit and Laursen, 2013). A typical HIIT session involves the repetition of periods of high-intensity exercise (i.e., the intervals) interspersed with periods of lower intensity (i.e., recovery periods) (Buchheit and Laursen, 2013). Long intervals (i.e., >3 min) are typically completed close to the speed/power associated with maximal oxygen consumption ( $\text{VO}_{2\text{max}}$ ), because it is believed that this is optimal stimulus for eliciting maximal cardiovascular and peripheral adaptations (Buchheit and Laursen, 2013). However, because  $\text{VO}_2$  and cardiac output (Qc) can be dissociated during intense exercise (Lepretre et al., 2004), and attaining and maintaining an elevated stroke volume (SV) is likely important for improving maximal cardiac function (Cooper, 1997), increasing time spent at maximal Qc (Qcmax) and/or training at an intensity associated with maximal SV may also be important (Lepretre et al., 2004).

The intensity of exercise that maximizes the time at maximal SV is difficult to predict (González-Alonso, 2008; Mortensen et al., 2005; Warburton and Gledhill,

2008). The most appropriate HIIT format inducing increased time at Qcmax in well-trained and elite athletes remains to be determined. In untrained males, compared with peak exercise during a graded exercise test, 30-s all-out sprints (typical of sprint interval training sessions (Buchheit and Laursen, 2013)), might allow attainment of similar Qc (effect size; ES = -0.1) and even largely higher SV (ES = +1.3) (Fontana et al., 2011). The recovery period is another key factor of HIIT with respect to cardiopulmonary responses (Buchheit and Laursen, 2013). Conjecture remains as to whether maximal SV is reached during the recovery period or during work periods—and whether this response is recovery-intensity dependent (Buchheit and Laursen, 2013; Cumming, 1972; González-Alonso, 2008; Warburton and Gledhill, 2008). Therefore, the aim of the current study was to examine the effect of recovery intensity on cardiovascular parameters during HIIT.

### Methods

#### Subjects

Fourteen endurance-trained male cyclists participated in the study (age,  $25 \pm 4$  years; body mass,  $69.6 \pm 4.9$  kg; height,  $1.77 \pm 0.04$  m;  $\text{VO}_{2\text{max}}$ ,  $66.6 \pm 4.2$  mL·kg<sup>-1</sup>·min<sup>-1</sup>; peak power output,  $405 \pm 28$  W). The experimental procedure was approved by the Human Research Ethics Committee at The University of Queensland.

#### Experimental overview

The complete methodology has been reported elsewhere (Stanley et al., 2014). Briefly, the cyclists completed an incremental cycling test to determine the power at  $\text{VO}_{2\text{max}}$  ( $\text{pVO}_{2\text{max}}$ ) and maximal Qc (Qcmax), and two cycling (Wattbike®, Wattbike Ltd., UK) HIIT sessions (12-min warm-up, three 3-min work periods (Ex) at 90%  $\text{pVO}_{2\text{max}}$ ) 5–7 days apart. The 2-min recovery periods were completed at either 30%  $\text{pVO}_{2\text{max}}$  (30%) or 60%  $\text{pVO}_{2\text{max}}$  (60%) in a randomized order with passive (PAS) recovery always following the final interval. Respiratory gas exchange (ParvoMedics TrueOne® 2400, Utah, USA) and HR, SV and Qc (PhysioFlow, Manatec Biomedical, France) (Charloux et al., 2000) were measured continuously. Oxygenation of *vastus lateralis* (tissue saturation index; TSI) was determined using near-infrared spectroscopy (PortaMon, Artinis Medical Systems BV, The Netherlands). Arteriovenous difference (a- $\text{vO}_2$ ) was calculated using the Fick equation. All data were sampled at 5-s

intervals, and the average for each work period (e.g., 3 min average for Ex, and 2 min average for the recovery period) was determined during the HIIT. Unfortunately due to technical difficulties,  $Q_{cmax}$  was only determined for 5 cyclists during the incremental test. The coefficient of variation (CV) for each variable during Ex was as follows: HR, 1.2%; SV, 5.5%;  $Q_c$ , 5.8%;  $VO_2$ , 3.4%; TSI, 9.9%; and  $a-vO_2$  diff, 2.5%.


### Data analysis

All data in the text and figures are presented as mean ( $n = 14$ ) or percentage ( $n = 5$ ) of incremental test peak values with 90% confidence limits (CL). Standardized differences/changes in all monitored variables were calculated using the pooled between-subject standard deviations. Threshold values were  $>0.2$  (small),  $>0.6$  (moderate) and  $>1.2$  (large) (Hopkins et al., 2009). Uncertainty in each effect was expressed as 90% CL and its probabilities that

the true difference was substantially greater or lower than the CV for each variable. These probabilities were used to make a qualitative mechanistic inference about the true effect (Hopkins et al., 2009). Data as a percentage of incremental test peak values were included to provide an indication of relative load and demonstrated similar trends to raw values. Statistical comparisons were not performed on data as a percentage of incremental test peak values because data was only available for 5 cyclists.

### Results

The effect of recovery intensity on cardiorespiratory variables (both mean and percentage of incremental test peak values) is displayed in Figure 1. No clear differences in SV were observed between recovery conditions. There was a possible small difference in SV between PAS and Ex.



**Figure 1.** Heart rate (HR), stroke volume (SV), cardiac output ( $Q_c$ ), oxygen consumption ( $VO_2$ ), arteriovenous oxygen difference ( $a-vO_2$  diff), and tissue saturation index (TSI) measured simultaneously during the different intensities during the HIIT session: pre-exercise rest (Rest), 3-min work periods at 90%  $pVO_{2max}$  (Ex), passive recovery (PAS), low-intensity recovery (30%), moderate-intensity recovery (60%). Data are presented as mean ( $n = 14$ ; circles) and percentage of incremental test peak values ( $n = 5$ ; squares) with 90% confidence intervals.  $VO_2$  values for Rest were calculated based on ( $5 \text{ mL}\cdot\text{kg}^{-1}$ ) (Medbø and Tabata, 1989) and were not available for PAS. TSI was not measured during rest or the max test. Letters 'a', 'b', and 'c' indicate a substantial difference versus PAS, 30%, and 60% respectively, with the number of letters representing a small (1 letter), moderate (2 letters), and large (3 letters) standardized differences respectively. If the 90% CL of the standardized differences overlapped small positive and negative values, the magnitude was deemed unclear; otherwise that magnitude was deemed to be the observed magnitude (Hopkins et al., 2009). For clarity, letters have been omitted for differences between Rest and all other intensities as all were clear and large. Data as a percentage of incremental test peaks were provided to indicate the relative load and demonstrated similar trends to raw values, however, because data was only available for 5 cyclists, statistical comparisons were not performed.

## Discussion

This is the first study to describe Qc and SV during a typical HIIT session performed by well-trained athletes. The main finding was that moderate-intensity recovery periods are not necessary to maintain high SV during the exercise intervals of HIIT.

In the present study, during the 3-min work periods at 90% pVO<sub>2max</sub>, Qc reached ~89% and SV reached ~102% of the peak values attained during the incremental test. Due to the limited number of data sets in the present study (n = 5), caution is advised when comparing these findings. Nevertheless, the Qc values in the present study are consistent with the ~85% observed during a 4-min exercise interval at an intensity ~90% pVO<sub>2max</sub> (Richard et al., 2004). However, present data contrast with that observed during a sprint interval session (30-s all-out sprints), where Qc reached ~102%, and SV ~134% of Qcmax and maximal SV respectively (Fontana et al., 2011). Nevertheless, HIIT is likely a better stimuli for cardiovascular improvements as the total time spent at or near maximal SV and VO<sub>2max</sub> is likely substantially greater during a HIIT session (e.g., 15-min HIIT: 3 × 3-min at 90% pVO<sub>2max</sub> interspersed with 2 min active recovery = >9 min near max) compared with a SIT session of comparable duration (e.g., 15-min SIT: 6 × 30-s all-out efforts interspersed with 2 min passive recovery (Buchheit and Laursen, 2013) = ~3 min near max). These data illustrate that cardiovascular responses during exercise are likely HIIT protocol-dependent.

During recovery periods we observed that SV did not surpass the levels attained during exercise (Figure 1). Because the statistical analysis included all 14 cyclists, we are confident with this finding. This is contrary to the belief that SV could reach maximal levels during the recovery periods of HIIT (Buchheit and Laursen, 2013; Cumming, 1972). In a well-trained cyclist, higher SV values were consistently observed during recovery periods, not work periods, irrespective of the type of exercise (incremental exercise, long intervals, or sprints) (Buchheit and Laursen, 2013). The reason for these disparities is not clear. However, differences in training status (well-trained vs. untrained) and mode of exercise (upright vs. supine) likely influenced venous return, cardiac preload, and peripheral resistance that can influence SV.

Interestingly, we observed that despite the expected differences in peripheral metabolic demands (as inferred from the greater a-vO<sub>2</sub> diff and lower TSI, figure 1), the SV response was not affected by recovery intensity (Figure 1). Our data suggest that SV already reached a plateau at a low intensity (González-Alonso, 2008; Warburton and Gledhill, 2008). Because active recovery involves rhythmical contraction of exercising muscles irrespective of intensity, it is likely that the effect of the muscle pump influenced peripheral resistance and venous return, such that a similarly high SV (compared with the exercise intervals) was maintained.

## Conclusion

The present study demonstrates that moderate-intensity

recovery periods are not necessary to maintain high SV during the exercise intervals HIIT. We acknowledge that the impedance method used to assess cardiovascular function in this study has not been validated during maximal exercise. Therefore, rather than compare our data in absolute terms (to previous studies), we have made comparisons with respect to difference in exercise intensity. Nevertheless, the CV for our data suggest similar reliability to that observed during sub-maximal exercise (Charloux et al., 2000). The practical implication of this finding is that reducing the intensity of the recovery period during a HIIT protocol may prolong the time to exhaustion (Dupont et al., 2003), potentially allowing completion of additional high-intensity intervals, which can, in turn, increase the time accumulated at Qcmax. Using the HIIT protocol in the current study as an example, completion of an additional two intervals would increase the time spent at Qcmax by 67%. Whether achieving a specific quantity of time at Qcmax is necessary to maximize cardiac adaptation remains unknown.

## Acknowledgements

The authors thank the cyclists for their generous time commitment and effort throughout the study. This study was supported by the Australian Institute of Sport High Performance Research Fund and the Centre of Excellence for Applied Sport Science Research at the Queensland Academy of Sport.

## References

- Buchheit, M. and Laursen, P.B. (2013) High-intensity interval training, solutions to the programming puzzle part i: Cardiopulmonary emphasis. *Sports Medicine* **43**, 313-338.
- Charloux, A., Lonsdorfer-Wolf, E., Richard, R., Lampert, E., Oswald-Mammosser, M., Mettauer, B., Geny, B. and Lonsdorfer, J. (2000) A new impedance cardiograph device for the non-invasive evaluation of cardiac output at rest and during exercise: Comparison with the "direct" fick method. *European Journal of Applied Physiology* **82**, 313-320.
- Cooper, G. (1997) Basic determinants of myocardial hypertrophy: A review of molecular mechanisms. *Annual Review of Medicine* **48**, 13-23.
- Cumming, G.R. (1972) Stroke volume during recovery from supine bicycle exercise. *Journal of Applied Physiology* **32**, 575-578.
- Dupont, G., Blondel, N. and Berthoin, S. (2003) Performance for short intermittent runs: Active recovery vs. Passive recovery. *European Journal of Applied Physiology* **89**, 548-554.
- Fontana, P., Betschon, K., Boutellier, U. & Toigo, M. (2011). Cardiac output but not stroke volume is similar in a wingate and vO<sub>2peak</sub> test in young men. *European Journal of Applied Physiology*, **111**, 155-158.
- González-Alonso, J. (2008) Point:Counterpoint: Stroke volume does/does not decline during exercise at maximal effort in healthy individuals. *Journal of Applied Physiology* **104**, 275-276.
- Hopkins, W.G., Marshall, S.W., Batterham, A.M. and Hanin, J. (2009) Progressive statistics for studies in sports medicine and exercise science. *Medicine and Science in Sports and Exercise* **41**, 3-13.
- Lepretre, P. M., Koralsztejn, J. P. & Billat, V. L. (2004). Effect of exercise intensity on relationship between VO<sub>2max</sub> and cardiac output. *Medicine and Science in Sports and Exercise* **36**, 1357-1363.
- Medbø, J. and Tabata, I. (1989) Relative importance of aerobic and anaerobic energy release during short-lasting exhausting bicycle exercise. *Journal of Applied Physiology* **67**, 1881-1886.
- Mortensen, S.P., Dawson, E.A., Yoshiga, C.C., Dalsgaard, M.K., Damsgaard, R., Secher, N.H. and Gonzalez-Alonso, J. (2005). Limitations to systemic and locomotor limb muscle oxygen delivery and uptake during maximal exercise in humans. *Journal of Physiology-London* **566**, 273-285.
- Richard, R., Lonsdorfer-Wolf, E., Dufour, S., Doutreleau, S., Oswald-



Mammosser, M., Billat, V. and Lonsdorfer, J. (2004) Cardiac output and oxygen release during very high-intensity exercise performed until exhaustion. *European Journal of Applied Physiology* **93**, 9-18.

Stanley, J., Peake, J., Coombes, J. and Buchheit, M. (2014) Central and peripheral adjustments during high-intensity exercise following cold water immersion. *European Journal of Applied Physiology* **114**(1), 147-163.

Warburton, D.E.R. and Gledhill, N. (2008) Counterpoint: Stroke volume does not decline during exercise at maximal effort in healthy individuals. *Journal of Applied Physiology* **104**, 276-278.

### Key points

- Moderate-intensity recovery periods may not be necessary to maintain high stroke volume during the exercise intervals of HIIT.
- Stroke volume did not surpass the levels attained during the exercise intervals during the recovery periods of HIIT.
- The practical implication of these finding is that reducing the intensity of the recovery period during a HIIT protocol may prolong the time to exhaustion, potentially allowing completion of additional high-intensity intervals increasing the time accumulated at maximal cardiac output.

### AUTHORS BIOGRAPHY



#### **Jamie STANLEY**

##### **Employment**

Post-Doctoral Research Fellow, School of Human Movement Studies, The University of Queensland and Centre of Excellence for Applied Sport Science Research, Queensland Academy of Sport, Brisbane, Australia. Research Scientist, Queensland Academy of Sport, Triathlon, Brisbane, Australia

##### **Degree**

PhD

##### **Research interest**

Exercise physiology, performance, recovery monitoring

**E-mail:** [j.stanley@uq.edu.au](mailto:j.stanley@uq.edu.au)



#### **Martin BUCHHEIT**

##### **Employment**

Physiologist & Sport Scientist, Myorobie Association, Montvalezan, France

##### **Degree**

PhD

##### **Research interest**

Exercise physiology, performance monitoring, progressive statistics, football, soccer, handball.

**E-mail:** [mb@martin-buchheit.net](mailto:mb@martin-buchheit.net)

#### ✉ **Jamie Stanley**

School of Human Movement Studies, The University of Queensland, Brisbane, Queensland 4072, Australia