

## **THE LIMITING BEHAVIOUR OF HANSKI'S INCIDENCE FUNCTION METAPOPOPULATION MODEL**

R. MCVINISH,\* *University of Queensland*

P.K. POLLETT,\*\* *University of Queensland*

### **Abstract**

Hanski's incidence function model is one of the most widely used metapopulation models in ecology. It models the presence/absence of a species at spatially distinct habitat patches as a discrete-time Markov chain whose transition probabilities are determined by the physical landscape. In this analysis, the limiting behaviour of the model is studied as the number of patches increases and the size of the patches decreases. Two different limiting cases are identified depending on whether or not the metapopulation is initially near extinction. Basic properties of the limiting models are derived.

*Keywords:* Extinction threshold; point process; stochastic patch occupancy model; SPOM; weak convergence

2000 Mathematics Subject Classification: Primary 92D40

Secondary 60J10;60J20;60F05

### **1. Introduction**

A metapopulation is a collection of local populations occupying spatially distinct habitat patches that evolves over time due to local extinctions and colonisation events. As noted in [12], the study of metapopulations using stochastic patch occupancy models (SPOM) is well established in the ecology literature. A SPOM is a discrete-time Markov chain that models the presence/absence of the focal species in a network of habitat patches. One of the most

---

\* Postal address: School of Mathematics and Physics, University of Queensland, Brisbane, QLD 4072, Australia.

\* Email address: r.mcvinish@uq.edu.au

\*\* Email address: pkp@maths.uq.edu.au

widely used SPOMs is Hanski's incidence function model (IFM) [10]. The IFM uses structural assumptions to relate the physical landscape to the transition probabilities of the Markov chain.

One drawback of using the IFM and realistic SPOMs in general is that they are difficult to analyse. Although simulation methods can be employed in specific cases, this approach provides no information on the behaviour of the model in general. To understand the general case, the modeller typically needs to resort to approximations. Two common classes of approximations for Markov chain models are deterministic approximations and branching process approximations.

Ovaskainen and Hanski [24] propose a system of deterministic difference equations as an approximation to the IFM and provide a detailed analysis of this system. However, it is not clear in what sense this system of deterministic difference equations approximates the original stochastic IFM. Our analysis is perhaps closer to Ovaskainen and Cornell's [23] analysis of a related continuous-time stochastic metapopulation model on a stochastic landscape. They showed that the fraction of occupied patches is well approximated by a one-dimensional ordinary differential equation called Levins' model [16] when the dispersal range is large.

If the spatial aspect of the metapopulation is ignored, then the behaviour of the metapopulation model near the extinction state resembles the initial behaviour of an epidemic. A number of researchers ([1, 2, 3, 4] among many others) have used branching processes to approximate the initial spread of an epidemic in a large population. The branching process approximation is less frequently applied in the context of metapopulation modelling, but [5] provides one example.

In this paper, we study the limiting behaviour of Hanski's IFM as the number of patches increases and the size of the patches decreases. The model is briefly described and the main assumptions stated in Section 2. In Section 3, we derive and study the deterministic limit. Our analysis differs from that of [23] in three ways: (i) a different scaling is used to study the limiting process, (ii) we allow the extinction probabilities to vary spatially, and (iii) we work directly with the discrete-time IFM. In Section 4, Hanski's IFM is studied when the metapopulation is initially close to extinction. Extending our earlier work [18], point process theory is used to study the limit and determine conditions under which the metapopulation has a positive probability of recovery. We conclude with a brief discussion of the results. All proofs are given in the Appendix.

## 2. Model description

The incidence function model of Hanski [10] for a metapopulation comprising  $n$  patches is a discrete-time Markov chain on  $\{0, 1\}^n$ . Denote this Markov chain by  $X_t^n = (X_{1,t}^n, \dots, X_{n,t}^n)$ , where  $X_{i,t}^n = 1$  if patch  $i$  is occupied at time  $t$  and  $X_{i,t}^n = 0$  otherwise. Patch  $i$  is described by three variables; its location  $z_i$ , a weight  $A_i > 0$  which may be interpreted as the size of the patch, and  $s_i$  which is the probability that the population occupying patch  $i$  survives one time step. Let  $z^n$ ,  $A^n$  and  $s^n$  denote the collections of patch variables for the entire metapopulation. We treat  $z^n$ ,  $A^n$  and  $s^n$  as random variables which is reasonable if we view the landscape as the result of some random process. A similar treatment of the landscape was adopted in the metapopulation model of Ovaskainen and Cornell [23]. The transition probabilities of the Markov chain are determined by how well the patches are connected to each other and by the probability of local extinction. The connectivity measure of patch  $i$  at time  $t$  is defined by

$$\tilde{C}_i(t) = \sum_{j \neq i} X_{j,t}^n D(z_i, z_j) A_j^b, \quad b > 0,$$

where  $D(z, \tilde{z}) \geq 0$  is a measure of how easily a patch located at  $z$  can be colonised by an individual from a patch located at  $\tilde{z}$ . Typically,  $D(z, \tilde{z})$  is a bounded decreasing function of  $\|z - \tilde{z}\|$  such as  $\exp(-\alpha\|z - \tilde{z}\|)$ ,  $\alpha > 0$ . Let  $f : [0, \infty) \mapsto [0, 1]$  such that  $f(0) = 0$ . Conditional on  $(X_t^n, z^n, A^n, s^n)$ , the  $X_{i,t+1}^n$  ( $i = 1, \dots, n$ ) are independent with transitions given by

$$P(X_{i,t+1}^n = 1 \mid X_t^n, z^n, A^n, s^n) = s_i X_{i,t}^n + f(\tilde{C}_i(t))(1 - X_{i,t}^n).$$

Since the connectivity measure of patch  $i$  is only important when patch  $i$  is unoccupied, the transition probabilities remain unchanged if  $\tilde{C}_i(t)$  is replaced by

$$C_i(t) = \sum_{j=1}^n X_{j,t}^n D(z_i, z_j) A_j^b.$$

There are a couple of important differences between the IFM and typical interacting particle systems [9, 25] used in spatial modelling. Apart from the discrete time setting, interacting particle systems usually evolve on a lattice of sites whereas the patches of an IFM are not naturally restricted to a lattice structure. More importantly, in interacting particle systems the instantaneous effect of change at one site is assumed to be limited to a finite neighbourhood of that site. The IFM does not usually have a finite range neighbourhood. For typical choices of  $D$ , changes at one patch immediately affect the transition probabilities for all other patches of the metapopulation.

Our analysis of the IFM will be based on the following assumptions:

- (A)  $A_i = a_i n^{-1/b}$  where  $a_i \in (0, A]$ . The collection of  $a_i$  will be denoted by  $a^n$ .
- (B)  $z_i \in \Omega$  where  $\Omega$  is a compact subset of  $\mathbb{R}^d$ .
- (C)  $D(z, \bar{z})$  is symmetric in its arguments and defines a uniformly bounded and equicontinuous family of functions on  $\Omega$ . That is, there exists a finite constant  $\bar{D}$  such that for all  $z_1, z_2 \in \Omega$ ,  $|D(z_1, z_2)| \leq \bar{D}$ , and for every  $\epsilon > 0$  there exists a  $\delta > 0$  such that for all  $z_1, z_2$  with  $\|z_1 - z_2\| < \delta$

$$\sup_{z \in \Omega} |D(z_1, z) - D(z_2, z)| < \epsilon.$$

- (D) The function  $f$  is Lipschitz continuous.

Assumption (A) implies that all patches are of a comparable size. Furthermore, in applications to butterfly metapopulations, estimates of  $b$  are typically less than one [11, 26]. Therefore, Assumption (A) corresponds to a decrease in the total area of the metapopulation as  $n \rightarrow \infty$  which is consistent with habitat fragmentation and destruction. Assumption (B) is made to simplify the analysis. Assumption (C) is satisfied for the examples given in [21, section 2.2.1]. In particular, it is satisfied for  $D(z, \bar{z}) = \exp(-\alpha\|z - \bar{z}\|)$ ,  $\alpha \geq 0$ . Typical forms for  $f$  include  $f(x) = x^2/(\beta + x^2)$ ,  $\beta > 0$ , and  $f(x) = 1 - \exp(-\beta x)$ ,  $\beta > 0$ , [21, section 2.2.3]. Both of these functions have bounded derivatives on  $[0, \infty)$ , so Assumption (D) is satisfied.

Another assumption that will be used in the analysis concerns the random variables  $(z^n, a^n, s^n)$ . Let  $C^+([0, 1] \times \Omega)$  denote the class of continuous functions  $h : [0, 1] \times \Omega \mapsto [0, \infty)$ . Note that since  $\Omega$  is a compact set from Assumption (B), the functions in  $C^+([0, 1] \times \Omega)$  are bounded. Consider the sequence of random measures  $\sigma_n$  defined by

$$\int h(s, z) \sigma_n(ds, dz) := n^{-1} \sum_{i=1}^n a_i^b h(s_i, z_i), \quad \text{for all } h \in C^+([0, 1] \times \Omega).$$

The measure  $\sigma_n$  describes the landscape of the  $n$  patch metapopulation model. It is purely atomic placing mass  $n^{-1}a_i^b$  at the point determined by patch  $i$ 's location  $z_i$  and its survival probability  $s_i$ . We assume that  $\sigma_n$  satisfies the following:

- (E) As  $n \rightarrow \infty$ ,  $\sigma_n \xrightarrow{d} \sigma$  for some non-random measure  $\sigma$ .

Applying the law of large numbers and Theorem 16.16 of [15], we see Assumption (E) is satisfied if, for example, the random vectors  $(z_i, a_i, s_i)$  are independent and identically distributed.

### 3. Deterministic limit

Consider the array of random measures  $\mu_{n,t}$  constructed from the Markov chain  $X_t^n$  by

$$\int h(s, z) \mu_{n,t}(ds, dz) := n^{-1} \sum_{i=1}^n a_i^b X_{i,t}^n h(s_i, z_i), \quad \text{for all } h \in C^+([0, 1] \times \Omega). \quad (3.1)$$

The measure  $\mu_{n,t}$  has a similar structure to  $\sigma_n$ , however,  $\mu_{n,t}$  only involves those patches that are occupied at time  $t$ . These measures can be used to determine quantities such as the proportion of occupied patches weighted by the patch size. The following theorem describes the behaviour of the metapopulation as the number of patches tends to infinity.

**Theorem 3.1.** *Suppose that Assumptions (A) – (E) hold and that  $\mu_{n,0} \xrightarrow{d} \mu_0$  for some non-random measure  $\mu_0$ . Then  $\mu_{n,t} \xrightarrow{d} \mu_t$  for all  $t = 0, 1, \dots$ , where  $\mu_t$  is defined by the recursion*

$$\begin{aligned} & \int h(s, z) \mu_{t+1}(ds, dz) \\ &= \int sh(s, z) \mu_t(ds, dz) + \int h(s, z) f \left( \int D(z, \bar{z}) \mu_t(d\bar{s}, d\bar{z}) \right) \sigma(ds, dz) \\ & \quad - \int h(s, z) f \left( \int D(z, \bar{z}) \mu_t(d\bar{s}, d\bar{z}) \right) \mu_t(ds, dz), \end{aligned} \quad (3.2)$$

for all  $h \in C^+([0, 1] \times \Omega)$ .

Consider the special case where  $D(z, \bar{z}) = 1$  for all  $(z, \bar{z}) \in \Omega \times \Omega$  and assuming  $a_i = 1$  for all  $i$ . Let  $h(s, z) = s^k$  for  $k = 0, 1, 2, \dots$ . Then recursion (3.2) becomes

$$d(t+1; k) = d(t; k+1) + f(d(t; 0)) (\bar{s}_k - d(t; k)). \quad (3.3)$$

where  $d(t; k) = \int s^k \mu_t(ds, dz)$  and  $\bar{s}_k = \int s^k \sigma(ds, dz)$ . This special case is related to the model studied in [18]. The difference between equation (3.3) and equation (2.1) in [18] is due to [18] assuming that extinction and colonisation events occur in distinct alternating phases.

Despite the complexity of the recursion (3.2), we are able to provide conditions for the existence of a non-zero fixed point and show that if a non-zero fixed point exists then it is unique. Fixed points are important in applications as they determine the equilibrium behaviour of the metapopulation. In particular, the presence of a non-zero fixed point implies that the metapopulation may persist for a long time.

Three further assumptions are introduced to study the fixed points of the recursion (3.2):

- (F) The function  $f$  is increasing, strictly concave and twice differentiable with bounded second derivative in a neighbourhood of 0.

(G) For some  $\epsilon > 0$  sufficiently small,  $\sigma([1 - \epsilon, 1] \times \Omega) = 0$ . Also, for every  $z \in \Omega$  and every open neighbourhood  $N_z$  of  $z$ ,  $\sigma([0, 1] \times N_z) > 0$ .

(H)  $D(z, \tilde{z}) > 0$  for all  $(z, \tilde{z}) \in \Omega \times \Omega$ .

Assumption (F) is satisfied for  $f(x) = 1 - \exp(-\beta x)$ ,  $\beta > 0$ , but not for  $f(x) = x^2/(\beta + x^2)$ ,  $\beta > 0$ . In ecological terms, by introducing Assumption (F) we are trying to exclude a metapopulation level Allee-like effect. An Allee-like effect refers to a metapopulation exhibiting a critical threshold in the occupancy level below which the metapopulation goes extinct [6, pages 103-105]. Note that even for simple models displaying an Allee-like effect, determining the number of fixed points can be a challenging problem. Assumption (G) excludes the possibility of any patches having a very large survival probability. Furthermore, for any  $\mu_{n,t}$  defined by equation (3.1) such that  $\mu_{n,t} \xrightarrow{d} \mu_t$ , Assumption (G) implies that  $\mu_t([1 - \epsilon, 1] \times \Omega) = 0$ . Assumption (H) means that all patches are connected to each other. Although some range limited forms of  $D$  have been used, the most common forms [21, section 2.2.1] satisfy Assumption (H).

Let  $\mu_\infty(ds, dz)$  denote a fixed point of (3.2) and defined  $\psi(z) = \int D(z, \tilde{z})\mu_\infty(d\tilde{s}, d\tilde{z})$ . Upon rearranging (3.2), we see that any fixed point satisfies

$$\int h(s, z)(1 - s + f(\psi(z)))\mu_\infty(ds, dz) = \int h(s, z)f(\psi(z))\sigma(ds, dz). \quad (3.4)$$

By definition  $\psi(z) \geq 0$  and, by Assumption (C),  $\psi$  is continuous on  $\Omega$ . Under Assumption (G) the support of  $\sigma$ , hence the support of  $\mu_\infty$ , is contained in  $[0, 1 - \epsilon] \times \Omega$ . Therefore, for any  $h \in C^+([0, 1] \times \Omega)$ , the function  $h(s, z)(1 - s + f(\psi(z)))^{-1}$  is a continuous function on the support of  $\mu_\infty$ . Using this fact together with equation (3.4), we see that for any  $h \in C^+([0, 1] \times \Omega)$ ,

$$\int h(s, z)\mu_\infty(ds, dz) = \int h(s, z)\frac{f(\psi(z))}{1 - s + f(\psi(z))}\sigma(ds, dz).$$

Clearly,  $\mu_\infty$  is absolutely continuous with respect to  $\sigma$  and has Radon-Nikodym derivative

$$\frac{\partial\mu_\infty}{\partial\sigma} = \frac{f(\psi(z))}{1 - s + f(\psi(z))},$$

where  $\psi$  satisfies the equation

$$\psi(z) = \int D(z, \tilde{z})\frac{f(\psi(\tilde{z}))}{1 - \tilde{s} + f(\psi(\tilde{z}))}\sigma(d\tilde{s}, d\tilde{z}). \quad (3.5)$$

The number of solutions to equation (3.5) is determined by the following theorem.

**Theorem 3.2.** *Suppose that Assumptions (B) – (D) and (F) – (H) hold. Let  $\mathcal{A} : C(\Omega) \mapsto C(\Omega)$  be the bounded linear operator*

$$\mathcal{A}\phi(z) = f'(0) \int \frac{D(z, \tilde{z})}{(1 - \tilde{s})} \phi(\tilde{z}) \sigma(d\tilde{s}, d\tilde{z}), \quad \phi \in C(\Omega), \quad (3.6)$$

and let  $r(\mathcal{A})$  be the spectral radius of  $\mathcal{A}$ . If  $r(\mathcal{A}) \leq 1$  then  $\psi = 0$  is the unique solution to equation (3.5). If  $r(\mathcal{A}) > 1$  then equation (3.5) has two solutions of which one is  $\psi = 0$ .

Theorem 3.2 simplifies considerably for certain special cases. As one example, consider the case where  $D(z, \tilde{z}) = \exp(-\theta(\|z - z_0\| + \|\tilde{z} - z_0\|))$  for some  $\theta > 0$  and  $z_0 \in \Omega$ . This form for  $D(z, \tilde{z})$  can be interpreted as requiring that individuals moving from a patch located at  $\tilde{z}$  to a patch located at  $z$  to first go through the point  $z_0$ . To apply Theorem 3.2, we note that  $\mathcal{A}$  maps continuous functions to functions of the form  $C \exp(-\theta\|z - z_0\|)$  for some  $C > 0$ . Therefore, the spectral radius of  $\mathcal{A}$  is easily computed to be

$$f'(0) \int \frac{\exp(-2\theta\|\tilde{z} - z_0\|)}{1 - \tilde{s}} \sigma(d\tilde{s}, d\tilde{z}).$$

Further discussion of metapopulations with  $D$  of this form is given in [20].

To end this section we comment on the problem of stability of the fixed points. Although we have been unable to determine the stability of these fixed points, we conjecture that if  $r(\mathcal{A}) > 1$  then the non-zero fixed point is asymptotically stable and the zero fixed point is unstable. Otherwise, if  $r(\mathcal{A}) \leq 1$  then the zero fixed point is asymptotically stable. In other words, we conjecture that this metapopulation will persist for a long time if  $r(\mathcal{A}) > 1$  but will go extinct quickly if  $r(\mathcal{A}) \leq 1$ , a property called the extinction threshold. In support of this conjecture, note that if the extinction and colonisation events were assumed to occur in distinct alternating phases as in [18, 20], then stability of the fixed points could be established using similar arguments to those used to prove Theorem 3 in [20].

#### 4. Point process limit

We now consider the case where initially only a small number of habitat patches are occupied. Although we could still apply Theorem 3.1, we would not obtain any useful information about the metapopulation. In that case,  $\mu_{n,0}$  converges to the trivial measure which implies that  $\mu_t$  is the trivial measure for all  $t \geq 0$ . A different analysis is needed to understand the behaviour of the metapopulation model near the extinction state. We replace Assumption (A) by

(A')  $A_i = n^{-1/b}$  for all  $i$ .

Assumption (A') implies that all habitat patches are of the same size. Note that, from Assumption (C),  $D(z, \tilde{z})$  can be continuously extended to all  $\mathbb{R}^d$  so that  $D(\cdot, \tilde{z}) = 0$  for all  $\tilde{z} \notin \Omega'$  where  $\Omega'$  is a compact subset of  $\mathbb{R}^d$  and  $\Omega \subset \Omega'$ .

As in [18], we define the point processes associated with the IFM using a random counting measure. Let  $\mu_{n,t}$  denote the random measure defined by

$$\mu_{n,t}(B) := \# \left\{ (s_i, z_i) \in B : X_{i,t}^n = 1 \right\}, \quad (4.1)$$

for any bounded Borel set  $B$ . Note that the definition of  $\mu_{n,t}$  given at (4.1) has two important differences to the one given at (3.1); all patches are of the same size and there is no longer a scaling by  $n^{-1}$ . The main tool that we shall use in the analysis of these point processes is the probability generating functional. Let  $\mathcal{V}$  denote the class of all real-valued Borel functions  $h$  on  $\mathbb{R}^{d+1}$  with  $1 - h$  vanishing outside some bounded set and satisfying  $0 \leq h(s, z) \leq 1$ , for all  $(s, z) \in \mathbb{R}^{d+1}$ . The probability generating functional of the point process  $\mu_{n,t}$  is defined by

$$G_{n,t}[h] := \mathbb{E} \left( \exp \left( \int \log h(s, z) \mu_{n,t}(ds, dz) \right) \right) = \mathbb{E} \left( \prod_{i=1}^n (X_{i,t}^n h(s_i, z_i) + 1 - X_{i,t}^n) \right).$$

Moyal [22] introduced a generalisation of branching processes called multiplicative population chains. Multiplicative population chains (MPC) are a family of point processes that evolve in discrete time such that the point process at time  $t + 1$  is a superposition of conditionally independent point processes representing the offspring from each element of the point process at time  $t$ . Let  $G_t[\cdot | (s, z)]$  denote the probability generating functional of the MPC at time  $t$  given the MPC at time 0 consisted of a single element locate at  $(s, z)$ . Moyal [22, Theorem 1.1] showed that the probability generating functional of an MPC satisfies the functional relation

$$G_t[h | (s, z)] = G_u[G_{t-u}[h | \cdot] | (s, z)], \quad u = 1, \dots, t - 1. \quad (4.2)$$

Given the probability generating functional at time 0, denoted by  $G_0$ , the probability generating functional at time  $t$  can be obtained from the recursion (4.2) by  $G_t[h] = G_0[G_t[h | (s, z)]]$ . In the following theorem we show that, under certain assumptions, the IFM converges to an MPC and give the recursion for the probability generating functional.

**Theorem 4.1.** *Suppose that Assumptions (A'), (B) – (F) hold. Assume that the sequence of point processes  $\{\mu_{n,0}\}_{n=1}^{\infty}$  converges weakly to some point process with probability generating*



functional  $G_0$  and that  $\sup_n \mathbb{E} \left( \exp \left( \alpha \sum_{i=1}^n X_{i,0}^n \right) \right) < \infty$ , for all  $\alpha > 0$ . The sequence of point processes  $\{\mu_{n,t}\}_{n=1}^\infty$  converges weakly to the point process with probability generating functional given by the recursion

$$G_{t+1}[h] = G_t[G_1[h | (s, z)]], \quad \text{for any } h \in \mathcal{V}, \quad (4.3)$$

where

$$G_1[h | (s, z)] = (1 - U(s)(1 - h(s, z))) \exp \left( -f'(0) \int D(\bar{z}, z) (1 - h(\bar{s}, \bar{z})) \sigma(d\bar{s}, d\bar{z}) \right) \quad (4.4)$$

and  $U$  is the distribution function of the standard uniform distribution.

As is the case with branching process approximations of Markov chains, of particular interest is the probability that the limiting MPC goes extinct in finite time. Moyal [22, Theorem 3.1] shows that to determine the extinction probability one needs to find the smallest nonnegative solution to the functional equation  $h = G_1[h | (s, z)]$ ,  $h \in \mathcal{V}$ . Clearly, one solution is given by  $h(s, z) = 1$  for all  $(s, z) \in \mathbb{R}^{d+1}$ . Therefore, we need to determine conditions under which a smaller solutions exist.

**Theorem 4.2.** *Suppose that Assumptions (B), (C), (G) and (H) hold. The probability that the limiting MPC goes extinct in finite time is less than one if and only if  $r(\mathcal{A}) > 1$ , where  $\mathcal{A}$  is the bounded linear operator defined by equation (3.6). If  $r(\mathcal{A}) > 1$  then the probability of extinction in finite time is given by*

$$G_0 \left[ \frac{(1 - U(s))\psi^*(z)}{1 - U(s)\psi^*(z)} \right],$$

where  $\psi^*$  is the smallest nonnegative solution to

$$\psi(z) = \exp \left( -f'(0) \int D(\bar{z}, z) \left( \frac{1 - \psi(\bar{z})}{1 - U(\bar{s})\psi(\bar{z})} \right) \sigma(d\bar{s}, d\bar{z}) \right), \quad \text{for all } z \in \Omega.$$

When  $D(z, \bar{z}) = 1$  for all  $(z, \bar{z}) \in \Omega \times \Omega$ , the results of this section parallel those of Section 3 in [18] with the small differences being due to the phase structure of colonisation and extinction events assumed in [18]. The proof of Theorem 4.2 (Appendix A.4) uses similar functional analytic tools to those used in the proof of Theorem 3.2 (Appendix A.2).

Theorem 4.2 provides further support to our conjecture concerning the stability of the fixed points of (3.2). Suppose that  $r(\mathcal{A}) > 1$ , then we conjectured that zero is an unstable fixed point of the recursion (3.2). If this were not the case then the metapopulation would display

very strange behaviour. Theorem 4.2 indicates that if the initial number of occupied patches is very small compared to the number of patches in the metapopulation and  $r(\mathcal{A}) > 1$ , then the number of occupied patches will increase and the metapopulation will persist for a long time with positive probability. However, if zero were a stable fixed point of recursion (3.2) and a small but positive proportion of patches are initially occupied in the metapopulation then there should be a rapid decrease to zero in the proportion of occupied patches.

## 5. Discussion

In this paper, we studied the convergence of the IFM under two different sets of initial conditions. One of the goals of studying convergence was to derive an approximation of the IFM which would allow a simpler analysis. Although both of the derived limiting processes are still quite complex, we have been able to establish some of their features. One benefit of determining the limiting processes is that the effect of the model parameters is more transparent. For example, it can be demonstrated that those factors that we would expect to improve the likelihood of the metapopulation persisting, such as a more rapidly increasing colonisation function and easier movement between patches, lead to an increase in the non-zero equilibrium of the deterministic limit from Theorem 3.1 and a decrease in the probability of extinction in the MPC limit from Theorem 4.1.

Determining the stability of the fixed points of (3.2) remains a significant challenge. We have conjectured that the non-zero fixed point of (3.2), when it exists, is stable and that when only the zero fixed point exists, the zero fixed point is stable. A resolution of this problem is necessary before more complex dynamics, such as a metapopulation Allee-like effect, can be incorporated.

## Appendix A. Proofs

### A.1. Proof of Theorem 3.1

We essentially proceed as in the proof of Theorem 2.1 of [18]. First note that if  $\int h d\mu_n \xrightarrow{d} \int h d\mu$  for all  $h \in C^+([0, 1] \times \Omega)$  then  $\mu_n \xrightarrow{d} \mu$  [15, Theorem 16.16]. We use induction on  $t$  to prove weak convergence of the random measures  $\mu_{n,t}$  to non-random measures  $\mu_t$ . By assumption  $\mu_{n,0} \xrightarrow{d} \mu_0$  for some non-random measure  $\mu_0$ . Suppose that  $\mu_{n,t} \xrightarrow{d} \mu_t$  for some

non-random measure  $\mu_t$ . Then

$$\begin{aligned}
\mathbb{E}\left(\int h d\mu_{n,t+1}|X_t^n, s^n, a^n, z^n\right) &= n^{-1} \sum_{i=1}^n a_i^b h(s_i, z_i) \mathbb{E}\left(X_{i,t+1}^n | X_t^n, s^n, a^n, z^n\right) \\
&= n^{-1} \sum_{i=1}^n a_i^b s_i h(s_i, z_i) X_{i,t}^n + n^{-1} \sum_{i=1}^n a_i^b h(s_i, z_i) f\left(n^{-1} \sum_{j=1}^n a_j^b D(z_i, z_j) X_{j,t}^n\right) (1 - X_{i,t}^n) \\
&= \int sh(s, z) \mu_{n,t}(ds, dz) + \int h(s, z) f\left(\int D(z, \tilde{z}) \mu_{n,t}(d\tilde{s}, d\tilde{z})\right) \sigma_n(ds, dz) \\
&\quad - \int h(s, z) f\left(\int D(z, \tilde{z}) \mu_{n,t}(d\tilde{s}, d\tilde{z})\right) \mu_{n,t}(ds, dz) \\
&= \int sh(s, z) \mu_{n,t}(ds, dz) + \int h(s, z) f\left(\int D(z, \tilde{z}) \mu_t(d\tilde{s}, d\tilde{z})\right) \sigma_n(ds, dz) \\
&\quad - \int h(s, z) f\left(\int D(z, \tilde{z}) \mu_t(d\tilde{s}, d\tilde{z})\right) \mu_{n,t}(ds, dz) + \epsilon_{n,t}(h),
\end{aligned}$$

where

$$\begin{aligned}
\epsilon_{n,t}(h) &= \int h(s, z) \left\{ f\left(\int D(z, \tilde{z}) \mu_{n,t}(d\tilde{s}, d\tilde{z})\right) - f\left(\int D(z, \tilde{z}) \mu_t(d\tilde{s}, d\tilde{z})\right) \right\} \sigma_n(ds, dz) \\
&\quad - \int h(s, z) \left\{ f\left(\int D(z, \tilde{z}) \mu_{n,t}(d\tilde{s}, d\tilde{z})\right) - f\left(\int D(z, \tilde{z}) \mu_t(d\tilde{s}, d\tilde{z})\right) \right\} \mu_{n,t}(ds, dz).
\end{aligned}$$

By Assumption (D)  $f$  is Lipschitz continuous, so there exists an  $L > 0$  such that

$$\begin{aligned}
|\epsilon_{n,t}(h)| &\leq L \int h(s, z) \left| \int D(z, \tilde{z}) \mu_{n,t}(d\tilde{s}, d\tilde{z}) - \int D(z, \tilde{z}) \mu_t(d\tilde{s}, d\tilde{z}) \right| (\sigma_n(ds, dz) + \mu_{n,t}(ds, dz)) \\
&\leq 2L \left( \int h(s, z) \sigma_n(ds, dz) \right) \sup_{z \in \Omega} \left| \int D(z, \tilde{z}) \mu_{n,t}(d\tilde{s}, d\tilde{z}) - \int D(z, \tilde{z}) \mu_t(d\tilde{s}, d\tilde{z}) \right|.
\end{aligned}$$

From a small modification of Theorem 3.1 of [27] and Assumption (C), it follows that if  $\mu_{n,t} \xrightarrow{d} \mu_t$ , a non-random measure, then

$$\sup_{z \in \Omega} \left| \int D(z, \tilde{z}) \mu_{n,t}(d\tilde{s}, d\tilde{z}) - \int D(z, \tilde{z}) \mu_t(d\tilde{s}, d\tilde{z}) \right| \xrightarrow{p} 0.$$

As  $h \in C^+([0, 1] \times \Omega)$  then both  $sh(s, z)$  and  $h(s, z) f\left(\int D(z, \tilde{z}) \mu_t(d\tilde{s}, d\tilde{z})\right)$  are in  $C^+([0, 1] \times \Omega)$ , the later being a consequence of Assumption (C). This implies

$$\begin{aligned}
\mathbb{E}\left(\int h d\mu_{n,t+1}|X_t^n, s^n, a^n, z^n\right) &\xrightarrow{p} \int sh(s, z) \mu_t(ds, dz) + \int h(s, z) f\left(\int D(z, \tilde{z}) \mu_t(d\tilde{s}, d\tilde{z})\right) \sigma(ds, dz) \\
&\quad - \int h(s, z) f\left(\int D(z, \tilde{z}) \mu_t(d\tilde{s}, d\tilde{z})\right) \mu_t(ds, dz).
\end{aligned}$$

Now to compute the conditional variance of  $\int h(s, z) \mu_{n,t+1}(ds, dz)$  given  $X_t^n$ :

$$\begin{aligned} \text{var} \left( \int h d\mu_{n,t+1} | X_t^n, s^n, a^n, z^n \right) &= \text{var} \left( n^{-1} \sum_{i=1}^n a_i^b X_{i,t+1}^n h(s_i, z_i) | X_t^n, s^n, a^n, z^n \right) \\ &= n^{-2} \sum_{i=1}^n a_i^{2b} h^2(s_i, z_i) \left( s_i(1-s_i)X_{i,t}^n + f(C_i(t))(1-f(C_i(t)))(1-X_{i,t}^n) \right) \leq n^{-2} \sum_{i=1}^n a_i^{2b} h^2(s_i, z_i). \end{aligned}$$

From Assumptions (A) and (B), the conditional variance goes to zero in probability. Applying a Chebyshev type inequality [19, Appendix C], we see  $\int h d\mu_{n,t+1}$  converges to  $\int h d\mu_{t+1}$  in probability which is equivalent to convergence in distribution as  $\int h d\mu_{t+1}$  is non-random.

## A.2. Proof of Theorem 3.2

The proof of Theorem 3.2 depends on the theory concerning positive operators and monotone operators. We recall some definitions and notation following [8, Chapter VIII Appendix].

Let  $C(\Omega)$  be the Banach space of continuous functions on  $\Omega$  (recall  $\Omega$  is compact by Assumption (B)). Let  $K$  denote the reproducing cone of functions  $h$  on  $\Omega$  such that  $h(z) \geq 0$  for all  $z \in \Omega$  and let  $\mathring{K}$  denote the interior of  $K$ . For any  $\phi, \chi \in K$  we write  $\phi \leq \chi$  if  $\chi - \phi \in K$ .

We have previously noted that the function  $\psi$  defined by  $\psi(z) = \int D(z, \bar{z}) \mu_\infty(d\bar{z})$  is in  $K$ . Let  $R : K \rightarrow K$  be the operator defined by the right-hand side of equation (3.5). The following properties of  $R$  will be used to prove Theorem 3.2.

**Proposition A.1.** *Suppose that Assumptions (B)–(D) and (F)–(H) hold. The operator  $R$  has the following properties:*

- (i) *continuity.*
- (ii) *order compactness, that is, for any  $\chi_1, \chi_2 \in K$ ,  $R$  maps the set  $\{\phi : \phi \leq \chi_2, \chi_1 \leq \phi\}$  to a relatively compact set.*
- (iii) *monotonicity, that is, if  $\phi_1 \leq \phi_2$  then  $R\phi_1 \leq R\phi_2$ .*
- (iv) *strong positivity, that is, if  $\phi \in K \setminus \{0\}$  then  $R\phi \in \mathring{K}$ .*
- (v) *strong sublinearity, that is, if  $\lambda \in (0, 1)$  and  $\phi \in \mathring{K}$  then  $R(\lambda\phi) - \lambda R\phi \in \mathring{K}$ .*

*Proof.* (i) *Continuity:* We wish to show that for any  $\epsilon > 0$  there exists a  $\delta$  such that for any  $\phi_1, \phi_2 \in K$ ,

$$\sup_{z \in \Omega} |\phi_1(z) - \phi_2(z)| \leq \delta \quad \implies \quad \sup_{z \in \Omega} |R\phi_1(z) - R\phi_2(z)| \leq \epsilon.$$

Now,

$$\begin{aligned}
|R\phi_1(z) - R\phi_2(z)| &= \left| \int D(z, \bar{z}) \left[ \frac{f(\phi_1(\bar{z}))}{1 - \bar{s} + f(\phi_1(\bar{z}))} - \frac{f(\phi_2(\bar{z}))}{1 - \bar{s} + f(\phi_2(\bar{z}))} \right] \sigma(d\bar{s}, d\bar{z}) \right| \\
&\leq \int D(z, \bar{z}) \frac{(1 - \bar{s}) |f(\phi_1(\bar{z})) - f(\phi_2(\bar{z}))|}{(1 - \bar{s} + f(\phi_1(\bar{z}))) (1 - \bar{s} + f(\phi_2(\bar{z})))} \sigma(d\bar{s}, d\bar{z}) \\
&\leq L \int D(z, \bar{z}) (1 - \bar{s})^{-1} |\phi_1(\bar{z}) - \phi_2(\bar{z})| \sigma(d\bar{s}, d\bar{z})
\end{aligned}$$

since  $f$  is Lipschitz continuous from Assumption (D). If  $\sup_{z \in \Omega} |\phi_1(z) - \phi_2(z)| \leq \delta$ , then  $|R\phi_1(z) - R\phi_2(z)| \leq L\delta\bar{D} \int (1 - \bar{s})^{-1} \sigma(d\bar{s}, d\bar{z})$  by Assumption (C). By Assumption (G), this integral is finite. Setting  $\delta = \epsilon(L\bar{D} \int (1 - \bar{s})^{-1} \sigma(d\bar{s}, d\bar{z}))^{-1}$ , continuity of  $R$  is established.

(ii) *Order compactness*: A subset  $Y$  of a metric space  $X$  is relatively compact if any bounded sequence in  $Y$  has a subsequence which converges in  $X$ . To prove relative compactness we use the Arzelà–Ascoli Theorem [15, Theorem A2.1] which, in this context, says that any sequence of functions  $\phi_1, \phi_2, \dots$  in  $C(\Omega)$  that is uniformly bounded and equicontinuous has a subsequence that converges in  $C(\Omega)$ .

Take any  $\chi_1, \chi_2 \in K$ . Let  $\phi_1, \phi_2, \dots$  be a sequence of functions in  $K$  such that  $\chi_1 \leq \phi_i \leq \chi_2$  for all  $i$ . It is sufficient to show that the sequence of functions  $R\phi_1, R\phi_2, \dots$  is uniformly bounded and equicontinuous. For any  $\phi \in K$  and for all  $z \in \Omega$ ,

$$R\phi(z) \leq \int D(z, \bar{z}) \sigma(d\bar{s}, d\bar{z}) \leq \bar{D} \int \sigma(d\bar{s}, d\bar{z}), \quad (\text{A.1})$$

from Assumption (C). Hence, the sequence  $R\phi_1, R\phi_2, \dots$  is uniformly bounded. To prove the sequence is equicontinuous, take  $z_1, z_2 \in \Omega$  and  $\phi \in K$ . Then

$$\begin{aligned}
|R\phi(z_1) - R\phi(z_2)| &\leq \left| \int (D(z_1, \bar{z}) - D(z_2, \bar{z})) \frac{f(\phi(\bar{z}))}{1 - \bar{s} + f(\phi(\bar{z}))} \sigma(d\bar{s}, d\bar{z}) \right| \\
&\leq \int |D(z_1, \bar{z}) - D(z_2, \bar{z})| \sigma(d\bar{s}, d\bar{z}).
\end{aligned}$$

It now follows from Assumption (C) that the sequence of functions  $R\phi_1, R\phi_2, \dots$  is equicontinuous. Hence,  $R$  is order compact.

(iii) *Monotonicity*: For a given  $s \in [0, 1 - \epsilon]$ , define the function

$$g_s(x) = f(x)(1 - s + f(x))^{-1}, \quad x \in [0, \infty). \quad (\text{A.2})$$

From Assumption (F),  $f(x)$  is increasing, hence  $g_s(x)$  is increasing. It follows that if  $\phi_1 \leq \phi_2$  then

$$\int D(z, \bar{z}) \frac{f(\phi_1(\bar{z}))}{1 - \bar{s} + f(\phi_1(\bar{z}))} \sigma(d\bar{s}, d\bar{z}) \leq \int D(z, \bar{z}) \frac{f(\phi_2(\bar{z}))}{1 - \bar{s} + f(\phi_2(\bar{z}))} \sigma(d\bar{s}, d\bar{z}),$$

for all  $z \in \Omega$ . Hence,  $R\phi_1 \leq R\phi_2$ .

(iv) *Strong positivity*: As  $\phi \in K \setminus \{0\}$  there exists a  $z_0 \in \Omega$  and  $\delta > 0$  such that  $\phi(z) > \delta$  for all  $z$  in an open neighbourhood of  $z_0$  denoted  $N_0$ . Therefore,

$$\begin{aligned} \inf_{z \in \Omega} R\phi(z) &\geq \inf_{z \in \Omega} \int_{[0,1] \times N_0} D(z, \bar{z}) \frac{f(\phi(\bar{z}))}{1 - \bar{s} + f(\phi(\bar{z}))} \sigma(d\bar{s}, d\bar{z}) \\ &\geq \left( \inf_{z \in \Omega} \inf_{\bar{z} \in N_0} D(z, \bar{z}) \right) \frac{f(\delta)}{1 + f(\delta)} \sigma([0, 1] \times N_0), \end{aligned}$$

which is positive from Assumptions (G) and (H). Hence,  $R\phi \in \mathring{K}$  for any  $\phi \in K \setminus \{0\}$ .

(v) *Strong sublinearity*: For any  $\lambda \in (0, 1)$  and  $\phi \in \mathring{K}$ ,

$$\begin{aligned} R(\lambda\phi) - \lambda R\phi &= \int D(z, \bar{z}) \left[ \frac{f(\lambda\phi(\bar{z}))}{1 - \bar{s} + f(\lambda\psi(\bar{z}))} - \frac{\lambda f(\lambda\phi(\bar{z}))}{1 - \bar{s} + f(\psi(\bar{z}))} \right] \sigma(d\bar{s}, d\bar{z}) \\ &= \int D(z, \bar{z}) \frac{f(\lambda\phi(\bar{z})) - \lambda f(\phi(\bar{z}))}{1 - \bar{s} + f(\lambda\psi(\bar{z}))} \sigma(d\bar{s}, d\bar{z}) \\ &\quad + \int D(z, \bar{z}) \left[ \frac{\lambda f(\phi(\bar{z}))}{1 - \bar{s} + f(\lambda\psi(\bar{z}))} - \frac{\lambda f(\phi(\bar{z}))}{1 - \bar{s} + f(\psi(\bar{z}))} \right] \sigma(d\bar{s}, d\bar{z}) \\ &\geq \int D(z, \bar{z}) \frac{f(\lambda\phi(\bar{z})) - \lambda f(\phi(\bar{z}))}{1 - \bar{s} + f(\lambda\psi(\bar{z}))} \sigma(d\bar{s}, d\bar{z}). \end{aligned}$$

From Assumption (F),  $f$  is strictly concave so  $f(\lambda x) - \lambda f(x) > 0$  for any  $x > 0$ . Arguing as in the proof of part (iv), we see that this last integral is positive for all  $z \in \Omega$ .

Let  $R^n\phi$  denote the  $n$ -th application of the operator  $R$  to  $\phi$ . From Proposition A.1, we can apply the cone limit trichotomy [13, Theorem 6.3] to show that one of the following hold:

- For any  $\phi \in K \setminus \{0\}$ , the sequence  $\{R^n\phi\}_{n=0}^\infty$  is unbounded.
- For any  $\phi \in K$ , the sequence  $\{R^n\phi\}_{n=0}^\infty$  converges to 0, the unique fixed point of  $R$ .
- For any  $\phi \in K \setminus \{0\}$ , the sequence  $\{R^n\phi\}_{n=0}^\infty$  converges to  $\psi$ , the unique non-zero fixed point of  $R$ .

From equation (A.1) we may exclude the first possibility. It remains to determine conditions that will decide which of the latter two possibilities hold. For this, we briefly summarise some theory from functional analysis which extends the Perron–Frobenius Theorem.

Let  $\Lambda : C(\Omega) \mapsto \mathbb{R}$  denote a linear functional on  $C(\Omega)$ . A positive linear functional is a linear functional with the property that  $\Lambda h \geq 0$  if  $h \geq 0$  on  $\Omega$ . Since  $\Omega$  is compact, the Riesz–Markov Theorem [17, Section A.5] states that a positive linear functional on  $C(\Omega)$  can be expressed as

$$\Lambda(h) = \int h \, dm^*, \quad \text{for all } g \in C(\Omega),$$

for a unique positive Borel measure  $m^*$  on  $\Omega$ . The set of positive Borel measures on  $\Omega$  forms the dual cone  $K^*$ . Let  $\mathring{K}^*$  denote the interior of  $K^*$ . If the unique Borel measure  $m^*$  is identified for a positive linear functional then we write the linear functional as  $\langle g, m^* \rangle := \Lambda(g)$ .

**Theorem A.1.** (Krein-Rutman Theorem.) *Let  $K$  be a reproducing cone, with interior  $\mathring{K} \neq \emptyset$ , and let  $B$  be a strongly positive compact bounded linear operator on  $K$ . Then the spectral radius of  $B$ ,  $r(B)$ , is a simple eigenvalue of  $B$  and  $B^*$  (the adjoint of  $B$ ), and their associated eigenvectors belong to  $\mathring{K}$  and  $\mathring{K}^*$ . Furthermore, all other eigenvalues are strictly less in absolute value than  $r(B)$ .*

**Corollary A.1.** *Let  $B$  be a strongly positive compact bounded linear operator,  $s$  a positive number, and  $\phi \in K \setminus \{0\}$  satisfying  $s\phi \leq B\phi$ . Then  $s \leq r(B)$  with equality if and only if  $s\phi = B\phi$ .*

*Proof.* Let  $u^*$  be the eigenvector of  $B^*$  associated with the eigenvalue  $r(B)$ . From Theorem A.1,  $u^* \in \mathring{K}^*$ . As  $B\phi - s\phi \in K$ , we have  $0 \leq \langle B\phi - s\phi, u^* \rangle$  with equality if and only if  $s\phi = B\phi$ . Now

$$\langle B\phi - s\phi, u^* \rangle = \langle B\phi, u^* \rangle - s \langle \phi, u^* \rangle = \langle \phi, B^* u^* \rangle - s \langle \phi, u^* \rangle = (r(B) - s) \langle \phi, u^* \rangle.$$

As  $u^* \in \mathring{K}^*$  and  $\phi \neq 0$ ,  $\langle \phi, u^* \rangle > 0$ . Hence,  $s \leq r(B)$  with  $s = r(B)$  if and only if  $B\phi = s\phi$ . This completes the proof.

Adapting the arguments of Proposition A.1, we see that, under Assumptions (C), (G) and (H),  $\mathcal{A}$  is a strongly positive compact operator. Therefore, the Krein-Rutman Theorem can be applied to  $\mathcal{A}$ . The next proposition shows that the operator  $R$  can be bounded by  $\mathcal{A}$ .

**Proposition A.2.** *For any  $\phi \in K$ ,  $R\phi \leq \mathcal{A}\phi$  with equality if and only if  $\phi = 0$ .*

*Proof.* From Assumption (F),  $f$  is strictly concave, so the function  $g$  defined in equation (A.2) is also strictly concave in  $x$ . For any  $x > 0$  and  $s \in [0, 1)$ ,

$$\frac{f(x)}{1 - s + f(x)} < \frac{f'(0)}{1 - s} x.$$

For any  $\phi \in K \setminus \{0\}$ , the set  $\{(s, z) : 0 < \phi(z)\}$  will have positive  $\sigma$  measure by Assumption (G). Therefore, for all  $\phi \in K$  and all  $z \in \Omega$ ,

$$\int D(z, \bar{z}) \frac{f(\phi(\bar{z}))}{1 - \bar{s} + f(\phi(\bar{z}))} \sigma(d\bar{s}, d\bar{z}) < \int D(z, \bar{z}) \frac{f'(0)}{(1 - \bar{s})} \phi(\bar{z}) \sigma(d\bar{s}, d\bar{z}),$$

by Assumption (H). In other words, the inequality  $R\phi \leq \mathcal{A}\phi$  is strict for any  $\phi \in K \setminus \{0\}$ . To complete the proof we simply note that if  $\phi = 0$  then  $R\phi = \mathcal{A}\phi$ .

We are now able to prove the first part of Theorem 3.2.

**Lemma A.1.** *If  $r(\mathcal{A}) \leq 1$  then  $\psi = 0$  is the unique solution to equation (3.5).*

*Proof.* Suppose that there exists a  $\phi \in K \setminus \{0\}$  such that  $\phi = R\phi$ . From Proposition A.2,  $\phi \leq \mathcal{A}\phi$ . From Corollary A.1, this is only possible if  $r(\mathcal{A}) \geq 1$ . So, if  $r(\mathcal{A}) < 1$  then the only solution to equation (3.5) is  $\psi = 0$ . Now suppose that  $r(\mathcal{A}) = 1$ . Then Corollary A.1 implies that  $\phi = \mathcal{A}\phi$ . From Proposition A.2, this is only possible if  $\phi = 0$ .

It remains to consider the case where  $r(\mathcal{A}) > 1$ .

**Lemma A.2.** *If  $r(\mathcal{A}) > 1$  then there exists a unique non-zero solution to equation (3.5).*

*Proof.* Let  $u$  be the eigenvector of  $\mathcal{A}$  associated with the eigenvalue  $r(\mathcal{A})$ . Take  $\delta > 0$ . Then

$$R(\delta u) = \mathcal{A}(\delta u) + \int D(x, \bar{z}) \left( \frac{f(\delta u(\bar{z}))}{1 - \bar{s} + f(\delta u(\bar{z}))} - \frac{f'(0)\delta u(\bar{z})}{1 - \bar{s}} \right) \sigma(d\bar{s}, d\bar{z}). \quad (\text{A.3})$$

Since  $f$  is twice differentiable in a neighbourhood of zero, from Assumption (F), and the support of  $\sigma$  is restrict to  $[0, 1 - \epsilon] \times \Omega$  by Assumption (G), the second term on the right-hand side of equation (A.3) is  $o(\delta\|u\|^2)$  as  $\delta \rightarrow 0$ . As  $u$  is an eigenvector of  $\mathcal{A}$ ,  $u \in \mathring{K}$  and  $R(\delta u) = r(\mathcal{A})\delta u + o(\delta\|u\|^2)$ . By assumption  $r(\mathcal{A}) > 1$ , so for a sufficiently small  $\delta > 0$ ,  $\delta u \leq R(\delta u)$ . The set of continuous functions bounded below by  $\delta u$  and bounded above by one form a closed convex set. It follows from the monotonicity of  $R$  (part (iii) of Proposition A.1) that  $R$  maps this set into itself. Since  $R$  is a compact map (part (ii) of Proposition A.1), we can apply the Schauder fixed point Theorem [14, Theorem 5.1.2] to conclude that there exists at least one non-zero solution to equation (3.5). To complete the proof, we note from the cone limit set trichotomy that  $R$  can have at most one non-zero fixed point.

### A.3. Proof of Theorem 4.1

*Proof.* First note that the functionals defined by equations (4.3) and (4.4) are indeed probability generating functionals. Specifically, each of these is the probability generating functional of an MPC where the offspring from the member of the population located at  $(s, z)$  at time  $t$  is generated according to an inhomogeneous Poisson process with intensity measure  $f'(0)D(\cdot, z)\sigma$  and the original member of the population survives to time  $t + 1$  with probability  $s$ .



The proof now proceeds by induction on  $t$ . Let  $\mathcal{V}_0$  denote the set  $\{h : h \in \mathcal{V}, \inf_{s,z} h(s, z) > 0\}$ . Since  $\{\mu_{n,0}\}_{n=1}^\infty$  converges weakly to some point process  $\mu_0$ , the probability generating functionals  $G_{n,0}[h]$  converge to  $G_0[h]$  for each continuous  $h \in \mathcal{V}_0$  [7, Proposition 11.1.VIII]. Now assume that for some  $T \geq 0$ ,  $G_{n,T}[h] \rightarrow G_T[h]$  for each continuous  $h \in \mathcal{V}_0$ . The probability generating functional of  $\mu_{n,T+1}$  is

$$\begin{aligned} G_{n,T+1}[h] &= \mathbb{E} \left( \prod_{i=1}^n (X_{i,T+1}^n h(s_i, z_i) + 1 - X_{i,T+1}^n) \right) \\ &= \mathbb{E} \left( \mathbb{E} \left( \prod_{i=1}^n (X_{i,T+1}^n h(s_i, z_i) + 1 - X_{i,T+1}^n) \mid X_T^n, s^n, z^n \right) \right). \end{aligned}$$

Note that  $\mathbb{E}(X_{i,T+1}^n \mid X_T^n, s^n, z^n) = U(s_i)X_{i,T}^n + f(C_i(T))(1 - X_{i,T}^n)$ . Therefore,

$$\begin{aligned} G_{n,T+1}[h] &= \mathbb{E} \left( \prod_{i=1}^n (X_{i,T}^n U(s_i) (h(s_i, z_i) - 1) + 1 + (1 - X_{i,T}^n) f(C_i(T)) (h(s_i, z_i) - 1)) \right) \\ &= \mathbb{E} \left( \prod_{i=1}^n (1 + X_{i,T}^n U(s_i) (h(s_i, z_i) - 1)) (1 + (1 - X_{i,T}^n) f(C_i(T)) (h(s_i, z_i) - 1)) \right). \end{aligned}$$

Define

$$\tilde{G}_{n,T+1}[h] := \mathbb{E} \left( \prod_{i=1}^n (1 + X_{i,T}^n U(s_i) (h(s_i, z_i) - 1)) (1 + f(C_i(T)) (h(s_i, z_i) - 1)) \right). \quad (\text{A.4})$$

We want to show that  $|G_{n,T+1}[h] - \tilde{G}_{n,T+1}[h]| \rightarrow 0$  as  $n \rightarrow \infty$ .

$$\begin{aligned} &|G_{n,T+1}[h] - \tilde{G}_{n,T+1}[h]| \\ &= \mathbb{E} \left| \left( \prod_{i=1}^n (1 + X_{i,T}^n U(s_i) (h(s_i, z_i) - 1)) (1 + (1 - X_{i,T}^n) f(C_i(T)) (h(s_i, z_i) - 1)) \right) \right. \\ &\quad \left. \times \left( 1 - \prod_{i=1}^n (1 + X_{i,T}^n f(C_i(T)) (h(s_i, z_i) - 1)) \right) \right| \\ &\leq \mathbb{E} \left| 1 - \prod_{i=1}^n (1 + X_{i,T}^n f(C_i(T)) (h(s_i, z_i) - 1)) \right|. \end{aligned} \quad (\text{A.5})$$

Clearly,  $\prod_{i=1}^n (1 + X_{i,T}^n f(C_i(T)) (h(s_i, z_i) - 1)) \leq 1$ . Let  $h_0 = \inf_{s,z} h(s, z)$ . Then

$$\begin{aligned} \mathbb{E} \left| 1 - \prod_{i=1}^n (1 + X_{i,T}^n f(C_i(T)) (h(s_i, z_i) - 1)) \right| &\leq 1 - \mathbb{E} \left( \exp \left( \sum_{i=1}^n \log (1 - (1 - h_0) X_{i,T}^n f(C_i(T))) \right) \right) \\ &\leq 1 - \mathbb{E} \left( \exp \left( (\log h_0) \sum_{i=1}^n X_{i,T}^n f(C_i(T)) \right) \right), \end{aligned}$$

where the second inequality follows from the bound  $\log(1 - (1 - h_0)x) \geq (\log h_0)x$  for  $x \in [0, 1]$ .

From Assumption (F),  $f(x) \leq f'(0)x$  for  $x \in [0, \infty)$ . Therefore,

$$\begin{aligned} \mathbb{E} \left| 1 - \prod_{i=1}^n \left( 1 + X_{i,T}^n f(C_i(T)) (h(s_i, z_i) - 1) \right) \right| &\leq 1 - \mathbb{E} \left( \exp \left( (\log h_0) f'(0) \sum_{i=1}^n X_{i,T}^n C_i(T) \right) \right) \\ &\leq 1 - \mathbb{E} \left( \exp \left( (\log h_0) f'(0) n^{-1} \bar{D} \left( \sum_{i=1}^n X_{i,t}^n \right)^2 \right) \right), \end{aligned}$$

and the final inequality comes from Assumption (C). Lemma A.3, given at the end of this proof, states that  $\exp(\alpha \sum_{i=1}^n X_{i,t}^n)$  is uniformly integrable for any  $\alpha > 0$ . Hence,  $(\sum_{i=1}^n X_{i,t}^n)^2$  is uniformly integrable and  $n^{-1} (\sum_{i=1}^n X_{i,t}^n)^2 \xrightarrow{P} 0$ . Therefore, the right-hand side of inequality (A.5) goes to zero as required.

To examine the limit of  $\widetilde{G}_{n,T+1}(h)$  given in equation (A.4) as  $n \rightarrow \infty$ , we note that

$$\begin{aligned} &\sum_{i=1}^n \log(1 + f(C_i(T)) (h(s_i, z_i) - 1)) = \sum_{i=1}^n f(C_i(T)) (h(s_i, z_i) - 1) + \epsilon_{n,1} \\ &= \sum_{i=1}^n f'(0) \left( n^{-1} \sum_{j=1}^n X_{j,T}^n D(z_i, z_j) \right) (h(s_i, z_i) - 1) + \epsilon_{n,1} + \epsilon_{n,2} \\ &= f'(0) \sum_{j=1}^n X_{j,T}^n \left( n^{-1} \sum_{i=1}^n D(z_i, z_j) (h(s_i, z_i) - 1) \right) + \epsilon_{n,1} + \epsilon_{n,2} \\ &= f'(0) \sum_{j=1}^n X_{j,T}^n \int D(z, z_j) (h(s, z) - 1) \sigma_n(ds, dz) + \epsilon_{n,1} + \epsilon_{n,2}, \end{aligned}$$

where

$$\begin{aligned} \epsilon_{n,1} &= \sum_{i=1}^n (\log(1 + f(C_i(t)) (h(s_i, z_i) - 1)) - f(C_i(t)) (h(s_i, z_i) - 1)) \\ \epsilon_{n,2} &= \sum_{i=1}^n (f(C_i(t)) (h(s_i, z_i) - 1) - f'(0) C_i(t) (h(s_i, z_i) - 1)). \end{aligned}$$

Let  $\widehat{G}_{n,T+1}[h]$  denote

$$\mathbb{E} \left( \left( \prod_{i=1}^n \left( 1 + X_{i,T}^n U(s_i) (h(s_i, z_i) - 1) \right) \right) \exp \left( f'(0) \sum_{j=1}^n X_{j,t}^n \int D(z, z_j) (h(s, z) - 1) \sigma_n(ds, dz) \right) \right).$$

To show  $|\widetilde{G}_{n,T+1}[h] - \widehat{G}_{n,T+1}[h]| \rightarrow 0$  as  $n \rightarrow \infty$  the difference is bounded by

$$\begin{aligned} &|\widetilde{G}_{n,T+1}[h] - \widehat{G}_{n,T+1}[h]| \leq \mathbb{E} \left( \left( \prod_{i=1}^n \left( 1 + X_{i,T}^n U(s_i) (h(s_i, z_i) - 1) \right) \right) \right. \\ &\quad \times \left. \exp \left( f'(0) \sum_{j=1}^n X_{j,T}^n \int D(z, z_j) (h(s, z) - 1) \sigma_n(ds, dz) \right) \times |(\exp(\epsilon_{n,1} + \epsilon_{n,2}) - 1)| \right) \\ &\leq \mathbb{E} |(\exp(\epsilon_{n,1} + \epsilon_{n,2}) - 1)| \leq \mathbb{E} \exp(|\epsilon_{n,1}| + |\epsilon_{n,2}|) - 1. \end{aligned} \tag{A.6}$$

Note that  $|\log(1-x) + x| \leq -\log(h_0)/(1-h_0)^2 x^2$  for  $x \in [0, 1-h_0]$ . From Assumptions (C) and (F),  $|\epsilon_{n,1}|$  is bounded by

$$|\epsilon_{n,1}| \leq \frac{-\log(h_0)}{(1-h_0)^2} \sum_{i=1}^n f^2(C_i(t)) \leq \frac{-\log(h_0)}{(1-h_0)^2} (f'(0)\bar{D})^2 n^{-1} \left( \sum_{i=1}^n X_{i,t} \right)^2.$$

From Assumption (F), there exists a  $c_1 > 0$  such that  $|f(x) - f'(0)x| \leq c_1 x^2$  for  $x \in [0, \infty)$ .

Therefore,  $\epsilon_{n,2}$  can be bounded by

$$|\epsilon_{n,2}| \leq \sum_{i=1}^n |f(C_i(T)) - f'(0)C_i(T)| \leq c_2 n^{-1} \left( \sum_{i=1}^n X_{i,T}^n \right)^2,$$

for some  $c_2 > 0$ , so  $|\epsilon_{n,1}| + |\epsilon_{n,2}| < c_3 n^{-1} \left( \sum_{j=1}^n X_{j,T}^n \right)^2$  for some  $c_3 > 0$ . Since  $\exp(\alpha \sum_{i=1}^n X_{i,T}^n)$  is uniformly integrable (Lemma A.3) for any  $\alpha > 0$  and  $n^{-1} \left( \sum_{i=1}^n X_{i,T}^n \right)^2 \xrightarrow{P} 0$ , the right-hand side of inequality (A.6) goes to zero.

We may now express  $G_{n,T}[G_1[h | (s, z)]]$  as

$$\begin{aligned} G_{n,T}[G_1[h | (s, z)]] &= \mathbb{E} \left( \left( \prod_{i=1}^n (X_{i,T}^n (1 - U(s_i)(1 - h(s_i, z_i))) + 1 - X_{i,T}^n) \right) \right. \\ &\quad \left. \times \exp \left( f'(0) \sum_{j=1}^n X_{j,T}^n \int D(z, z_j) (h(s, z) - 1) \sigma(ds, dz) \right) \right). \end{aligned}$$

We now show that  $|\widehat{G}_{n,T+1}[h] - G_{n,T}[G_1[h | (s, z)]]| \rightarrow 0$  as  $n \rightarrow \infty$ . Let  $\mathcal{G}(h; X_T^n, \sigma_n)$  denote

$$\sum_{j=1}^n X_{j,T}^n \left( \int D(z, z_j) (h(s, z) - 1) \sigma_n(ds, dz) - \int D(z, z_j) (h(s, z) - 1) \sigma(ds, dz) \right).$$

Noting that  $1 + X_{i,T} U(s_i)(h(s_i, z_i) - 1) = X_{i,T}(1 - U(s_i)(1 - h(s_i, z_i))) + 1 - X_{i,T}$ , we see that

$$\begin{aligned} & \left| \widehat{G}_{n,T+1}[h] - G_{n,T}[G_1[h | (s, z)]] \right| \\ &= \left| \mathbb{E} \left( \left( \prod_{i=1}^n (X_{i,T}^n (1 - U(s_i)(1 - h(s_i, z_i))) + 1 - X_{i,T}^n) \right) \right. \right. \\ &\quad \left. \left. \times \exp \left( f'(0) \sum_{j=1}^n X_{j,T}^n \int D(z, z_j) (h(s, z) - 1) \sigma(ds, dz) \right) (\exp(\mathcal{G}(h; X_T^n, \sigma_n)) - 1) \right) \right| \\ &\leq \mathbb{E} |\exp(\mathcal{G}(h; X_T^n, \sigma_n)) - 1|. \end{aligned} \tag{A.7}$$

From Assumption (C),  $|\mathcal{G}(h; X_T^n, \sigma_n)| \leq 2\bar{D} \sum_{j=1}^n X_{j,T}^n$  so  $\exp(\mathcal{G}(h; X_T^n, \sigma_n))$  is uniformly integrable. To show that the right-hand side of (A.7) converges to zero as  $n \rightarrow \infty$ , it is sufficient to show that  $\mathcal{G}(h; X_T^n, \sigma_n)$  converges to zero in probability. Define the measures  $\sigma_n^{(h)}$  and  $\sigma^{(h)}$

as being the measures with Radon-Nikodym derivative  $(1 - h(s, z))$  with respect to  $\sigma_n$  and  $\sigma$ , respectively. As  $h$  is a continuous function in  $\mathcal{V}_0$  and from Assumption (E)  $\sigma_n \xrightarrow{d} \sigma$ , it follows that  $\sigma_n^{(h)} \xrightarrow{d} \sigma^{(h)}$ . By Assumption (C), the kernel  $D(\bar{z}, z)$  defines a uniformly bounded and equicontinuous family of functions on  $\Omega$ . Therefore, we can apply a small modification of Theorem 3.1 of [27] to show that

$$\sup_{z \in \Omega} \left( \int D(\bar{z}, z) (h(\bar{s}, \bar{z}) - 1) \sigma_n(d\bar{s}, d\bar{z}) - \int D(\bar{z}, z) (h(\bar{s}, \bar{z}) - 1) \sigma(d\bar{s}, d\bar{z}) \right) \xrightarrow{p} 0.$$

As  $\exp(\alpha \sum_{i=1}^n X_{i,t}^n)$  is uniformly integrable for any  $\alpha > 0$  (Lemma A.3), so is  $\sum_{i=1}^n X_{i,t}^n$ . Hence,  $\mathcal{G}(h; X^n, \sigma_n) \xrightarrow{p} 0$ . Therefore,  $\left| \widehat{G}_{n,T+1}[h] - G_{n,T}[G_1[h | (s, z)]] \right| \rightarrow 0$ .

It is easily checked that if  $h$  is a continuous function in  $\mathcal{V}_0$  then  $G_1[h | (s, z)]$  is also a continuous function in  $\mathcal{V}_0$ . Applying the induction hypothesis, we see  $G_{n,T}[G_1[h | (s, z)]] \rightarrow G_T[G_1[h | (s, z)]]$  for all continuous  $h \in \mathcal{V}_0$ . Hence,  $G_{n,T+1}[h] \rightarrow G_T[G_1[h | (s, z)]] = G_{T+1}[h]$  for all continuous  $h \in \mathcal{V}_0$ .

**Lemma A.3.** *Suppose that for all  $\alpha > 0$   $\sup_n \mathbb{E}(\exp(\alpha \sum_{i=1}^n X_{i,0}^n)) < \infty$ . Then for all  $\alpha > 0$  and all  $t \geq 0$   $\sup_n \mathbb{E}(\exp(\alpha \sum_{i=1}^n X_{i,t}^n)) < \infty$ .*

*Proof.* The proof proceeds by induction on  $t$ . By assumption  $\sup_n \mathbb{E}(\exp(\alpha \sum_{i=1}^n X_{i,0}^n)) < \infty$  for all  $\alpha > 0$ . Suppose now that for some  $T \geq 0$ ,  $\sup_n \mathbb{E}(\exp(\alpha \sum_{i=1}^n X_{i,T}^n)) < \infty$  for all  $\alpha > 0$ .

Then

$$\begin{aligned} \mathbb{E} \left( \exp \left( \alpha \sum_{i=1}^n X_{i,T+1}^n \right) \right) &= \mathbb{E} \left( \mathbb{E} \left( \exp \left( \alpha \sum_{i=1}^n X_{i,T+1}^n \right) \middle| X_T^n, s^n, z^n \right) \right) \\ &= \mathbb{E} \left( \prod_{i=1}^n \left( (1 - s_i + s_i \exp(\alpha)) X_{i,T}^n + (1 - f(C_i(T)) + f(C_i(T)) \exp(\alpha)) (1 - X_{i,T}^n) \right) \right) \\ &\leq \mathbb{E} \left( \prod_{i=1}^n \left( X_{i,T}^n \exp(\alpha) + 1 + f(C_i(T)) (\exp(\alpha) - 1) \right) \right) \\ &\leq \mathbb{E} \left( \exp \left( \sum_{i=1}^n \left( \exp(\alpha) X_{i,T}^n + f(C_i(T)) (\exp(\alpha) - 1) \right) \right) \right) \\ &\leq \mathbb{E} \left( \exp \left( \sum_{i=1}^n \left( \exp(\alpha) X_{i,T}^n + f'(0) \bar{D} n^{-1} (\exp(\alpha) - 1) \sum_{j=1}^n X_{j,T}^n \right) \right) \right) \\ &\leq \mathbb{E} \left( \exp \left( \exp(\alpha) (1 + f'(0) \bar{D}) \sum_{i=1}^n X_{i,T}^n \right) \right). \end{aligned} \tag{A.8}$$

Since  $\sup_n \mathbb{E}(\exp(\alpha \sum_{i=1}^n X_{i,T}^n)) < \infty$  for all  $\alpha > 0$ , inequality (A.8) implies that, for all  $\alpha > 0$ ,  $\sup_n \mathbb{E}(\exp(\alpha \sum_{i=1}^n X_{i,T+1}^n)) < \infty$ .

#### A.4. Proof of Theorem 4.2

*Proof.* Let  $K$  denote the cone of continuous positive functions on  $\Omega$  and let  $K_1 = \{\phi : \phi \in K, \phi \leq 1\}$ . Define the operator

$$R\phi(z) = \exp\left(-f'(0) \int D(\bar{z}, z) \left(\frac{1 - \phi(\bar{z})}{1 - U(\bar{s})\phi(\bar{z})}\right) \sigma(d\bar{s}, d\bar{z})\right), \quad \phi \in K_1. \quad (\text{A.9})$$

Note that under Assumption (G),  $1 - U(s)\phi(z) > 0$  on the support of  $\sigma$  for any  $\phi \in K_1$  and  $R : K_1 \mapsto K_1$ . From Theorem 3.1 of Moyal [22], the asymptotic extinction probability of a multiplicative population chain starting at  $(s, z)$  is given by the smallest nonnegative solution to  $h(s, z) = G_1[h \mid (s, z)]$ . Denote this solution by  $h^*(s, z)$  and define

$$\psi[h](z) = \exp\left(-f'(0) \int D(\bar{z}, z) (1 - h(\bar{s}, \bar{z})) \sigma(d\bar{s}, d\bar{z})\right). \quad (\text{A.10})$$

Note that for any  $h \in K_1$ ,  $\psi[h] \in K_1$ . Substituting equation (A.10) into  $G_1[h \mid (s, z)]$ , we obtain

$$h^*(s, z) = (1 - U(s) + U(s)h^*(s, z)) \psi[h^*](z). \quad (\text{A.11})$$

Hence,  $h^*$  satisfies

$$h^*(s, z) = \frac{(1 - U(s)) \psi[h^*](z)}{1 - U(s) \psi[h^*](z)} \quad (\text{A.12})$$

on the support of  $\sigma$ . Substituting this back into equation (A.10), we see that  $\psi[h^*](z)$  satisfies the equation  $\psi(z) = R\psi(z)$ . From the definition of  $R$ ,  $\psi(z) = 1$  for some  $z \in \Omega$  if and only if  $h = 1$ ,  $\sigma - a.e.$  Since the right-hand side of equation (A.12) is increasing in  $\psi$ ,  $h^*$  will be determined by the smallest nonnegative fixed point of the operator  $R$ .

Recall  $1 - \exp(-x) \leq x$  for all  $x \geq 0$  with equality if and only if  $x = 0$ . Therefore,

$$\begin{aligned} 1 - R\psi(z) &\leq f'(0) \int D(\bar{z}, z) \left(\frac{1 - \psi(\bar{z})}{1 - U(\bar{s})\psi(\bar{z})}\right) \sigma(d\bar{s}, d\bar{z}) \\ &\leq f'(0) \int \frac{D(\bar{z}, z)}{(1 - U(\bar{s}))} (1 - \psi(\bar{z})) \sigma(d\bar{s}, d\bar{z}) = \mathcal{A}(1 - \psi)(z), \end{aligned} \quad (\text{A.13})$$

where the equality follows as  $D$  is symmetric with respect to its arguments from Assumption (C). Note that we have equality if and only if  $\psi(z) = 1$  for all  $z$ . As in the proof of Theorem 3.2, the operator  $\mathcal{A}$  is strongly positive on  $K$ . If  $\psi^*$  is the smallest nonnegative fixed point of  $R$  then

$$1 - \psi^* = 1 - R\psi^* \leq \mathcal{A}(1 - \psi^*). \quad (\text{A.14})$$

Let  $r(\mathcal{A})$  denote the spectral radius of  $\mathcal{A}$ . As  $\psi^* \in K_1$ ,  $1 - \psi^* \in K$  and we can apply Corollary A.1 to conclude that if  $r(\mathcal{A}) < 1$  then  $\psi^*(z) = 1$  for all  $z \in \Omega$ . Furthermore, if  $r(\mathcal{A}) = 1$  then

$1 - \psi^* = \mathcal{A}(1 - \psi^*)$ . As the inequality (A.14) is strict for  $\psi^* \neq 1$ , it follows that  $\psi^*(z) = 1$ , for all  $z \in \Omega$ . Hence, if  $r(\mathcal{A}) \leq 1$  then  $h^*(s, z) = 1$  on the support of  $\sigma$ .

We now consider the case of  $r(A) > 1$ . We aim to find a convex subset of  $K_1$  not containing the function 1 such that  $R$  maps this set to itself. We can then apply the Schauder fixed point Theorem [14, Theorem 5.1.2] to show that  $\psi^* \neq 1$ . Clearly  $R0 \in K_1$ . Since the function  $(1 - x)(1 - sx)^{-1}$  is decreasing in  $x > 0$  for any  $s \in [0, 1 - \epsilon]$ , it follows that for any  $\phi_1, \phi_2 \in K_1$  such that  $\phi_1 \leq \phi_2$ ,  $R\phi_1 \leq R\phi_2$ . From the Krein–Rutman Theorem (see Theorem A.1)  $r(\mathcal{A})$  is an eigenvalue of  $\mathcal{A}$  and has an associated eigenvector  $u \in K$ . For any  $\delta > 0$  sufficiently small,  $R(1 - \delta u) = R(1) - \delta \mathcal{A}u + \epsilon_{\delta, u} = 1 - \delta r(A)u + \epsilon_{\delta, u}$ , where

$$\begin{aligned} \epsilon_{\delta, u} &= R(1 - \delta u) - 1 + f'(0) \int D(\tilde{z}, z) \frac{\delta u(\tilde{z})}{1 - \tilde{s} + \tilde{s}\delta u(\tilde{z})} \sigma(d\tilde{s}, d\tilde{z}) \\ &\quad + \delta \int D(\tilde{z}, z) u(\tilde{z}) \left( \frac{1}{1 - \tilde{s} + \delta \tilde{s} u(\tilde{z})} - \frac{1}{1 - \tilde{s}} \right) \sigma(d\tilde{s}, d\tilde{z}). \end{aligned}$$

It is seen that  $\epsilon_{\delta, u}$  is  $O(\delta^2)$  since

$$\begin{aligned} &\left| R(1 - \delta u) - 1 + f'(0) \int D(\tilde{z}, z) \frac{\delta u(\tilde{z})}{1 - \tilde{s} + \tilde{s}\delta u(\tilde{z})} \sigma(d\tilde{s}, d\tilde{z}) \right| \\ &\leq (f'(0))^2 \left( \int D(\tilde{z}, z) \frac{\delta u(\tilde{z})}{1 - \tilde{s} + \tilde{s}\delta u(\tilde{z})} \sigma(d\tilde{s}, d\tilde{z}) \right)^2 \leq (f'(0))^2 \bar{D}^2 \delta^2 \left( \int \frac{u(\tilde{z})}{1 - \tilde{s}} \sigma(d\tilde{s}, d\tilde{z}) \right)^2 \end{aligned}$$

and

$$\int D(\tilde{z}, z) u(\tilde{z}) \left| \frac{1}{1 - \tilde{s} + \delta \tilde{s} u(\tilde{z})} - \frac{1}{1 - \tilde{s}} \right| \sigma(d\tilde{s}, d\tilde{z}) \leq \delta \bar{D} \int \frac{u(\tilde{z})}{(1 - \tilde{s})^2} \sigma(d\tilde{s}, d\tilde{z}),$$

where both integrals are finite from Assumption (G). Therefore, for  $\delta > 0$  sufficiently small,  $R(1 - \delta u) \leq 1 - \delta u$  when  $r(\mathcal{A}) > 1$ . Take the convex set of functions  $K_u := \{\phi : \phi \in K, \phi \leq 1 - \delta u\}$ . We see that  $R : K_u \mapsto K_u$  and hence, from the Schauder fixed point Theorem, there exists a  $\psi^* \in K_u$  satisfying  $\psi = R\psi$  if  $r(\mathcal{A}) > 1$ . Hence, if  $r(\mathcal{A}) > 1$  then  $h^*(s, z) < 1$  for all  $(s, z)$  in the support of  $\sigma$ . Finally,  $h^*(s, z)$  gives the asymptotic extinction probability of a multiplicative population chain starting with a single point at  $(s, z)$ . As the chains starting from two different points are conditionally independent, the asymptotic probability of extinction for the limiting metapopulation is  $\mathbb{E}(\prod_i h^*(s_i, z_i)) = G_0[h^*]$ , where the product is over the occupied patches in the limiting metapopulation at  $t = 0$ .

### Acknowledgements

We are grateful to the referee for helpful comments and suggestions. This work is supported by the Australian Research Council (Discovery grant DP120102398 and Centre of Excellence for Mathematics and Statistics of Complex Systems).

### References

- [1] BALL, F. (1983). The threshold behaviour of epidemic models. *J. Appl. Prob.* **20**, 227-241.
- [2] BALL, F. AND CLANCY, D. (1993). The final size and severity of a generalised stochastic multitype epidemic model. *J. Appl. Prob.* **25**, 721-736.
- [3] BALL, F. AND DONNELLY, P. (1995). Strong approximations for epidemic models. *Stoch. Proc. Appl.* **55**, 121.
- [4] BARBOUR, A. D. (2010). Coupling a branching process to an infinite dimensional epidemic process. *ESAIM-Prob. Stat.* **14**, 53-64.
- [5] BUCKLEY, F. M. AND POLLETT, P. K. (2010). Limit theorems for discrete-time metapopulation models. *Probability Surveys*, **7**, 53-83.
- [6] COURCHAMP, F., BEREC, L. AND GASCOIGNE, J. (2008). *Allee effects in ecology and conservation*. Oxford University Press, New York.
- [7] DALEY, D. J. AND VERE-JONES, D. (2008). *An Introduction to the Theory of Point Processes, Volume 2*, Springer, New York.
- [8] DAUTRAY, R. AND LIONS, J.-L. (1990). *Mathematical Analysis and Numerical Methods for Science and Technology, Volume 3*. Springer-Verlag, Berlin.
- [9] DURRETT, R. (1999). Stochastic spatial models. *SIAM Rev.* **41**, 677-718.
- [10] HANSKI, I. (1994). A practical model of metapopulation dynamics. *J. Anim. Ecol.* **63**, 151-162.
- [11] HANSKI, I., ALHO, J. AND MOILANEN, A. (2000). Estimating the parameters of survival and migration of individuals in metapopulations. *Ecology* **81**, 239-251.
- [12] HANSKI, I. AND OVASKAINEN, O. (2003). Metapopulation theory for fragmented landscapes. *Theoret. Popn Biol.* **64**, 119-127.
- [13] HIRSCH, M.W. AND SMITH, H. (2005). Monotone maps: a review. *J. Differ. Equ. Appl.* **11**, 379-398.
- [14] ISTRATESCU, V. I. (1981). *Fixed point theory : an introduction*. Kluwer, Boston.
- [15] KALLENBERG, O. (2002). *Foundations of modern probability*, 2nd edn. Springer, New York.

- [16] LEVINS, R. (1969). Some demographic and genetic consequences of environmental heterogeneity for biological control. *Bull. Entomol. Soc. Am.* **15**, 237-240.
- [17] MACCLUER, B.D. (2009). *Elementary Functional Analysis*, Springer, New York.
- [18] McVINISH, R. AND POLLETT, P. K. (2010). Limits of large metapopulations with patch dependent extinction probabilities. *Adv. Appl. Prob.* **42**, 1172-1186.
- [19] McVINISH, R. AND POLLETT, P. K. (2012). The limiting behaviour of a mainland-island metapopulation. *J. Math. Biol.* **64**, 775-801.
- [20] McVINISH, R. AND POLLETT, P. K. (2012). The limiting behaviour of a stochastic patch occupancy model. To appear in *J. Math. Biol.* [dx.doi.org/10.1007/s00285-012-0568-z](https://doi.org/10.1007/s00285-012-0568-z)
- [21] MOILANEN, A. (2004). SPOMSIM: software for stochastic patch occupancy models of metapopulation dynamics. *Ecol. Model.* **179**, 533-550.
- [22] MOYAL, J. E. (1962). Multiplicative population chains. *Proc. R. Soc. Ser. A* **266**, 518-526.
- [23] OVASKAINEN, O. AND CORNELL, S. J. (2006). Asymptotically exact analysis of stochastic metapopulation dynamics with explicit spatial structure. *Theoret. Popn Biol.* **69**, 13-33.
- [24] OVASKAINEN, O. AND HANSKI, I. (2001). Spatially structured metapopulation models: Global and local assessment of metapopulation capacity. *Theoret. Popn Biol.* **60**, 281-302.
- [25] PENROSE, M.D. (2008). Existence and spatial limit theorems for lattice and continuum particle systems. *Probability Surveys*, **5**, 1-36.
- [26] PETTIT, S., MOILANEN, A., HANSKI, I. AND BAGUETTE, M. (2001). Metapopulation dynamics of the bog fritillary butterfly: Movements between habitat patches. *Oikos* **92**, 491-500.
- [27] RANGA RAO, R. (1962). Relations between weak and uniform convergence of measures with applications. *Ann. Math. Statist.* **33**, 659-680.