

Molecular Epidemiology of NDM-1-Producing *Enterobacteriaceae* and *Acinetobacter baumannii* Isolates from Pakistan

Anna L. Sartor,^a Muhammad W. Raza,^b Shahid A. Abbasi,^c Kathryn M. Day,^b John D. Perry,^b David L. Paterson,^a Hanna E. Sidjabat^a

University of Queensland, UQ Centre for Clinical Research, Herston, Queensland, Australia^a; Department of Microbiology, Freeman Hospital, Newcastle upon Tyne, United Kingdom^b; Microbiology Department, Armed Forces Institute of Pathology, Rawalpindi, Pakistan^c

The molecular epidemiology of 66 NDM-producing isolates from 2 Pakistani hospitals was investigated, with their genetic relatedness determined using repetitive sequence-based PCR (Rep-PCR). PCR-based replicon typing and screening for antibiotic resistance genes encoding carbapenemases, other β -lactamases, and 16S methylases were also performed. Rep-PCR suggested a clonal spread of *Enterobacter cloacae* and *Escherichia coli*. A number of plasmid replicon types were identified, with the incompatibility A/C group (IncA/C) being the most common (78%). 16S methylase-encoding genes were coharbored in 81% of NDM-producing *Enterobacteriaceae*.

With the worldwide spread of the NDM-1 gene and its variants (NDM-2 to NDM-8) (1, 2), molecular epidemiological studies of global isolates using various genotyping techniques are essential for gaining a better understanding of how this spread is occurring. India, Pakistan, and Bangladesh are clearly major reservoir countries for *bla*_{NDM}, with numerous factors, such as antibiotic selection pressure, contributing to this current situation (3, 4). This study examines a group of 66 NDM-1-producing isolates from Pakistan for their genetic relatedness, phylotype, plasmid replicon type, and plasmid transferability.

All isolates were acquired from stool samples from 37 distinct patients at two military hospitals in Rawalpindi, Pakistan (5). The samples were collected from inpatients (35%) and outpatients (65%). The isolates were tested for susceptibility to 17 antimicrobials using the Vitek 2 system. The MICs for meropenem, doripenem, fosfomycin, and amdinocillin were determined using a standard agar dilution methodology (5).

The isolates were reconfirmed for the presence of the carbapenem resistance gene *bla*_{NDM-1} by PCR, as previously described (6). PCR was also performed to detect *bla*_{OXA-48}, *bla*_{OXA-23}, *bla*_{VIM}, *bla*_{IMP}, *bla*_{KPC}, *bla*_{CTX-M-15}, *bla*_{SHV}, *bla*_{TEM}, *bla*_{OXA-1} group, AmpC β -lactamases, *bla*_{CMY-2}, and the 16S rRNA methylase genes *armA*,

rmtB, *rmtC*, and *rmtF* (6–9). The phylogenetic groups of *Escherichia coli* were determined using a multiplex (PCR)-based method (10).

Repetitive sequenced-based PCR (Rep-PCR)-based typing by the DiversiLab system (bioMérieux, Oakleigh, Australia) was used for assessing clonal relatedness. A cluster of closely related isolates was defined as isolates sharing >95% similarity and indistinguishable isolates of >97% (11, 12). PCR-based replicon typing analysis (PBRT) was performed to determine the plasmid incompatibility (Inc) groups for all *Enterobacteriaceae* isolates (13).

Ten genetically diverse *E. coli* isolates, based on different Rep-PCR profiles and phylogroups, were selected for transformation studies and typing by multilocus sequence typing (MLST). MLST

Received 4 February 2014 Returned for modification 7 April 2014

Accepted 21 June 2014

Published ahead of print 30 June 2014

Address correspondence to Anna L. Sartor, anna.sartor22@gmail.com.

Copyright © 2014, American Society for Microbiology. All Rights Reserved.

doi:10.1128/AAC.02425-14

TABLE 1 Resistance genes and plasmid replicon types of NDM-1-producing isolates

NDM-producing isolates (no.)	No. (%) of antibiotic resistance genes								Plasmid replicon type(s) (no.)
	β -Lactamases ^a				16S rRNA methylases				
	<i>bla</i> _{CMY-2}	<i>bla</i> _{CTX-M-15}	<i>bla</i> _{SHV} <i>bla</i> _{TEM}	<i>bla</i> _{OXA-1}	<i>rmtB</i>	<i>rmtC</i>	<i>rmtF</i>	<i>armA</i>	
<i>Escherichia coli</i> (30)	26 (87)	16 (53)	0, 19 (63)	9 (30)	8 (27)	20 (67)	0	15 (50)	IncHI1 (10), IncI1 (2), IncL/M (2), IncN (4), IncFIA (5), IncFIB (6), IncY (2), IncA/C (25), IncFII (9)
<i>Enterobacter cloacae</i> (21)	16 (76)	21 (100)	0, 20 (95)	20 (95)	0	15 (71)	0	14 (67)	IncA/C (19), untypeable (2)
<i>Citrobacter freundii</i> (4)	3 (75)	2 (50)	0, 2 (50)	2 (50)	0	0	0	1 (14)	IncA/C (2), IncFII (1), untypeable (1)
<i>Acinetobacter baumannii</i> (3)	0	0	0	0	0	0	0	0	ND ^b
<i>Klebsiella pneumoniae</i> (3)	1 (33)	3 (100)	3 (100), 3 (100)	3 (100)		1 (33)	0	1 (33)	IncN (1), IncA/C (1), untypeable (1)
<i>Pseudocitrobacter faecalis</i> (2)	0	2 (100)	0, 2 (100)	2 (100)	0	0	0	0	IncN (2)
<i>Providencia rettgeri</i> (2)	2 (100)	0	0, 0	1 (50)	0	1 (50)	0	2 (100)	IncA/C (1), IncN (2), IncY (1)
<i>Citrobacter braakii</i> (1)	1 (100)	0	0, 0	0	0	1 (100)	0	0	IncA/C (1)
Total no. of isolates ^c	49 (78)	44 (70)	3 (5), 46 (73)	37 (59)	8 (13)	38 (60)	0	33 (52)	IncA/C (49)

^a All isolates were negative for *bla*_{OXA-48}, *bla*_{IMP}, *bla*_{VIM}, and *bla*_{KPC}.

^b ND, not determined.

^c *A. baumannii* excluded from this total.




FIG 1 Southern blot hybridization of the PFGE gel with a specific *bla*_{NDM-1} probe. The black arrows indicate positive signals with the NDM-1 probe in each *E. coli* clinical isolate and its corresponding transformant. Lanes 1 to 8, PN1, PN1 TF1, PN7, PN7 TF1, PN14, PN14 TF1, PN18, and PN18 TF1, respectively.

included seven conserved housekeeping genes and was performed according to the *E. coli* MLST Database (see <http://mlst.warwick.ac.uk/mlst/dbs/Ecoli>).

The transferability of *bla*_{NDM-1}-carrying plasmids was investigated by electroporation. Plasmid DNA was prepared and electroporated into the recipient TOP10 *E. coli* (Invitrogen, Melbourne, Victoria, Australia), as previously described (6). Successful electrotransformants carrying *bla*_{NDM-1} were confirmed by PCR. The plasmid replicon type of the transformants acquiring *bla*_{NDM-1}-carrying plasmids was confirmed by PBRT. Plasmid size was determined by performing S1 endonuclease (Promega; Madison, WI, USA) restriction digestion using pulsed-field gel electrophoresis (PFGE) (14). PCR-amplified DNA probes of *bla*_{NDM-1} were labeled with digoxigenin nucleic acid (Roche, Mannheim, Germany).

Widespread dissemination of NDM-1 in Pakistan was first described in 2010 (15). In this study, we investigated the molecular epidemiology of a group of NDM-1-producing isolates from Pakistan, and we report here on the clonal relatedness of these isolates, providing an insight into the molecular characterization of *bla*_{NDM-1}-carrying plasmids.

The majority of the NDM-1-producing *Enterobacteriaceae* isolates coharbored an extended-spectrum β -lactamase (ESBL) gene,




FIG 2 Dendrogram analysis of DiversiLab Rep-PCR fingerprint of NDM-1-producing *E. cloacae* isolates.




FIG 3 Dendrogram analysis of DiversiLab Rep-PCR fingerprint of NDM-1-producing *E. coli* isolates by phylotype.

*bla*_{CTX-M-15} (70%). *bla*_{CMY-2} and the cooccurrence of 16S rRNA methylase genes encoding broad-spectrum aminoglycoside resistance, *rmtB*, *rmtC*, or *armA*, was detected in 47 (75%) isolates (Table 1). AmpC β-lactamase production coexisting with an ESBL was also high ($n = 36$ [73%]). The novel 16S rRNA methylase *rmtF* (16) was not detected; however, *rmtB* was found in 8 *E. coli* strains. A strong association between NDM-producing isolates harboring a 16S rRNA methylase-encoding gene has been well documented, particularly with *rmtC*. More recent studies in India, the United Kingdom, South Africa, and Nepal have reported the carriage of *rmtF* among NDM-harboring isolates (9, 17–19). *rmtB* together with *bla*_{NDM-1} is a rare association, although a recent outbreak of *E. coli* isolates in Bulgaria, harboring both *bla*_{NDM-1} and *rmtB* genes, raises new concerns for the acquisition of resistance determinants (20). Our results suggest that NDM-1-producing *Enterobacteriaceae* from Rawalpindi, Pakistan, have acquired a broad spectrum of singular and distinct resistance genes.

The plasmid incompatibility types HI1, I1, L/M, N, FIA, FIB, Y, A/C, and FII were identified among the NDM-1-producing isolates (Table 1). Plasmid replicon typing revealed the dominance of two incompatibility groups, IncA/C (78%) and IncF (33%), with

IncA/C occurring in multiple NDM-carrying species. IncA/C plasmids carrying *bla*_{NDM} have been reported in Pakistan (21).

Of the 10 *E. coli* isolates subjected to electroporation, the *bla*_{NDM-1} plasmids in 4 isolates were successfully electroporated. Plasmid replicon typing of these transformants confirmed that *bla*_{NDM-1} resides on IncA/C-, IncN-, IncFIB-, and IncFII-type plasmids. These replicon types have been reported in *Enterobacteriaceae* in many regions of the world (4).

Southern hybridization (Fig. 1) of the *E. coli* donors and their transformants revealed *bla*_{NDM-1} plasmid sizes ranging from ~50 kb to ~350 kb. The majority of the *bla*_{NDM-1} plasmids were ~140 kb in size. Among the 30 *E. coli* isolates, there was a predominance of the phylogenetic group B1 (57%), followed by phylotypes A (40%) and D (3%). It has been suggested that the distribution of *E. coli* phylotypes may be geographically dependent (22). Mushtaq et al. (23) found a prevalence of phylotype B1 among NDM isolates in Pakistan and no phylotype B2. Our study shows similar results.

Rep-PCR revealed two dominant clones among *Enterobacter cloacae*, one large cluster ($n = 17$), designated ECLI, and one small cluster of three isolates (ECLII) (see Fig. 2). Three clonal types were observed among 17 *E. coli* isolates, and the remaining *E. coli*




FIG 4 Dendrogram analysis of DiversiLab Rep-PCR fingerprint of NDM-1-producing *Citrobacter* isolates. *, novel species, *Pseudocitrobacter faecalis*.

isolates were diverse (Fig. 3). The three *Klebsiella pneumoniae* isolates were genetically diverse, while the three *Acinetobacter baumannii* isolates were considered identical (>99% similarity) (data not shown). Figure 4 shows the dendrogram for *Citrobacter* spp. and *Pseudocitrobacter faecalis* isolates (>99% similarity).

MLST differentiated the 10 representative *E. coli* strains into seven sequence types and one unknown sequence type (ST) (un-typeable). The sequence types included ST10 ($n = 2$), ST101 ($n = 2$), and single isolates representing STs 211, 226, 1431, 2598, and 3032. MLST studies on NDM-1-producing *E. coli* in the literature provide an incomplete and heterogeneous global distribution, suggesting a nonclonal pattern of spread for *bla*_{NDM-1} (24). In this study, the clinical isolates of *E. coli* representing STs 211, 226, 1431, 2598, and 3032 to our knowledge have not been reported in NDM-1-producing *E. coli*.

There were a number of limitations in our study, including the lack of clinical patient data and using fecal samples from 2 hospitals at a single point in time. It is difficult in this respect to obtain a clear epidemiological picture of NDM-producing isolates more widely in Pakistan.

The spread of *bla*_{NDM-1} is frequently associated with common and highly promiscuous plasmids resulting in a diverse range of species and clones harboring *bla*_{NDM-1}. However, the molecular epidemiology of our study may indicate that *bla*_{NDM-1} additionally disseminates via dominant clones. The potential role of such dominant clones as a factor of *bla*_{NDM-1} spread may be underrepresented due to the lack of large-scale surveillance and molecular epidemiological studies monitoring *bla*_{NDM-1} dissemination. In this scenario, we can see a situation in which a single clone of NDM-1 may become epidemic or pandemic.

REFERENCES

- Göttig S, Hamprecht AG, Christ S, Kempf VA, Wichelhaus TA. 2013. Detection of NDM-7 in Germany, a new variant of the New Delhi metallo- β -lactamase with increased carbapenemase activity. *J. Antimicrob. Chemother.* 68:1737–1740. <http://dx.doi.org/10.1093/jac/dkt088>.
- Tada T, Miyoshi-Akiyama T, Dahal RK, Sah MK, Ohara H, Kirikae T, Pokhrel BM. 2013. NDM-8 metallo- β -lactamase in a multidrug-resistant *Escherichia coli* strain isolated in Nepal. *Antimicrob. Agents Chemother.* 57:2394–2396. <http://dx.doi.org/10.1128/AAC.02553-12>.
- Nordmann P. 2013. Carbapenemase-producing *Enterobacteriaceae*: overview of a major public health challenge. *Med. Mal. Infect.* <http://dx.doi.org/10.1016/j.medmal.2013.11.007>.
- Wailan AM, Paterson DL. 2013. The spread and acquisition of NDM-1: a multifactorial problem. *Expert Rev. Anti Infect. Ther.* 12:91–115. <http://dx.doi.org/10.1586/14787210.2014.856756>.
- Perry JD, Naqvi SH, Mirza IA, Alizai SA, Hussain A, Ghirardi S, Orensa S, Wilkinson K, Woodford N, Zhang J, Livermore DM, Abbasi SA, Raza MW. 2011. Prevalence of faecal carriage of *Enterobacteriaceae* with NDM-1 carbapenemase at military hospitals in Pakistan, and evaluation of two chromogenic media. *J. Antimicrob. Chemother.* 66:2288–2294. <http://dx.doi.org/10.1093/jac/dkr299>.
- Sidjabat H, Nimmo GR, Walsh TR, Binotto E, Htin A, Hayashi Y, Li J, Nation RL, George N, Paterson DL. 2011. Carbapenem resistance in *Klebsiella pneumoniae* due to the New Delhi metallo- β -lactamase. *Clin. Infect. Dis.* 52:481–484. <http://dx.doi.org/10.1093/cid/ciq178>.
- Poirel L, Héritier C, Tolün V, Nordmann P. 2004. Emergence of oxacillinase-mediated resistance to imipenem in *Klebsiella pneumoniae*. *Antimicrob. Agents Chemother.* 48:15–22. <http://dx.doi.org/10.1128/AAC.48.1.15-22.2004>.
- Muzaheed, Doi Y, Adams-Haduch JM, Endimiani A, Sidjabat HE, Gaddad SM, Paterson DL. 2008. High prevalence of CTX-M-15-producing *Klebsiella pneumoniae* among inpatients and outpatients with urinary tract infection in southern India. *J. Antimicrob. Chemother.* 61:1393–1394. <http://dx.doi.org/10.1093/jac/dkn109>.
- Hidalgo L, Hopkins KL, Gutierrez B, Ovejero CM, Shukla S, Douthwaite S, Prasad KN, Woodford N, Gonzalez-Zorn B. 2013. Association of the novel aminoglycoside resistance determinant RmtF with NDM carbapenemase in *Enterobacteriaceae* isolated in India and the UK. *J. Antimicrob. Chemother.* 68:1543–1550. <http://dx.doi.org/10.1093/jac/dkt078>.
- Clermont O, Bonacorsi S, Bingen E. 2000. Rapid and simple determination of the *Escherichia coli* phylogenetic group. *Appl. Environ. Microbiol.* 66:4555–4558. <http://dx.doi.org/10.1128/AEM.66.10.4555-4558.2000>.
- Brolund A, Hæggman S, Edquist PJ, Gezelius L, Olsson-Liljequist B, Wisell KT, Giske CG. 2010. The DiversiLab system versus pulsed-field gel electrophoresis: characterisation of extended spectrum β -lactamase-producing *Escherichia coli* and *Klebsiella pneumoniae*. *J. Microbiol. Methods* 83:224–230. <http://dx.doi.org/10.1016/j.mimet.2010.09.004>.
- Pitout JDD, Campbell L, Church DL, Wang PW, Guttman DS, Gregson DB. 2009. Using a commercial DiversiLab semiautomated repetitive sequence-based PCR typing technique for identification of *Escherichia coli* clone ST131 producing CTX-M-15. *J. Clin. Microbiol.* 47:1212–1215. <http://dx.doi.org/10.1128/JCM.02265-08>.

13. Carattoli A, Bertini A, Villa L, Falbo V, Hopkins KL, Threlfall EJ. 2005. Identification of plasmids by PCR-based replicon typing. *J. Microbiol. Methods* 63:219–228. <http://dx.doi.org/10.1016/j.mimet.2005.03.018>.
14. Sidjabat HE, Paterson DL, Adams-Haduch JM, Ewan L, Pasculle AW, Muto CA, Tian GB, Doi Y. 2009. Molecular epidemiology of CTX-M-producing *Escherichia coli* isolates at a tertiary medical center in western Pennsylvania. *Antimicrob. Agents Chemother.* 53:4733–4739. <http://dx.doi.org/10.1128/AAC.00533-09>.
15. Kumarasamy KK, Toleman MA, Walsh TR, Bagaria J, Butt F, Balakrishnan R, Chaudhary U, Doumith M, Giske CG, Irfan S, Krishnan P, Kumar AV, Maharjan S, Mushtaq S, Noorie T, Paterson DL, Pearson A, Perry C, Pike R, Rao B, Ray U, Sarma JB, Sharma M, Sheridan E, Thirunarayan MA, Turton J, Upadhyay S, Warner M, Welfare W, Livermore DM, Woodford N. 2010. Emergence of a new antibiotic resistance mechanism in India, Pakistan, and the UK: a molecular, biological, and epidemiological study. *Lancet Infect. Dis.* 10:597–602. [http://dx.doi.org/10.1016/S1473-3099\(10\)70143-2](http://dx.doi.org/10.1016/S1473-3099(10)70143-2).
16. Galimand M, Courvalin P, Lambert T. 2012. RmtF, a new member of the aminoglycoside resistance 16S rRNA N7 G1405 methyltransferase family. *Antimicrob. Agents Chemother.* 56:3960–3962. <http://dx.doi.org/10.1128/AAC.00660-12>.
17. Rahman M, Shukla SK, Prasad KN, Ovejero CM, Pati BK, Tripathi A, Singh A, Srivastava AK, Gonzalez-Zorn B. 2014. Prevalence and molecular characterisation of New Delhi metallo- β -lactamases NDM-1, NDM-5, NDM-6 and NDM-7 in multidrug-resistant *Enterobacteriaceae* from India. *Int. J. Antimicrob. Agents* 44:30–37. <http://dx.doi.org/10.1016/j.ijantimicag.2014.03.003>.
18. Rubin JE, Peirano G, Peer AK, Govind CN, Pitout JD. 2014. NDM-1-producing *Enterobacteriaceae* from South Africa: moving towards endemicity? *Diagn. Microbiol. Infect. Dis.* 79:378–380. <http://dx.doi.org/10.1016/j.diagmicrobio.2014.04.003>.
19. Tada T, Miyoshi-Akiyama T, Dahal RK, Mishra SK, Ohara H, Shimada K, Kirikae T, Pokhrel BM. 2013. Dissemination of multidrug-resistant *Klebsiella pneumoniae* clinical isolates with various combinations of carbapenemases (NDM-1 and OXA-72) and 16S rRNA methylases (ArmA, RmtC and RmtF) in Nepal. *Int. J. Antimicrob. Agents* 42:372–374. <http://dx.doi.org/10.1016/j.ijantimicag.2013.06.014>.
20. Poirel L, Savov E, Nazli A, Trifonova A, Todorova I, Gergova I, Nordmann P. 2014. Outbreak caused by NDM-1- and RmtB-producing *Escherichia coli* in Bulgaria. *Antimicrob. Agents Chemother.* 58:2472–2474. <http://dx.doi.org/10.1128/AAC.02571-13>.
21. Poirel L, Dortet L, Bernabeu S, Nordmann P. 2011. Genetic features of *bla*_{NDM-1}-positive *Enterobacteriaceae*. *Antimicrob. Agents Chemother.* 55:5403–5407. <http://dx.doi.org/10.1128/AAC.00585-11>.
22. Nielsen KL, Dynesen P, Larsen P, Frimodt-Møller N. 2014. Faecal *Escherichia coli* from patients with *E. coli* urinary tract infection and healthy controls who have never had a urinary tract infection. *J. Med. Microbiol.* 63:582–589. <http://dx.doi.org/10.1099/jmm.0.068783-0>.
23. Mushtaq S, Irfan S, Sarma JB, Doumith M, Pike R, Pitout J, Livermore DM, Woodford N. 2011. Phylogenetic diversity of *Escherichia coli* strains producing NDM-type carbapenemases. *J. Antimicrob. Chemother.* 66:2002–2005. <http://dx.doi.org/10.1093/jac/dkr226>.
24. Bushnell G, Mitrani-Gold F, Mundy LM. 2013. Emergence of New Delhi metallo- β -lactamase type 1-producing *Enterobacteriaceae* and non-*Enterobacteriaceae*: global case detection and bacterial surveillance. *Int. J. Infect. Dis.* 17:e325–333. <http://dx.doi.org/10.1016/j.ijid.2012.11.025>.