

Accepted Manuscript

Healthcare 2020: Reengineering Healthcare Delivery to Combat Chronic Disease

Richard V. Milani , MD, Carl J. Lavie , MD

PII: S0002-9343(14)01035-3

DOI: [10.1016/j.amjmed.2014.10.047](https://doi.org/10.1016/j.amjmed.2014.10.047)

Reference: AJM 12764

To appear in: *The American Journal of Medicine*

Received Date: 14 October 2014

Revised Date: 17 October 2014

Accepted Date: 17 October 2014

Please cite this article as: Milani RV, Lavie CJ, Healthcare 2020: Reengineering Healthcare Delivery to Combat Chronic Disease, *The American Journal of Medicine* (2014), doi: 10.1016/j.amjmed.2014.10.047.

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.



Healthcare 2020: Reengineering Healthcare Delivery to Combat Chronic Disease

Richard V. Milani*, MD, Carl J. Lavie MD*†

*Department of Cardiovascular Diseases, John Ochsner Heart and Vascular Institute, Ochsner Clinical School – University of Queensland School of Medicine, New Orleans, Louisiana; and the

†Department of Preventive Medicine, Pennington Biomedical Research Center, Louisiana State University System, Baton Rouge, LA

Running Title: Healthcare 2020

Word Count: 2,793

Key Words: Healthcare delivery, chronic disease, information technology, social networking

Funding Source: none

Both authors have collaborated on the development of this manuscript and are in support of content. Authors have no relevant conflicts of interest to disclose relevant to this manuscript.

Correspondence to:

Richard V. Milani, M.D.
Ochsner Heart and Vascular Institute
Ochsner Clinic Foundation
1514 Jefferson Highway
New Orleans, LA. 70121
Tel: (504) 842-5874
Fax: (504) 842-5875
e-mail: rmilani@ochsner.org

Abstract

Chronic disease has become the great epidemic of our times, responsible for 75% of total health care costs and the majority of deaths in the United States. Our current delivery model is poorly constructed to manage chronic disease as evidenced by low adherence to quality indicators and poor control of treatable conditions. New technologies have emerged that can engage patients and offer additional modalities in the treatment of chronic disease. Modifying our delivery model to include team-based care in concert with patient-centered technologies offer great promise in managing the chronic disease epidemic.

ACCEPTED MANUSCRIPT

Chronic Disease Burden

Chronic disease represents the major driver of illness and healthcare utilization in the United States (U.S.), and its prevalence in the population is increasing. In 2010, chronic disease was responsible for seven of every ten deaths in the U.S. and accounted for over 75% of total health care costs.^{1,2} In just a five year span, from 2005 to 2010, the prevalence of chronic disease increased from 46% to 47% of the U.S. population, equivalent to an additional 8 million Americans, and by 2020 it is projected to increase by an additional 16 million comprising 48% of the population.³ It is noteworthy that over half of these individuals, or approximately 81 million of the U.S. population, will have multiple chronic conditions.⁴ Total cost of health care has also risen steadily over this period, and it is estimated that two-thirds of this escalation is due to the increased prevalence of chronic disease.⁵

As a rule, the proportion of the population diagnosed with chronic conditions will increase with age and today there are more Americans age 65 and older than at any other time in U.S. history.⁶ According to the Census Bureau, there were 40.3 million people age 65 and older in 2010, up 15.1 percent from 35 million in 2000 (compared to just a 9.7 percent increase for the total U.S. population).⁷ By 2020 the US population 65 years and older is projected to reach 53 million, with continual increases to 89 million by 2050.⁸ More significantly, the proportion of Americans 65 and older who report having one or more chronic diseases also rose, from 86.9 percent in 1998 to 92.2 percent in 2008.⁶ With the combination of increasing longevity and high rates of obesity and physical inactivity, this trend is expected to continue.^{6,9}

Current projections suggest that by 2020, there will probably be an additional fifteen million Americans with hypertension, twelve million with diabetes, four million with coronary heart disease, two million with stroke, and two million with heart failure.^{10,11} According to the World Health Organization and the Center for Disease Control and Prevention, the root cause of the epidemic in chronic disease is lack of physical activity and poor nutrition, which alone or in combination contributes to obesity and its attendant consequences. In the past 30 years, adult obesity rates in the U.S. have more than doubled, and today, more than two-thirds of American adults are either overweight or obese.¹² Nationally, 38% and 23% of adults and 36% and 37% of adolescents report consuming fruits and vegetables, respectively, less than one time daily.¹³ These risk factors

coupled with tobacco use and excess alcohol consumption represent the major lifestyle factors leading to the pandemic of chronic disease in the U.S. and globally.¹⁴

Treatment Options in Chronic Disease

Although chronic disease represents the leading cause of death in the U.S., 40% of all premature death is due to behaviors amenable to change. Maximizing disease outcomes will therefore require the necessary time and expertise needed for a careful assessment and modification of lifestyle factors.¹⁵ In the primary care setting, the median length of a physician visit is less than 15 minutes during which a median of six topics will be covered, leaving little if any time to formally assess and address the root causes of many chronic diseases including poor nutrition and physical inactivity.^{16,17} A survey conducted in 2006 revealed for example, that only 65% of obese patients were provided advice to lose weight by their physicians, and recommendations for physical activity are also rarely addressed.¹⁸ However, when lifestyle modification advice is provided, patient adherence rates regarding weight loss, smoking cessation, or dietary changes are remarkably low, and healthcare professionals have identified a lack of knowledge, skills, and practical tools as major barriers to successful intervention.¹⁹ Higher success rates for lifestyle modification have been achieved however through group programs utilizing non-physician personnel such as cardiac rehabilitation and exercise training. A contributing factor in the success of these programs is related to the impact of the group dynamic and social support created when patients are exposed to other individuals with the same condition at various stages of lifestyle change.^{20,21} Although these formal structured programs have proven successful in effecting lifestyle change and result in significant improvements in excess weight, smoking cessation, exercise capacity, blood pressure, insulin sensitivity, and lipids, they are unfortunately limited to a small number of patients who present following a cardiac event.

Since assessment and treatment of lifestyle behaviors for chronic disease are infrequent, and when provided, poorly adhered to, physicians are often limited to the sole therapeutic option of medication in order to treat the secondary conditions created from poor lifestyle such as hypertension or diabetes. Although medication is clearly a proven and much needed therapeutic in the management of many chronic diseases, an unfortunate consequence of its overreliance is that medication-related events have now become a major health concern, particularly among the

elderly where chronic disease is most prevalent. On average, individuals aged 65-69 years old take 14 prescriptions per year (80-84 years take 18 prescriptions per year), and medication-related problems are now one of the top five causes of death in this age group. It is estimated that 28% of hospitalizations among seniors are due to adverse drug reactions, and 32,000 seniors suffer hip fractures each year due to falls caused by medication-related problems.^{22,23} These issues would be best addressed by additionally providing non-pharmacologic alternatives, and the use of emerging technologies has recently demonstrated potential in this regard.¹⁹

Health-focused mobile application software (apps) and wearable devices (wearables) designed for increasing exercise, nutrition education and counseling, smoking cessation and weight loss programs have demonstrated positive results in effecting lifestyle change in patients presenting with chronic disease.²⁴⁻²⁶ Apps and wearables have the opportunity to provide appropriate level, tailored education, patient-friendly data visualization tools, exciting gamification strategies, regular feedback with prompts, and other impactful tactics to positively create healthy behaviors. Many apps also encourage patients to engage in social networks where patients have the option of interacting with other individuals seeking similar behavioral change; these social interactions are not trivial, and have been shown to be important in maintaining motivation, a key component of successful behavior change.²⁷⁻²⁹ Apps and wearables have now been successfully implemented in secondary prevention as a virtual form of cardiac rehabilitation and exercise training programs, and have demonstrated promising results.³⁰ Utilization of these technologies have been shown to better engage patients in the care process, leading to improved satisfaction with the healthcare system, and converts the patient from a passive recipient to an active partner on the healthcare team.³¹ This is in keeping with the Office of the National Coordinator's 2020 vision for health information technology: *the power of each individual is developed and unleashed to be active in managing their health and partnering in their health care, enabled by information and technology.*³² The opportunities afforded by apps and wearables will significantly expand the physician armamentarium, and provide a cost-effective, non-pharmacologic strategy that can create positive behavior change.

Chronic Disease Outcomes and Primary Care Services

Physician adherence to the current evidence base in the management of chronic disease is poor, and patients diagnosed with a chronic disease typically receive only half of the recommended process of care.^{33,34} In the case of the two most common chronic diseases impacting the population, fewer than 1 in 3 patients with hypertension and hypercholesterolemia attain control of both disorders.³⁴ These gaps in care have been subsequently shown to lead to higher clinical events and added health care costs. The causes for this deficiency in care are multifactorial, but are primarily due to four factors: physician time demands, rapidly expanding medical database, therapeutic inertia, and lack of supporting infrastructure.

The current U.S. model for delivery of chronic disease care typically rests on the back of the primary care physician, whose time for face-to-face patient care has become progressively constrained; it is estimated that direct patient care accounts for only 55% of the average workday.^{16,35} Studies evaluating the time necessary to achieve the recommendations of national practice guidelines for just 10 chronic diseases estimate that this alone would require 10.6 hours a day, more time than primary care physicians have available for patient care overall.³⁶

Compounding these data are the expected shortage of primary care physicians in the U.S.; by 2025 it is estimated that an additional 52,000 primary care physicians will be needed to care for the growing and aging population, yet there is little evidence to suggest that these needs will be met.³⁷

The second factor compromising chronic disease care is the rapidly evolving medical database, which has grown logarithmically in the last 4 decades. In the mid-1960s, there were approximately 100,000 peer-reviewed articles published in the medical literature per year. By 2012, there were 28,100 active scholarly peer-reviewed journals collectively publishing about 1.8-1.9 million articles a year.^{38,39} Further confounding the widening breadth of medical information is that a significant percentage of published studies contradict current medical practice, or what has been labeled a *medical reversal*.⁴⁰ The ability therefore to keep up with the current and accepted evidence base across the broad range of medical conditions comprising chronic disease is clearly a major challenge for any busy practicing clinician.

The third factor influencing poor chronic disease care is due to what has been labeled *therapeutic inertia*, which occurs when a provider fails to increase or modify therapy when treatment goals are

unmet.⁴¹ In uncontrolled hypertension, the prevalence of therapeutic inertia has been reported to be as high as 86.9% of visits when the blood pressure was $\geq 140/90$ mmHg.⁴² Failure to intensify therapy in patients with abnormal blood glucose, blood lipids, or blood pressure technically fits the definition of a medical error as defined by the Institute of Medicine, and contributes to the widespread failure to achieve evidence-based goals. The causes of therapeutic inertia are multiple, and involve the clinician, the patient, and the healthcare system (Table 1).

The fourth and final factor is the care model supporting the patient and physician including the reactive and episodic nature of care delivery.⁴³ Studies covering a variety of medical conditions consistently show that providing the primary care physician with a team-based infrastructure of specialized, non-physician caregivers whose role is to provide a continuous framework of monitoring and management, improves adherence to quality measures and yields superior outcomes, cost, and patient satisfaction.⁴⁴ Moreover, non-physician caregivers following evidence-based guidelines are less likely to be impeded by therapeutic inertia.^{35,45,46} Management of warfarin is an excellent case in point. When compared to physician-management, pharmacist-directed care resulted in the highest attainment of quality indicators and patient satisfaction while yielding the lowest adverse clinical events and cost.⁴⁷ Similarly, management of hypertension, chronic obstructive lung disease, diabetes and heart failure have also demonstrated superior process measures and clinical outcomes when supplementing the primary care physician with a supporting infrastructure of specialized, non-physician caregivers working in a focused factory model of care delivery.^{45,48,49}

Social Networks

The importance of social network influences on behavior is now well established, having demonstrated considerable impact on smoking, diet, exercise, depression, medication adherence and obesity.^{50,51} Decisions to quit smoking, begin an exercise program, as well as other health-related behaviors are not completely made by isolated persons but rather reflect choices made by groups connected to each other.⁵⁰ This influence can be extensive, often reaching up to three degrees of separation. Because patients are embedded within social networks suggest that both good and bad behaviors could spread over a range of social ties, and that network involvement in health improvement and/or disease management could lead to positive and sustainable effects

over time.⁵¹ A recent study in patients with either chronic heart disease or diabetes revealed that higher levels of social network involvement was linked to the maintenance of healthy behaviors over time, leading to reductions in hospitalizations and total cost of care.⁵² Social networks appear adaptable and responsive to levels of health need, thus harnessing and sustaining the capacity of these networks offers promise as a cost-effective way of supporting behavior change and long-term chronic disease management.^{52,53} Use of electronic communications and disease-centric social networks permit large-scale unobtrusive measures of network activity along with behavior change information that can accelerate improvement in health behaviors and disease management.⁵³ Recent innovations such as patient portals offer healthcare providers a new avenue of accessing large groups of patients and favorably influencing health behaviors.⁵⁴ Successful disease management strategies may utilize the potential of social networks in creating sustainable and cost-effective solutions for patients with chronic disease.

Infrastructure Requirements Going Forward

As healthcare moves from a volume-to-value strategy, the need to collect and manage data will continually increase, and with it, the necessity to provide analysis, data visualization tools and education/training surrounding the use of new technologies and the data they generate. Healthcare systems that develop both discipline and efficiency in data analytics and reporting (as well as the ability to train it's staff in the capabilities of these ever-evolving systems) will best manage both sides of the value equation. Real-time, metric-driven process improvement utilizing outcome measures will facilitate a continuously learning health care system (Table 2). Decision support software and knowledge management tools incorporated as key components of the delivery system can ensure that decisions are informed by the best evidence.⁵⁵ Payments should reward desired care outcomes with incentives to provide the best care at lower cost.^{44,55,56} Physicians will willingly utilize non-physician support services as long as such services deliver superior outcomes in an efficient, patient-focused, cost-effective manner.⁴⁷

Changing the current delivery model by incorporating a *generalized* care-team supporting the PCP (medical home model) has unfortunately demonstrated limited improvements in quality and overall outcomes.⁵⁷ More successful approaches have utilized *specialized* integrated practice units (IPUs), each employing non-physician personnel who are dedicated to a specific disease condition for the full cycle of care.^{56,58} Members of the care team may include pharmacists, advanced practice

clinicians, nurses, health educators, dieticians, social workers, counselors and therapists, all organized around the patient's medical condition. In this model, patients can be more frequently and effectively connected to the health delivery system utilizing apps as well as home-based and wearable devices, and communication can be consistent and at regular intervals between the care team and the patient.^{45,48,59} (Figure 1) IPU will have the capacity to care for the spectrum of patients within a disease category, but may concentrate greater efforts in high-risk patients who often consume the highest percentage of healthcare resources. Patients can achieve a higher level of engagement in the care process via enhanced education, real-time feedback via wearable and home-based devices, and enriched communication with both the care team and other patients via social networks, thus achieving a higher level of satisfaction with the healthcare system. This new delivery model is in keeping with current patient expectations; results of a recently released global study of healthcare suggest that individuals worldwide want to see their biological makeup and individual behaviors used to make receiving care more effective and efficient.⁶⁰ Systems and just-in-time communication will be capable of delivering customized care efficiently across broad populations with chronic disease cost-effectively when delivery systems are compensated using value-based payment models. Moreover, broadening the care team will reduce waste and improve primary care access and capacity at a time when the primary care workforce is diminishing.

Conclusion

Throughout history, the healthcare system has continually reengineered itself to meet the medical needs of the time. Isolation wards were created in the late 19th and early 20th centuries to meet the crisis in infectious disease, particularly during the typhoid and influenza epidemics, and mobile army surgical hospitals (MASH) were crafted in 1945 to better manage surgical emergencies during war. Today, healthcare must reengineer its care delivery model to manage the chief medical crisis of the 21st century, chronic disease. The capacity of the stand-alone physician to produce high-quality, evidenced-based care, yielding meaningful and lasting change in lifestyle behaviors has proven illusive. A new model of team-based care organized as an IPU, will have the ability to deliver comprehensive consistent treatment and advice using a focused-factory approach. The IPU will employ the latest in technology innovation, thus better engaging patients, in addition

to providing high-quality, consistent, personalized care delivery, and accelerate consequential lifestyle change.

ACCEPTED MANUSCRIPT

Bibliography

1. Heron M. Deaths: leading causes for 2010. *National vital statistics reports : from the Centers for Disease Control and Prevention, National Center for Health Statistics, National Vital Statistics System*. 2013;62(6):1-97.
2. Moses H, 3rd, Matheson DH, Dorsey ER, George BP, Sadoff D, Yoshimura S. The anatomy of health care in the United States. *JAMA : the journal of the American Medical Association*. 2013;310(18):1947-1963.
3. Wu S, Green A. *Projection of chronic illness prevalence and cost inflation*. RAND Corporation;2000.
4. Bodenheimer T, Chen E, Bennett HD. Confronting the growing burden of chronic disease: can the U.S. health care workforce do the job? *Health affairs*. 2009;28(1):64-74.
5. Thorpe KE. The rise in health care spending and what to do about it. *Health affairs*. 2005;24(6):1436-1445.
6. Dall TM, Gallo PD, Chakrabarti R, West T, Semilla AP, Storm MV. An aging population and growing disease burden will require a large and specialized health care workforce by 2025. *Health affairs*. 2013;32(11):2013-2020.
7. Schiller JS, Lucas JW, National Health Interview Survey (U.S.), National Center for Health Statistics (U.S.). Division of Health Interview Statistics. *Summary health statistics for U.S. adults : National Health Interview Survey, 2010 : data from the National Health Interview Survey*. Hyattsville, MD: U.S. Dept. of Health and Human Services, Centers for Disease Control and Prevention, National Center for Health Statistics; 2012.
8. Christ G, Diwan S. Chronic Illness and Aging. 2013; <http://www.cswe.org/File.aspx?id=25462>.
9. Hung WW, Ross JS, Boockvar KS, Siu AL. Recent trends in chronic disease, impairment and disability among older adults in the United States. *BMC geriatrics*. 2011;11:47.
10. Heidenreich PA, Trogon JG, Khavjou OA, et al. Forecasting the future of cardiovascular disease in the United States: a policy statement from the American Heart Association. *Circulation*. 2011;123(8):933-944.
11. Boyle JP, Thompson TJ, Gregg EW, Barker LE, Williamson DF. Projection of the year 2050 burden of diabetes in the US adult population: dynamic modeling of incidence, mortality, and prediabetes prevalence. *Population health metrics*. 2010;8:29.
12. Fryar CD, Carroll MD, Ogden CL. Prevalence of Overweight, Obesity, and Extreme Obesity Among Adults: United States, Trends 1960–1962 Through 2009–2010. *NCHS data brief*. 2012:1-8.
13. McGuire S. State Indicator Report on Fruits and Vegetables, 2013, Centers for Disease Control and Prevention, Atlanta, GA. *Advances in nutrition*. 2013;4(6):665-666.
14. Danaei G, Ding EL, Mozaffarian D, et al. The preventable causes of death in the United States: comparative risk assessment of dietary, lifestyle, and metabolic risk factors. *PLoS medicine*. 2009;6(4):e1000058.
15. Schroeder SA. Shattuck Lecture. We can do better--improving the health of the American people. *The New England journal of medicine*. 2007;357(12):1221-1228.
16. Gottschalk A, Flocke SA. Time spent in face-to-face patient care and work outside the examination room. *Annals of family medicine*. 2005;3(6):488-493.

17. Tai-Seale M, McGuire TG, Zhang W. Time allocation in primary care office visits. *Health services research*. 2007;42(5):1871-1894.
18. Thande NK, Hurstak EE, Sciacca RE, Giardina EG. Management of obesity: a challenge for medical training and practice. *Obesity*. 2009;17(1):107-113.
19. Rao G, Burke LE, Spring BJ, et al. New and emerging weight management strategies for busy ambulatory settings: a scientific statement from the American Heart Association endorsed by the Society of Behavioral Medicine. *Circulation*. 2011;124(10):1182-1203.
20. Milani RV, Lavie CJ. Impact of cardiac rehabilitation on depression and its associated mortality. *The American journal of medicine*. 2007;120(9):799-806.
21. House JS, Landis KR, Umberson D. Social relationships and health. *Science*. 1988;241(4865):540-545.
22. American Society of Consultant Pharmacists Fact Sheet. 2013; <https://http://www.ascp.com/articles/about-ascp/ascp-fact-sheet>. Accessed Jan, 2014.
23. Lucado J, Paez K, Elixhauser A. Medication-Related Adverse Outcomes in U.S. Hospitals and Emergency Departments, 2008: Statistical Brief #109. *Healthcare Cost and Utilization Project (HCUP) Statistical Briefs*. Rockville (MD)2011.
24. Spring B, Duncan JM, Janke EA, et al. Integrating technology into standard weight loss treatment: a randomized controlled trial. *JAMA internal medicine*. 2013;173(2):105-111.
25. Connelly J, Kirk A, Masthoff J, MacRury S. The use of technology to promote physical activity in Type 2 diabetes management: a systematic review. *Diabetic medicine : a journal of the British Diabetic Association*. 2013;30(12):1420-1432.
26. Fry JP, Neff RA. Periodic prompts and reminders in health promotion and health behavior interventions: systematic review. *Journal of medical Internet research*. 2009;11(2):e16.
27. Deci EL, Ryan RM. The "What" and "Why" of Goal Pursuits: Human Needs and the Self-Determination of Behavior. *Psychological Inquiry*. 2000;11(4):227-268.
28. Lefebvre RC, Bornkessel AS. Digital social networks and health. *Circulation*. 2013;127(17):1829-1836.
29. Ba S, Wang L. Digital health communities: The effect of their motivation mechanisms. *Decision Support Systems*. 2013;55(4):941-947.
30. Beatty AL, Fukuoka Y, Whooley MA. Using mobile technology for cardiac rehabilitation: a review and framework for development and evaluation. *Journal of the American Heart Association*. 2013;2(6):e000568.
31. Greene J, Hibbard JH, Sacks R, Overton V. When seeing the same physician, highly activated patients have better care experiences than less activated patients. *Health affairs*. 2013;32(7):1299-1305.
32. Atkinson NL, Olinger L, Pan E. *Unleashing the Power of Each Individual to Manage Their Health and Partner in Their Health Care, Enabled by Information and Technology*. Office of the National Coordinator for Health Information Technology. February 21, 2013.
33. McGlynn EA, Asch SM, Adams J, et al. The quality of health care delivered to adults in the United States. *The New England journal of medicine*. 2003;348(26):2635-2645.
34. Egan BM, Li J, Qanungo S, Wolfman TE. Blood pressure and cholesterol control in hypertensive hypercholesterolemic patients: national health and nutrition examination surveys 1988-2010. *Circulation*. 2013;128(1):29-41.
35. Yarnall KS, Ostbye T, Krause KM, Pollak KI, Gradison M, Michener JL. Family physicians as team leaders: "time" to share the care. *Preventing chronic disease*. 2009;6(2):A59.

36. Ostbye T, Yarnall KS, Krause KM, Pollak KI, Gradison M, Michener JL. Is there time for management of patients with chronic diseases in primary care? *Annals of family medicine*. 2005;3(3):209-214.
37. Petterson SM, Liaw WR, Phillips RL, Jr., Rabin DL, Meyers DS, Bazemore AW. Projecting US primary care physician workforce needs: 2010-2025. *Annals of family medicine*. 2012;10(6):503-509.
38. Druss BG, Marcus SC. Growth and decentralization of the medical literature: implications for evidence-based medicine. *Journal of the Medical Library Association : JMLA*. 2005;93(4):499-501.
39. Ware M. *The STM report. An overview of scientific and scholarly journal publishing*. 2012.
40. Prasad V, Cifu A, Ioannidis JP. Reversals of established medical practices: evidence to abandon ship. *JAMA : the journal of the American Medical Association*. 2012;307(1):37-38.
41. Milani RV, Lavie CJ. Lipid Control in the Modern Era: An Orphan's Tale of Rags to Riches. *Journal of the American College of Cardiology*. 2013;62(23):2185-2187.
42. Okonofua EC, Simpson KN, Jesri A, Rehman SU, Durkalski VL, Egan BM. Therapeutic inertia is an impediment to achieving the Healthy People 2010 blood pressure control goals. *Hypertension*. 2006;47(3):345-351.
43. Asch DA, Muller RW, Volpp KG. Automated hovering in health care--watching over the 5000 hours. *The New England journal of medicine*. 2012;367(1):1-3.
44. Thorpe KE. Team approach to care of chronic conditions is key to long-term health system fix. Interviewed by Lois A Bowers. *Medical economics*. 2013;90(1):58.
45. Margolis KL, Asche SE, Bergdall AR, et al. Effect of home blood pressure telemonitoring and pharmacist management on blood pressure control: a cluster randomized clinical trial. *JAMA : the journal of the American Medical Association*. 2013;310(1):46-56.
46. Horning KK, Hoehns JD, Doucette WR. Adherence to clinical practice guidelines for 7 chronic conditions in long-term-care patients who received pharmacist disease management services versus traditional drug regimen review. *Journal of managed care pharmacy : JMCP*. 2007;13(1):28-36.
47. Saokaew S, Permsuwan U, Chaiyakunapruk N, Nathisuwan S, Sukonthasarn A. Effectiveness of pharmacist-participated warfarin therapy management: a systematic review and meta-analysis. *Journal of thrombosis and haemostasis : JTH*. 2010;8(11):2418-2427.
48. Steventon A, Bardsley M, Billings J, et al. Effect of telehealth on use of secondary care and mortality: findings from the Whole System Demonstrator cluster randomised trial. *Bmj*. 2012;344:e3874.
49. Magid DJ, Green BB. Home blood pressure monitoring: take it to the bank. *JAMA : the journal of the American Medical Association*. 2013;310(1):40-41.
50. Christakis NA, Fowler JH. The collective dynamics of smoking in a large social network. *The New England journal of medicine*. 2008;358(21):2249-2258.
51. Christakis NA, Fowler JH. The spread of obesity in a large social network over 32 years. *The New England journal of medicine*. 2007;357(4):370-379.
52. Reeves D, Blickem C, Vassilev I, et al. The contribution of social networks to the health and self-management of patients with long-term conditions: a longitudinal study. *PloS one*. 2014;9(6):e98340.
53. Valente TW. Network interventions. *Science*. 2012;337(6090):49-53.

54. Urowitz S, Wiljer D, Dupak K, et al. Improving diabetes management with a patient portal: a qualitative study of diabetes self-management portal. *Journal of medical Internet research*. 2012;14(6):e158.
55. Smith MD, Institute of Medicine (U.S.). Committee on the Learning Health Care System in America. *Best care at lower cost : the path to continuously learning health care in America*. Washington, D.C.: National Academies Press; 2013.
56. Porter ME. A strategy for health care reform--toward a value-based system. *The New England journal of medicine*. 2009;361(2):109-112.
57. Friedberg MW, Schneider EC, Rosenthal MB, Volpp KG, Werner RM. Association between participation in a multipayer medical home intervention and changes in quality, utilization, and costs of care. *JAMA : the journal of the American Medical Association*. 2014;311(8):815-825.
58. Porter ME, Pabo EA, Lee TH. Redesigning primary care: a strategic vision to improve value by organizing around patients' needs. *Health affairs*. 2013;32(3):516-525.
59. Magid DJ, Olson KL, Billups SJ, Wagner NM, Lyons EE, Kroner BA. A pharmacist-led, American Heart Association Heart360 Web-enabled home blood pressure monitoring program. *Circulation. Cardiovascular quality and outcomes*. 2013;6(2):157-163.
60. *Global Innovation Barometer*. Intel Corporation. December, 2013.

Table 1: Factors Leading to Therapeutic Inertia

Clinician	Patient	Health System
Failure to initiate treatment	Medication side effects	Lack of clinical guideline
Failure to titrate to goal	Too many medications	Lack of care coordination
Failure to set clear goals	Cost of medications	No visit planning
Underestimation of patient need	Denial of disease	Lack of decision support
Failure to identify and manage comorbid conditions (such as depression)	Denial of disease severity	Poor communication between physician and office staff
Insufficient time	Forgetfulness	No disease registry
Insufficient focus or emphasis on goal attainment	Perception of low susceptibility	No active outreach
Reactive rather than proactive	Absence of disease symptoms	Perverse incentives
	Poor communication with physician	
	Mistrust of clinician	
	Depression, mental illness, substance abuse	
	Low health literacy	

Adapted from: Milani RV, Lavie CJ. JACC 2013; 62:2185-7.⁴¹

Table 2: Characteristics of a Continuously Learning Health Care System

Science and Informatics

- Real-time access to knowledge
- Digital capture of the care experience

Patient-Clinician Relationships

- Engaged, empowered patients

Incentives


- Incentives aligned for value
- Full transparency

Culture

- Leadership-instilled culture of learning
- Supportive system competencies

Adapted from reference 55

Figure 1: Models of Care Delivery in Chronic Disease Management



- Chronic disease is responsible for 75% of total health care costs and the majority of deaths in the United States.
- Existing delivery models are poorly constructed to manage chronic disease as evidenced by low adherence to quality and control indicators.
- New technologies have emerged that can engage patients and offer additional modalities in treating chronic disease.
- Modifying healthcare delivery to include team-based care combined with patient-centered technologies offer great promise.