

MATHEMATICAL SCIENCES AND APPLICATIONS E-NOTES

4 (1) 31-39 (2016) ©MSAEN

A Discretization of the Hadamard fractional derivative

Ricardo Almeida* and Nuno R. O. Bastos

(Communicated by Murat TOSUN)

Abstract

We present a new discretization for the Hadamard fractional derivative, that simplifies the computations. We then apply the method to solve a fractional differential equation and a fractional variational problem with dependence on the Hadamard fractional derivative.

Keywords: fractional calculus, discretization methods.

AMS Subject Classification (2010): Primary: 26A33; Secondary: 49M25, 49M25.

*Corresponding author

1. Introduction

Similarly to ordinary calculus, we can find in the literature distinct definitions for fractional derivatives and for fractional integrals, which are generalizations of the integer-order derivatives and multiple integrals, respectively. The most common ones and consequently more studied are the Riemann–Liouville, Caputo and Grünwald–Letnikov definitions. We deal in this paper with the Hadamard fractional derivative, introduced in [4]. Recently, it has call the attention of researchers and numerous results have appeared, with an extensive study of properties of such kind of operators [1, 2, 8]. For recent results we suggest [5, 10, 11].

Due the complexity of solving equations involving fractional operators, in most cases is impossible to determine the exact solution and so numerical methods are used to determine an approximated solution of the problem. This is an emerging field, and we can find already in the literature several methods to deal with these problems, at least for the most common fractional derivative types. For the Hadamard fractional derivative, we mention the recent paper [9], where the fractional operator is replaced by a finite sum involving only integer-order derivatives of the function. Replacing the fractional derivative by this sum, we rewrite the initial problem in terms of integer-order derivatives and thus we are able to apply classical known methods. In [3] another approximation formula is obtained, using also integer-order derivatives only. The disadvantage is

Received : 25–September–2015, *Accepted* : 14–January–2016

This article is the written version of author's plenary talk delivered on September 03- August 31, 2015 at 4th International Eurasian Conference on Mathematical Sciences and Applications (IECMSA-2015) in Athens, Greece.

that in order to have a good approximation, we need to use higher-order derivatives, which may not be adequate for fractional problems. In this paper we follow a different path, by discretizing the fractional derivative, and then convert continuous problems into discrete ones.

To start, let us recall the definition of the Hadamard fractional derivative.

Definition 1.1. Let a, b be two reals with $0 < a < b$, and $x : [a, b] \rightarrow \mathbb{R}$ be a function. For $\alpha \in (0, 1)$, the left Hadamard fractional derivative of order α is defined by

$${}_a\mathcal{D}_t^\alpha x(t) = \frac{t}{\Gamma(1-\alpha)} \frac{d}{dt} \int_a^t \left(\ln \frac{t}{\tau}\right)^{-\alpha} \frac{x(\tau)}{\tau} d\tau,$$

while the right Hadamard fractional derivative of order α by

$${}_t\mathcal{D}_b^\alpha x(t) = \frac{-t}{\Gamma(1-\alpha)} \frac{d}{dt} \int_t^b \left(\ln \frac{\tau}{t}\right)^{-\alpha} \frac{x(\tau)}{\tau} d\tau,$$

where Γ denotes the Gamma function.

When x is an absolutely continuous function, there exists an equivalent definition (cf. [6])

$${}_a\mathcal{D}_t^\alpha x(t) = \frac{x(a)}{\Gamma(1-\alpha)} \left(\ln \frac{t}{a}\right)^{-\alpha} + \frac{1}{\Gamma(1-\alpha)} \int_a^t \left(\ln \frac{t}{\tau}\right)^{-\alpha} \dot{x}(\tau) d\tau,$$

and

$${}_t\mathcal{D}_b^\alpha x(t) = \frac{x(b)}{\Gamma(1-\alpha)} \left(\ln \frac{b}{t}\right)^{-\alpha} - \frac{1}{\Gamma(1-\alpha)} \int_t^b \left(\ln \frac{\tau}{t}\right)^{-\alpha} \dot{x}(\tau) d\tau.$$

More properties can be found in references at the end. The paper is organized in the following way. In Section 2 we present the main result of the paper. Starting with the definition, and with an appropriate grid on time, we present a new discrete version for the left Hadamard fractional derivative. To show the efficiency of the method, in Section 3 we compare the exact expression of a fractional derivative with some numerical experiments, for different values of α and different step sizes n . In Section 4 we apply the technique to solve a fractional differential equation and a fractional calculus of variation problem.

2. The discretization method

The discretization method is described in the following way. Given a function $x : [a, b] \rightarrow \mathbb{R}$, fix a positive integer n , and define the time step

$$\Delta T = \frac{\ln \frac{b}{a}}{n}.$$

Given $N \in \{0, 1, \dots, n\}$, denote the time and space grid by

$$t_N = a \exp(N\Delta T) = a \sqrt[n]{\left(\frac{b}{a}\right)^N} \quad \text{and} \quad x_N = x(t_N).$$

Theorem 2.1. Let $x : [a, b] \rightarrow \mathbb{R}$ be a function of class C^2 and $n \in \mathbb{N}$. Denote

$$\psi = \frac{(\Delta T)^{1-\alpha}}{a(1 - \exp(-\Delta T))\Gamma(2-\alpha)} \quad \text{and} \quad (\omega_k^\alpha) = k^{1-\alpha} - (k-1)^{1-\alpha}.$$

Then, for all $N \in \{1, \dots, n\}$,

$${}_a\mathcal{D}_{t_N}^\alpha x_N = {}_a\tilde{\mathcal{D}}_{t_N}^\alpha x_N + O(\Delta T),$$

where

$${}_a\tilde{\mathcal{D}}_{t_N}^\alpha x_N = \frac{x(a)}{\Gamma(1-\alpha)} \left(\ln \frac{t_N}{a}\right)^{-\alpha} + \psi \sum_{k=1}^N (\omega_{N-k+1}^\alpha) \frac{x_k - x_{k-1}}{\exp(k\Delta T)} \cdot t_k, \quad (2.1)$$

and

$$\lim_{\Delta T \rightarrow 0} O(\Delta T) = 0.$$

Proof.

$$\begin{aligned} {}_a\mathcal{D}_{t_N}^\alpha x_N &= \frac{x(a)}{\Gamma(1-\alpha)} \left(\ln \frac{t_N}{a}\right)^{-\alpha} + \frac{1}{\Gamma(1-\alpha)} \int_a^{t_N} \left(\ln \frac{t_N}{\tau}\right)^{-\alpha} \dot{x}(\tau) d\tau \\ &= \frac{x(a)}{\Gamma(1-\alpha)} \left(\ln \frac{t_N}{a}\right)^{-\alpha} + \frac{1}{\Gamma(1-\alpha)} \sum_{k=1}^N \int_{t_{k-1}}^{t_k} \left(\ln \frac{t_N}{\tau}\right)^{-\alpha} \frac{1}{\tau} (\dot{x}(\tau)\tau) d\tau \\ &\approx \frac{x(a)}{\Gamma(1-\alpha)} \left(\ln \frac{t_N}{a}\right)^{-\alpha} \\ &\quad + \frac{1}{\Gamma(1-\alpha)} \sum_{k=1}^N \int_{t_{k-1}}^{t_k} \left(\ln \frac{t_N}{\tau}\right)^{-\alpha} \frac{1}{\tau} \left(\frac{x_k - x_{k-1}}{a(1 - \exp(-\Delta T)) \exp(k\Delta T)} \cdot t_k \right) d\tau \\ &= \frac{x(a)}{\Gamma(1-\alpha)} \left(\ln \frac{t_N}{a}\right)^{-\alpha} \\ &\quad + \frac{1}{a(1 - \exp(-\Delta T))\Gamma(1-\alpha)} \sum_{k=1}^N \frac{x_k - x_{k-1}}{\exp(k\Delta T)} \cdot t_k \int_{t_{k-1}}^{t_k} \left(\ln \frac{t_N}{\tau}\right)^{-\alpha} \frac{1}{\tau} d\tau \\ &= \frac{x(a)}{\Gamma(1-\alpha)} \left(\ln \frac{t_N}{a}\right)^{-\alpha} \\ &\quad + \frac{(\Delta T)^{1-\alpha}}{a(1 - \exp(-\Delta T))\Gamma(2-\alpha)} \sum_{k=1}^N \frac{x_k - x_{k-1}}{\exp(k\Delta T)} \cdot t_k [(N-k+1)^{1-\alpha} - (N-k)^{1-\alpha}]. \end{aligned}$$

Thus, we get the desired approximation formula. Now, let us determine an upper bound for the error when we use formula (2.1). The error is given by

$$E = \frac{1}{\Gamma(1-\alpha)} \sum_{k=1}^N \int_{t_{k-1}}^{t_k} \left(\ln \frac{t_N}{\tau}\right)^{-\alpha} \frac{1}{\tau} \left| \dot{x}(\tau)\tau - \frac{x_k - x_{k-1}}{t_k - t_{k-1}} \cdot t_k \right| d\tau.$$

Let

$$M_i = \max_{\tau \in [a, b]} |x^{(i)}(\tau)|, \quad i = 1, 2.$$

Then, using Taylor's Theorem, we get that, for all $k \in \{1, \dots, N\}$ and for all $\tau \in [t_{k-1}, t_k]$,

$$\begin{aligned} &\left| \dot{x}(\tau)\tau - \frac{x_k - x_{k-1}}{t_k - t_{k-1}} \cdot t_k \right| \\ &= \left| \dot{x}(\tau)\tau - \left(\dot{x}(t_{k-1}) + \ddot{x}(\xi_1) \frac{t_k - t_{k-1}}{2} \right) \cdot t_k \right| \\ &\leq |\dot{x}(\tau)\tau - \dot{x}(t_{k-1})t_k| + M_2 \frac{t_k - t_{k-1}}{2} \cdot t_k \\ &= |(\dot{x}(t_{k-1}) + \ddot{x}(\xi_2)(\tau - t_{k-1}))\tau - \dot{x}(t_{k-1})t_k| + M_2 \frac{t_k - t_{k-1}}{2} \cdot t_k \\ &\leq M_1(t_k - \tau) + M_2(\tau - t_{k-1})\tau + M_2 \frac{t_k - t_{k-1}}{2} \cdot t_k \\ &\leq (t_k - t_{k-1}) \left[M_1 + \frac{3}{2}M_2b \right]. \end{aligned}$$

Therefore, the error is bounded by

$$\begin{aligned} E &\leq \frac{1}{\Gamma(1-\alpha)} \sum_{k=1}^N \int_{t_{k-1}}^{t_k} \left(\ln \frac{t_N}{\tau}\right)^{-\alpha} \frac{1}{\tau} (t_k - t_{k-1}) \left[M_1 + \frac{3}{2}M_2b \right] d\tau \\ &= \frac{M_1 + \frac{3}{2}M_2b}{\Gamma(1-\alpha)} \sum_{k=1}^N (t_k - t_{k-1}) \int_{t_{k-1}}^{t_k} \left(\ln \frac{t_N}{\tau}\right)^{-\alpha} \frac{1}{\tau} d\tau. \end{aligned}$$

Having into consideration that, for all $k \in \{1, \dots, N\}$ and for all $\tau \in [t_{k-1}, t_k)$,

$$0 \leq \left(\ln \frac{t_N}{\tau} \right)^{-\alpha} \leq \left(\ln \frac{t_k}{\tau} \right)^{-\alpha},$$

we have that

$$0 \leq \int_{t_{k-1}}^{t_k} \left(\ln \frac{t_N}{\tau} \right)^{-\alpha} \frac{1}{\tau} d\tau \leq \frac{(\Delta T)^{1-\alpha}}{1-\alpha}.$$

Also, since

$$t_k - t_{k-1} = a(1 - \exp(-\Delta T)) \exp(k\Delta T),$$

then

$$\begin{aligned} E &\leq \frac{M_1 + \frac{3}{2}M_2b}{\Gamma(2-\alpha)} a(1 - \exp(-\Delta T)) (\Delta T)^{1-\alpha} \sum_{k=1}^N \exp(k\Delta T) \\ &= \frac{M_1 + \frac{3}{2}M_2b}{\Gamma(2-\alpha)} a(\Delta T)^{1-\alpha} (\exp(N\Delta T) - 1) \\ &\leq \frac{M_1 + \frac{3}{2}M_2b}{\Gamma(2-\alpha)} a(\Delta T)^{1-\alpha} \left(\frac{b}{a} - 1 \right). \end{aligned}$$

In conclusion, we obtain the upper bound formula for our approximation (2.1):

$$E \leq \frac{M_1 + \frac{3}{2}M_2b}{\Gamma(2-\alpha)} (b-a) (\Delta T)^{1-\alpha} \quad (2.2)$$

which converges to zero as $\Delta T \rightarrow 0$. □

In opposite to the classical case, where the concept of derivative is local, a fractional derivative contains memory, and thus to compute the approximation obtained in Eq. (2.1) at a point t_N , we need to know the values of $x(t_n)$ from the beginning until the end-point, i.e., from $n = 0$ to $n = N$.

For the right Hadamard fractional derivative, we have in a similar way the following approximation formula:

$${}_{t_N} \mathcal{D}_b^\alpha x_N \approx \frac{x(b)}{\Gamma(1-\alpha)} \left(\ln \frac{b}{t_N} \right)^{-\alpha} - \psi \sum_{k=N+1}^n (\omega_{k-N}^\alpha) \frac{x_k - x_{k-1}}{\exp(k\Delta T)} \cdot t_k.$$

3. Example

Let $x(t) = \ln t$, for $t \in [1, 2]$. Then (see [7])

$${}_1 \mathcal{D}_t^\alpha x(t) = \frac{(\ln t)^{1-\alpha}}{\Gamma(2-\alpha)}.$$

In Figure 1 we show the accuracy of the procedure, for different values of $\alpha \in \{0.2, 0.5, 0.7, 0.9\}$ and for different values of $n \in \{10, 30, 50\}$. The error of the numerical experiments is measured using the norm

$$d(x, y) = \frac{\sum_{k=1}^n |x_k - y_k|}{n}. \quad (3.1)$$

We can see that, for a greater value of n , the error decreases as expected.




Figure 1. Analytic vs. numerical approximation.

4. Applications

Example 4.1. Consider a fractional differential equation with dependence on the left Hadamard fractional derivative:

$$\begin{cases} f(t, x(t), {}_a\mathcal{D}_t^\alpha x(t)) = 0, & t \in [a, b] \\ x(a) = x_a. \end{cases}$$

The procedure to solve numerically the system is described next. Fix $n \in \mathbb{N}$ and for $N \in \{1, \dots, n\}$, define

$$t_N = a \exp(N\Delta T), \quad x_N = x(t_N) \quad \text{with} \quad \Delta T = \frac{\ln \frac{b}{a}}{n}.$$

Replacing the fractional operator by the approximation given in Eq. (2.1), we obtain a classical difference equation with n unknown points x_1, \dots, x_n ,

$$\begin{cases} f(t_N, x_N, {}_a\tilde{\mathcal{D}}_{t_N}^\alpha x_N) = 0, & N \in \{1, \dots, n\} \\ x_0 = x_a. \end{cases}$$

For example, consider the system

$$\begin{cases} {}_1\mathcal{D}_t^\alpha x(t) + x(t) = \frac{(\ln t)^{1-\alpha}}{\Gamma(2-\alpha)} + \ln t, & t \in [1, 2] \\ x(1) = 0. \end{cases}$$

The obvious solution is $\bar{x}(t) = \ln t$. Applying the discussed method, we obtain

$$\begin{cases} \psi \sum_{k=1}^N (\omega_{N-k+1}^\alpha) \frac{x_k - x_{k-1}}{\exp(k\Delta T)} \cdot t_k + x_N = \frac{(\ln t_N)^{1-\alpha}}{\Gamma(2-\alpha)} + \ln t_N, & N \in \{1, \dots, n\} \\ x_0 = 0, \end{cases}$$

with

$$\Delta T = \frac{\ln 2}{n}, \quad t_N = \exp(N\Delta T), \quad \psi = \frac{(\Delta T)^{1-\alpha}}{(1 - \exp(-\Delta T))\Gamma(2-\alpha)} \quad \text{and} \quad \omega_k = k^{1-\alpha} - (k-1)^{1-\alpha}.$$

In Figure 2 we show the numerical results, for different values of $\alpha \in \{0.2, 0.5, 0.7, 0.9\}$ and for different values of $n \in \{5, 15, 30\}$.




Figure 2. Analytic and approximated solutions for problem of Example 4.1.

Example 4.2. For our next application, we show how to solve fractional variational problems with the Lagrangian depending on the Hadamard fractional derivative. Consider the functional

$$J(x) = \int_a^b L(t, x(t), {}_a\mathcal{D}_t^\alpha x(t)) dt,$$

on the set on functions that satisfy the boundary conditions

$$x(a) = x_a \quad \text{and} \quad x(b) = x_b,$$

where x_a, x_b are two fixed reals. The procedure how to find a numerical approximation is explained next. First, divide the interval $[a, b]$ into n subintervals $[t_{N-1}, t_N]$ for $N \in \{1, \dots, n\}$, where

$$t_N = a \exp(N \Delta T) \quad \text{and} \quad \Delta T = \frac{\ln \frac{b}{a}}{n}.$$

Denoting $x_N = x(t_N)$, applying the trapezoidal rule and taking into consideration Eq. (2.1), the variational integral is approximated as

$$\begin{aligned} J(x) &= \sum_{N=1}^n \int_{t_{N-1}}^{t_N} f(t, x(t), {}_a\mathcal{D}_t^\alpha x(t)) dt \\ &\approx \frac{f(t_0, x_0, {}_a\mathcal{D}_{t_0}^\alpha x_0)(t_1 - t_0) + f(t_n, x_n, {}_a\mathcal{D}_{t_n}^\alpha x_n)(t_n - t_{n-1})}{2} \\ &\quad + \sum_{N=1}^{n-1} \frac{f(t_N, x_N, {}_a\mathcal{D}_{t_N}^\alpha x_N)(t_{N+1} - t_{N-1})}{2} \\ &\approx \frac{f(a, x_a, {}_a\tilde{\mathcal{D}}_{t_1}^\alpha x_1)(t_1 - t_0) + f(t_n, x_n, {}_a\tilde{\mathcal{D}}_{t_n}^\alpha x_n)(t_n - t_{n-1})}{2} \\ &\quad + \sum_{N=1}^{n-1} \frac{f(t_N, x_N, {}_a\tilde{\mathcal{D}}_{t_N}^\alpha x_N)(t_{N+1} - t_{N-1})}{2}. \end{aligned}$$

Observe that we used here the approximation

$${}_a\mathcal{D}_a^\alpha x_a \approx {}_a\tilde{\mathcal{D}}_{t_1}^\alpha x_1.$$

We can regard this sum as a function of $n - 1$ unknown variables $\Psi(x_1, \dots, x_{n-1})$, and then to find the optimal solution one needs to solve the system

$$\frac{\partial \Psi}{\partial x_N} = 0, \quad \text{for } N \in \{1, \dots, n-1\},$$

and with this we track the desired values (x_1, \dots, x_{n-1}) . Observe that, in opposite to the classical case, $\partial \Psi / \partial x_N$ depends on the points $x_N, x_{N+1}, \dots, x_{n-1}$.

For example, we want the global minimizer for

$$J(x) = \int_1^2 \left({}_1\mathcal{D}_t^\alpha x(t) - \frac{(\ln t)^{1-\alpha}}{\Gamma(2-\alpha)} \right)^2 dt,$$

with the restrictions

$$x(1) = 0 \quad \text{and} \quad x(2) = \ln 2.$$

The optimal solution is $\bar{x}(t) = \ln t$ since the functional takes only non-negative values and vanishes when evaluated at \bar{x} .

In Figure 3 we show the solution of the problem, for different values of $\alpha \in \{0.2, 0.5, 0.7, 0.9\}$ and for different values of $n \in \{5, 15, 30\}$.

5. Conclusions

For all numerical experiments presented above we used MatLab to obtain the results. In Examples 4.1 and 4.2, when we take a small number of mesh points ($n = 5$) we get a not so good solution. However, increasing the value of n , the error decreases and the numerical solution approaches the analytic solution, converging to it. From the numerical results we also notice that, for the same values of n , as α increases the error also increases, which makes sense taking into account formula (2.2). We fix all parameters except α , it is easy to check that the maximum for error value increases as we increase the value of α .




Figure 3. Analytic and approximated solutions for problem of Example 4.2.

	α	n	E	n	E	n	E
Example 4.1	0.2	5	0.013201	15	0.0044804	30	0.0022501
	0.5	5	0.015074	15	0.005199	30	0.002621
	0.7	5	0.016174	15	0.0056458	30	0.0028553
	0.9	5	0.017093	15	0.0060434	30	0.003068
Example 4.2	0.2	5	0.018783	15	0.0081261	30	0.0041641
	0.5	5	0.026079	15	0.013255	30	0.0066303
	0.7	5	0.037751	15	0.021007	30	0.01066
	0.9	5	0.058731	15	0.031629	30	0.016684

Table 1. Number of mesh points, n , with corresponding error, E from formula (3.1).

Acknowledgments

Work supported by Portuguese funds through the CIDMA - Center for Research and Development in Mathematics and Applications, and the Portuguese Foundation for Science and Technology (FCT-Fundação para a Ciência e a Tecnologia), within project UID/MAT/04106/2013.

References

- [1] Butzer, P.L., Kilbas, A. A. and Trujillo, J. J., Mellin transform analysis and integration by parts for Hadamard-type fractional integrals, *J. Math. Anal. Appl.* 270 (2002), no. 1, 1-15.
- [2] Butzer, P.L., Kilbas, A. A. and Trujillo, J. J., Fractional calculus in the Mellin setting and Hadamard-type fractional integrals, *J. Math. Anal. Appl.* 269 (2002), no. 1, 1-27.
- [3] Butzer, P.L., Kilbas, A. A. and Trujillo, J. J., Stirling functions of the second kind in the setting of difference and fractional calculus, *Numer. Funct. Anal. Optim.* 24 (2003), no. 7-8, 673-711.
- [4] Hadamard, J., Essai sur l'étude des fonctions données par leur développement de Taylor, *J. Pure Appl. Math.* 4 (1892), no. 8, 101-186.
- [5] Jarad, F., Abdeljawad, T. and Baleanu, D., Caputo-type modification of the Hadamard fractional derivatives, *Advances in Difference Equations* August 2012 (2012), 2012–142.
- [6] Kilbas, A.A., Hadamard-type fractional calculus, *J. Korean Math. Soc.* 38 (2001), no. 6, 1191-1204.
- [7] Kilbas, A.A., Srivastava, H.M. and Trujillo, J.J., Theory and applications of fractional differential equations. North-Holland Mathematics Studies, 204. Elsevier Science B.V., Amsterdam, 2006.
- [8] Kilbas, A.A. and Titioura, A.A., Nonlinear differential equations with Marchaud-Hadamard-type fractional derivative in the weighted space of summable functions, *Math. Model. Anal.* 12 (2007), no. 3, 343-356.
- [9] Pooseh, S., Almeida, R. and Torres, D. F. M., Expansion formulas in terms of integer-order derivatives for the Hadamard fractional integral and derivative, *Numer. Funct. Anal. Optim.* 33 (2012), no. 3, 301–319.
- [10] Qassim, M. D., Furati, K. M. and Tatar, N.E., On a Differential Equation Involving Hilfer-Hadamard Fractional Derivative, *Abstract and Applied Analysis* vol. 2012 (2012), Article ID 391062, 17 pages, doi:10.1155/2012/391062.
- [11] Qian, D., Gong, Z. and Li, C., A generalized Gronwall inequality and its application to fractional differential equations with Hadamard derivatives, 3rd Conference on Nonlinear Science and Complexity (NSC10), Cankaya University, Ankara, Turkey, 28–31 July, 2010.

Affiliations

RICARDO ALMEIDA

ADDRESS: Center for Research and Development in Mathematics and Applications (CIDMA)
Department of Mathematics, University of Aveiro, 3810–193 Aveiro, Portugal.

E-MAIL: ricardo.almeida@ua.pt

NUNO R. O. BASTOS

ADDRESS: Centre for the Study of Education, Technologies and Health (CI&DETS), Department
of Mathematics, School of Technology and Management of Viseu
Polytechnic Institute of Viseu, 3504–510 Viseu, Portugal.

E-MAIL: nbastos@estv.ipv.pt