

PROCEEDINGS OF THE
 AMERICAN MATHEMATICAL SOCIETY
 Volume 00, Number 0, Pages 000–000
 S 0002-9939(XX)0000-0

A REMARK ON SCHATTEN CLASS TOEPLITZ OPERATORS ON BERGMAN SPACES

JORDI PAU

ABSTRACT. We provide a new characterization (valid for all $0 < p < \infty$) of Schatten class membership of Toeplitz operators induced by positive measures acting on weighted Bergman spaces on the unit ball. The result is new even for the case of the unit disk.

Let \mathbb{C}^n denote the Euclidian space of complex dimension n . For any two points $z = (z_1, \dots, z_n)$ and $w = (w_1, \dots, w_n)$ in \mathbb{C}^n we write $\langle z, w \rangle = z_1 \bar{w}_1 + \dots + z_n \bar{w}_n$, and $|z| = \sqrt{\langle z, z \rangle} = \sqrt{|z_1|^2 + \dots + |z_n|^2}$. The set $\mathbb{B}_n = \{z \in \mathbb{C}^n : |z| < 1\}$ is the open unit ball in \mathbb{C}^n . Denote by dv the usual Lebesgue volume measure on \mathbb{B}_n , normalized so that the volume of \mathbb{B}_n is one. Throughout this paper we fix a real parameter α with $\alpha > -1$ and write $dv_\alpha(z) = c_\alpha (1 - |z|^2)^\alpha dv(z)$, where c_α is a positive constant chosen so that $v_\alpha(\mathbb{B}_n) = 1$.

The weighted Bergman space $A_\alpha^2(\mathbb{B}_n)$ consists of those functions f holomorphic on \mathbb{B}_n that are in the Lebesgue space $L^2(\mathbb{B}_n, dv_\alpha)$. It is a Hilbert space with inner product

$$\langle f, g \rangle_\alpha = \int_{\mathbb{B}_n} f(z) \overline{g(z)} dv_\alpha(z).$$

The corresponding norm will be denoted by $\|f\|_\alpha$. The orthogonal projection $P_\alpha : L^2(\mathbb{B}_n, dv_\alpha) \rightarrow A_\alpha^2(\mathbb{B}_n)$ is an integral operator given by

$$P_\alpha f(z) = \int_{\mathbb{B}_n} \frac{f(w) dv_\alpha(w)}{(1 - \langle z, w \rangle)^{n+1+\alpha}}, \quad f \in L^2(\mathbb{B}_n, dv_\alpha).$$

Given a finite positive Borel measure μ on \mathbb{B}_n , the Toeplitz operator T_μ with symbol μ acting on $A_\alpha^2(\mathbb{B}_n)$ is the densely defined integral operator

$$T_\mu f(z) = \int_{\mathbb{B}_n} \frac{f(w) d\mu(w)}{(1 - \langle z, w \rangle)^{n+1+\alpha}}, \quad f \in A_\alpha^2(\mathbb{B}_n).$$

The operator T_μ is well defined on $H^\infty(\mathbb{B}_n)$, the algebra of all bounded holomorphic functions in \mathbb{B}_n . It is easy to check that $H^\infty(\mathbb{B}_n)$ is dense in $A_\alpha^2(\mathbb{B}_n)$.

For $z \in \mathbb{B}_n$, the Berezin transform of μ is

$$\tilde{\mu}(z) = \int_{\mathbb{B}_n} \frac{(1 - |z|^2)^{n+1+\alpha}}{|1 - \langle z, w \rangle|^{2(n+1+\alpha)}} d\mu(w).$$

2010 *Mathematics Subject Classification.* 32A36, 47B10, 47B35.

Key words and phrases. Bergman spaces, Toeplitz operators, Schatten classes.

The author is supported by SGR grant 2009SGR 420 (Generalitat de Catalunya) and DGICYT grant MTM2011-27932-C02-01 (MCyT/MEC).

We also need another averaging function. For $z \in \mathbb{B}_n$ and $r > 0$ define

$$\widehat{\mu}_r(z) = \frac{\mu(D(z, r))}{v_\alpha(D(z, r))},$$

where $D(z, r) = \{w \in \mathbb{B}_n : \beta(z, w) < r\}$ is the Bergman metric ball at z with radius r (here $\beta(z, w)$ denotes the Bergman or hyperbolic metric). The boundedness and compactness of T_μ on $A_\alpha^2(\mathbb{B}_n)$ for positive μ , can be described using the averaging functions $\widehat{\mu}_r$ and $\widetilde{\mu} : T_\mu$ is bounded on $A_\alpha^2(\mathbb{B}_n)$ if and only if $\widetilde{\mu}$ is bounded on \mathbb{B}_n if and only if $\widehat{\mu}_r$ is bounded on \mathbb{B}_n ; and T_μ is compact on $A_\alpha^2(\mathbb{B}_n)$ if and only if $\widetilde{\mu} \in C_0(\mathbb{B}_n)$ if and only if $\widehat{\mu}_r \in C_0(\mathbb{B}_n)$ (see [5]). Here $C_0(\mathbb{B}_n)$ is the space of all complex-valued continuous functions f on \mathbb{B}_n such that $f(z) \rightarrow 0$ as $|z| \rightarrow 1^-$.

Concerning the membership of T_μ in the Schatten-Von Neumann ideals S_p , there is a complete description using the averaging function $\widehat{\mu}_r$, but a description using the Berezin transform $\widetilde{\mu}$ can only be obtained for a certain ranges of p . We recall that, for $0 < p < \infty$, a compact operator T acting on a separable Hilbert space H belongs to the Schatten class S_p if its sequence of singular numbers belongs to the sequence space ℓ^p (the singular numbers are the square roots of the eigenvalues of the positive operator T^*T , where T^* is the Hilbert adjoint of T). We refer to [7, Chapter 1] for a brief account on Schatten classes. Part of the result stated below is due to D. Luecking [2] when $n = 1$, and the general form is due to K. Zhu [8].

Theorem A. *Let $0 < p < \infty$, $r > 0$ and μ be a finite positive Borel measure on \mathbb{B}_n . Then*

- (a) T_μ is in S_p if and only if $\widehat{\mu}_r \in L^p(\mathbb{B}_n, d\lambda)$.
- (b) If $p > n/(n + 1 + \alpha)$, then T_μ is in S_p if and only if $\widetilde{\mu} \in L^p(\mathbb{B}_n, d\lambda)$.

Here

$$d\lambda(z) = \frac{dv(z)}{(1 - |z|^2)^{n+1}}$$

is the Möbius invariant volume measure on \mathbb{B}_n . The cut-off point $n/(n + 1 + \alpha)$ in part (b) is best possible [8], and it is related to the finiteness of the measure $(1 - |z|^2)^{p(n+1+\alpha)}d\lambda(z)$. One could ask if it is possible to obtain a description in terms of some other integral transform (similar to the Berezin transform) being valid for all $0 < p < \infty$. The positive answer given below is the main result of the paper.

Theorem 1. *Let $\alpha > -1$, $0 < p < \infty$, and μ be a finite positive Borel measure on \mathbb{B}_n . Then T_μ is in the Schatten class S_p of $A_\alpha^2(\mathbb{B}_n)$ if and only if, for each (or some) $t \geq 0$ with $p(n + 1 + \alpha + 2t) > n$, the function*

$$(B^t\mu)(z) = \int_{\mathbb{B}_n} \frac{(1 - |z|^2)^{n+1+\alpha+2t}}{|1 - \langle z, w \rangle|^{2(n+1+\alpha+t)}} d\mu(w)$$

belongs to $L^p(\mathbb{B}_n, d\lambda)$.

The necessity of the condition $B^t\mu \in L^p(\mathbb{B}_n, d\lambda)$ follows easily from Theorem A, so that the interesting case is the sufficiency, especially for small p . Throughout the paper, constants are often given without computing their exact values, and the value of a constant C may change from one occurrence to the next.

We need the following well known integral estimate that has become very important in this area of analysis (see [6, Theorem 1.12] for example).

Lemma B. *Let $t > -1$ and $s > 0$. There is a positive constant C such that*

$$\int_{\mathbb{B}_n} \frac{(1 - |w|^2)^t dv(w)}{|1 - \langle z, w \rangle|^{n+1+t+s}} \leq C (1 - |z|^2)^{-s}$$

for all $z \in \mathbb{B}_n$.

We also need the concept of an r -lattice in the Bergman metric. Let $r > 0$. A sequence $\{a_k\}$ of points in \mathbb{B}_n is called an r -lattice, if the unit ball is covered by the Bergman metric balls $\{D(a_k, r)\}$, and $\beta(a_i, a_j) \geq r/2$ for all i and j with $i \neq j$. If $\{a_k\}$ is an r -lattice in \mathbb{B}_n , then it also has the following property: for any $R > 0$ there exists a positive integer N (depending on r and R) such that every point in \mathbb{B}_n belongs to at most N sets in $\{D(a_k, R)\}$. There are elementary constructions of r -lattices in \mathbb{B}_n .

Proof of the necessity in Theorem 1. The case $p \geq 1$ (or $p(n+1+\alpha) > n$) follows directly from part (b) of Theorem A since $B^t \mu(z) \leq C \tilde{\mu}(z)$. If $0 < p \leq 1$, consider an r -lattice $\{a_k\}$. Since $|1 - \langle z, w \rangle|$ is comparable to $|1 - \langle z, a_k \rangle|$ for $w \in D(a_k, r)$, we have

$$\begin{aligned} \int_{\mathbb{B}_n} B^t \mu(z)^p d\lambda(z) &\leq \int_{\mathbb{B}_n} \left(\sum_k \int_{D(a_k, r)} \frac{d\mu(w)}{|1 - \langle z, w \rangle|^{2(n+1+\alpha+t)}} \right)^p (1 - |z|^2)^{p(n+1+\alpha+2t)} d\lambda(z) \\ &\leq C \int_{\mathbb{B}_n} \left(\sum_k \frac{\mu(D(a_k, r))}{|1 - \langle z, a_k \rangle|^{2(n+1+\alpha+t)}} \right)^p (1 - |z|^2)^{p(n+1+\alpha+2t)-n-1} dv(z) \\ &\leq C \sum_k \mu(D(a_k, r))^p \int_{\mathbb{B}_n} \frac{(1 - |z|^2)^{p(n+1+\alpha+2t)-n-1} dv(z)}{|1 - \langle z, a_k \rangle|^{2p(n+1+\alpha+t)}}. \end{aligned}$$

Since $p(n+1+\alpha+2t) > n$ we can apply Lemma B and the fact that $v_\alpha(D(a_k, r))$ is comparable to $(1 - |a_k|^2)^{n+1+\alpha}$ in order to obtain

$$\int_{\mathbb{B}_n} B^t \mu(z)^p d\lambda(z) \leq C \sum_k \frac{\mu(D(a_k, r))^p}{(1 - |a_k|^2)^{(n+1+\alpha)p}} \leq C \sum_k \hat{\mu}_r(a_k)^p.$$

Now, it has been already proved (see [8, p. 305]) that

$$\sum_k \hat{\mu}_r(a_k)^p \leq CN \int_{\mathbb{B}_n} \hat{\mu}_{2r}(z)^p d\lambda(z),$$

and this together with part (a) of Theorem A finishes the proof of the necessity. \square

Before going to the proof of the sufficiency we need several properties on reproducing kernels. Recall that $A_\alpha^2(\mathbb{B}_n)$ is a reproducing kernel Hilbert space with the reproducing kernel function given by

$$K_z^\alpha(w) = \frac{1}{(1 - \langle w, z \rangle)^{n+1+\alpha}}, \quad z, w \in \mathbb{B}_n$$

with norm $\|K_z^\alpha\|_\alpha = \sqrt{K_z^\alpha(z)} = (1 - |z|^2)^{-(n+1+\alpha)/2}$ (see [6]). The normalized kernel functions are denoted by $k_z^\alpha = K_z^\alpha / \|K_z^\alpha\|_\alpha$. If $\{e_k\}$ is any orthonormal set of $A_\alpha^2(\mathbb{B}_n)$ then

$$(1) \quad \sum_k |e_k(z)|^2 \leq \|K_z^\alpha\|_\alpha^2,$$

with equality if $\{e_k\}$ is an orthonormal basis of $A_\alpha^2(\mathbb{B}_n)$.

Proof of the sufficiency in Theorem 1: case $p \geq 1$. Assume that $B^t \mu$ is in $L^p(\mathbb{B}_n, d\lambda)$. In order to show that T_μ is in S_p , it is enough to see that there is a positive constant C such that

$$\sum_k \langle T_\mu e_k, e_k \rangle_\alpha^p \leq C$$

for all orthonormal sets $\{e_k\}$ of $A_\alpha^2(\mathbb{B}_n)$ (see [7]). Since

$$\langle T_\mu e_k, e_k \rangle_\alpha = \int_{\mathbb{B}_n} |e_k(z)|^2 d\mu(z),$$

the use of the estimate (a consequence of [6, Lemma 2.24] and the comparability of the quantities $(1 - |z|^2)$, $|1 - \langle z, w \rangle|$ and $1 - |w|^2$ for $w \in D(z, r)$)

$$|e_k(z)|^2 \leq C \int_{\mathbb{B}_n} \frac{|e_k(w)|^2 (1 - |w|^2)^{n+1+\alpha+2t} dv_\alpha(w)}{|1 - \langle z, w \rangle|^{2(n+1+\alpha+t)}}$$

together with Fubini's theorem yields

$$\langle T_\mu e_k, e_k \rangle_\alpha \leq C \int_{\mathbb{B}_n} |e_k(w)|^2 B^t \mu(w) dv_\alpha(w).$$

Thus, by Hölder's inequality and (1)

$$\begin{aligned} \sum_k \langle T_\mu e_k, e_k \rangle_\alpha^p &\leq C \int_{\mathbb{B}_n} \left(\sum_k |e_k(w)|^2 \right) B^t \mu(w)^p dv_\alpha(w) \\ &\leq C \int_{\mathbb{B}_n} B^t \mu(w)^p \|K_w^\alpha\|_\alpha^2 dv_\alpha(w) \leq C \|B^t \mu\|_{L^p(\mathbb{B}_n, d\lambda)}^p. \end{aligned}$$

□

In order to deal with the remaining case $0 < p < 1$ of the sufficiency, we need to introduce some “derivatives” of the kernel functions. For $z, w \in \mathbb{B}_n$, define

$$K_z^{\alpha, t}(w) = \frac{1}{(1 - \langle w, z \rangle)^{n+1+\alpha+t}}$$

and let $k_z^{\alpha, t}$ denote its normalization, that is, $k_z^{\alpha, t} = K_z^{\alpha, t} / \|K_z^{\alpha, t}\|_\alpha$. Notice that

$$K_z^{\alpha, t}(w) = \overline{R^{\alpha, t} K_w^\alpha(z)},$$

where $R^{\alpha, t}$ is the unique continuous linear operator on $H(\mathbb{B}_n)$, the space of all holomorphic functions on \mathbb{B}_n , satisfying

$$R^{\alpha, t} \left(\frac{1}{(1 - \langle z, w \rangle)^{n+1+\alpha}} \right) = \frac{1}{(1 - \langle z, w \rangle)^{n+1+\alpha+t}}$$

for all $w \in \mathbb{B}_n$ (see [7, Section 1.4]). In particular, since $f(z) = \langle f, K_z^\alpha \rangle_\alpha$ whenever $f \in A_\alpha^2(\mathbb{B}_n)$, one has

$$(2) \quad R^{\alpha, t} f(z) = \langle f, K_z^{\alpha, t} \rangle_\alpha, \quad f \in A_\alpha^2(\mathbb{B}_n).$$

Here we want to notice that

$$\tilde{\mu}(z) = \langle T_\mu k_z^\alpha, k_z^\alpha \rangle_\alpha = \int_{\mathbb{B}_n} |k_z^\alpha(w)|^2 d\mu(w)$$

and, since $\|K_z^{\alpha,t}\|_\alpha^2$ is comparable to $(1 - |z|^2)^{-n-1-\alpha-2t}$, we have that $B^t\mu(z)$ corresponds essentially to the quantity

$$\langle T_\mu k_z^{\alpha,t}, k_z^{\alpha,t} \rangle_\alpha = \int_{\mathbb{B}_n} |k_z^{\alpha,t}(w)|^2 d\mu(w).$$

Taking this into account, the sufficiency for $0 < p < 1$ will follow from the next lemma. Similar results involving first order derivatives of reproducing kernels appear in [1], [3] and [4].

Lemma 2. *Let $T : A_\alpha^2(\mathbb{B}_n) \rightarrow A_\alpha^2(\mathbb{B}_n)$ be a positive operator. For $t \geq 0$ set*

$$\widetilde{T}^t(z) = \langle T k_z^{\alpha,t}, k_z^{\alpha,t} \rangle_\alpha, \quad z \in \mathbb{B}_n.$$

- (a) *Let $0 < p \leq 1$. If $\widetilde{T}^t \in L^p(\mathbb{B}_n, d\lambda)$ then T is in S_p .*
- (b) *Let $p \geq 1$. If T is in S_p then $\widetilde{T}^t \in L^p(\mathbb{B}_n, d\lambda)$.*

Proof. Let $p > 0$. The positive operator T is in S_p if and only if T^p is in the trace class S_1 . Fix an orthonormal basis $\{e_k\}$ of $A_\alpha^2(\mathbb{B}_n)$. Since T^p is positive, it belongs to the trace class if and only if

$$\sum_k \langle T^p e_k, e_k \rangle_\alpha < \infty.$$

Let $S = \sqrt{T^p}$. Then

$$\sum_k \langle T^p e_k, e_k \rangle_\alpha = \sum_k \|S e_k\|_\alpha^2,$$

and, by [6, Theorem 2.19], this is comparable to

$$\sum_k \|R^{\alpha,t} S e_k\|_{\alpha+2t}^2.$$

Now, by (2), Fubini's theorem and Parseval's identity, we have

$$\begin{aligned} \sum_k \|R^{\alpha,t} S e_k\|_{\alpha+2t}^2 &= \sum_k \int_{\mathbb{B}_n} |R^{\alpha,t} S e_k(z)|^2 dv_{\alpha+2t}(z) \\ &= \sum_k \int_{\mathbb{B}_n} |\langle S e_k, K_z^{\alpha,t} \rangle_\alpha|^2 dv_{\alpha+2t}(z) \\ &= \int_{\mathbb{B}_n} \left(\sum_k |\langle e_k, S K_z^{\alpha,t} \rangle_\alpha|^2 \right) dv_{\alpha+2t}(z) \\ &= \int_{\mathbb{B}_n} \|S K_z^{\alpha,t}\|_\alpha^2 dv_{\alpha+2t}(z) \\ &= \int_{\mathbb{B}_n} \langle T^p K_z^{\alpha,t}, K_z^{\alpha,t} \rangle_\alpha dv_{\alpha+2t}(z) \\ &= \int_{\mathbb{B}_n} \langle T^p k_z^{\alpha,t}, k_z^{\alpha,t} \rangle_\alpha \|K_z^{\alpha,t}\|_\alpha^2 dv_{\alpha+2t}(z). \end{aligned}$$

Putting all together and taking into account that $\|K_z^{\alpha,t}\|_\alpha^2 (1 - |z|^2)^{\alpha+2t}$ is comparable to $(1 - |z|^2)^{-(n+1)}$, we have that T is in S_p if and only if

$$\int_{\mathbb{B}_n} \langle T^p k_z^{\alpha,t}, k_z^{\alpha,t} \rangle_\alpha d\lambda(z) < \infty.$$

Now, both (a) and (b) are deduced from the inequalities (see [7, Proposition 1.31])

$$\langle T^p k_z^{\alpha,t}, k_z^{\alpha,t} \rangle_\alpha \leq [\langle T k_z^{\alpha,t}, k_z^{\alpha,t} \rangle_\alpha]^p = [\widetilde{T}^t(z)]^p, \quad 0 < p \leq 1$$

and

$$[\widetilde{T}^t(z)]^p = [\langle T k_z^{\alpha,t}, k_z^{\alpha,t} \rangle_\alpha]^p \leq \langle T^p k_z^{\alpha,t}, k_z^{\alpha,t} \rangle_\alpha, \quad p \geq 1.$$

This finishes the proof of the lemma. \square

Proof of the sufficiency in Theorem 1: case $0 < p < 1$. Since $B^t \mu(z)$ is comparable to $\widetilde{T}^t(z)$, the result follows directly from Lemma 2, part (a). \square

Acknowledgments: The author would like to thank the referee for his/her valuable comments that improved the final version of the paper.

REFERENCES

- [1] Z. Harper and M.P. Smith, *Testing Schatten class Hankel operators, Carleson embeddings and weighted composition operators on reproducing kernels*, J. Operator Theory 55 (2006), 349–371.
- [2] D.H. Luecking, *Trace ideal criteria for Toeplitz operators*, J. Funct. Anal. 73 (1987), 345–368.
- [3] J. Pau and J.A. Peláez, *Schatten classes of integration operators on Dirichlet spaces*, J. Anal. Math., to appear.
- [4] M.P. Smith, *Testing Schatten class Hankel operators and Carleson embeddings via reproducing kernels*, J. London Math. Soc. 71 (2005), 172–186.
- [5] K. Zhu, *Positive Toeplitz operators on weighted Bergman spaces of bounded symmetric domains*, J. Operator Theory 20 (1988), 329–357.
- [6] K. Zhu, ‘Spaces of Holomorphic Functions in the Unit Ball’, Springer-Verlag, New York, 2005.
- [7] K. Zhu, ‘Operator Theory in Function Spaces’, Second Edition, Math. Surveys and Monographs, Vol. 138, American Mathematical Society: Providence, Rhode Island, 2007.
- [8] K. Zhu, *Schatten class Toeplitz operators on weighted Bergman spaces of the unit ball*, New York J. Math. 13 (2007), 299–316.

JORDI PAU, DEPARTAMENT DE MATEMÀTICA APLICADA I ANALISI, UNIVERSITAT DE BARCELONA,
 GRAN VIA 585, 08007 BARCELONA, SPAIN
E-mail address: jordi.pau@ub.edu