



Mitogen-Activated Protein Kinases and Mitogen Kinase Phosphatase 1: A Critical Interplay in Macrophage Biology

Jorge Lloberas, Lorena Valverde-Estrella, Juan Tur, Tania Vico and Antonio Celada *

Departament of Cell Biology, Physiology and Immunology, Universitat de Barcelona, Barcelona, Spain

Macrophages are necessary in multiple processes during the immune response or inflammation. This review emphasizes the critical role of the mitogen-activated protein kinases (MAPKs) and mitogen kinase phosphatase-1 (MKP-1) in the functional activities of macrophages. While the phosphorylation of MAPKs is required for macrophage activation or proliferation, MKP-1 dephosphorylates these kinases, thus playing a balancing role in the control of macrophage behavior. MKP-1 is a nuclear-localized dual-specificity phosphatase whose expression is regulated at multiple levels, including at the transcriptional and post-transcriptional level. The regulatory role of MKP-1 in the interplay between MAPK phosphorylation/dephosphorylation makes this molecule a critical regulator of macrophage biology and inflammation.

Keywords: kinases, phosphatases, inflammation, macrophages, immune response, signal transduction

OPEN ACCESS

Edited by:

Ana Cuenda,
Spanish National Research Council,
Spain

Reviewed by:

Eusebio Perdiguero,
University Pompeu Fabra, Spain
Jose M. Lizcano,
Universitat Autònoma de Barcelona,
Spain

*Correspondence:

Antonio Celada
acelada@ub.edu

Specialty section:

This article was submitted to
Cellular Biochemistry,
a section of the journal
Frontiers in Molecular Biosciences

Received: 15 February 2016

Accepted: 13 June 2016

Published: 28 June 2016

Citation:

Lloberas J, Valverde-Estrella L, Tur J,
Vico T and Celada A (2016)
Mitogen-Activated Protein Kinases
and Mitogen Kinase Phosphatase 1: A
Critical Interplay in Macrophage
Biology. *Front. Mol. Biosci.* 3:28.
doi: 10.3389/fmolb.2016.00028

MACROPHAGES A KEY ELEMENT OF THE IMMUNE SYSTEM

Monocytes are the precursors of macrophages and they originate from undifferentiated stem cells in the bone marrow. Differentiation is induced in response to growth factors such as M-CSF, IL-3, and GM-CSF. M-CSF being the only specific one because the M-CSF receptor (CSF1R, M-CSF-R, or CD115 is encoded by the *c-fms* proto-oncogene) is present only in the monocyte/macrophage lineage. Monocytes reach all the tissues in the body via the blood and then they become macrophages (Epelman et al., 2014). In response to various factors, including cytokines, cell-cell contacts, and extracellular matrix interactions, macrophages differentiate to more specialized cells, depending on the tissue. For example, in response to RANKL, these cells differentiate to osteoclasts in the bone (Edwards and Mundy, 2011), to Kupffer cells in the liver, to Langerhans cells in the skin, to bone marrow macrophages in the bone, and to crypt macrophages in the intestine. The activities of differentiated macrophages vary in function of the tissue. The interaction of macrophages with factors such as GM-CSF, IL-4, and TGF- β induces differentiation to dendritic cells. These cells are key elements in antigen presentation in the lymph node a process that induces T cell activation and the acquired immune response. The immune system responds to stress induced by chemical, physical, and infectious agents by producing inflammation. In this process, macrophages play a key role, initially through their capacity to remove bacteria or parasites. This activity is achieved through many components that can indirectly damage the surrounding tissues. After this pro-inflammatory activity (classical activation or M1), macrophages remove all the damaged tissues and start the reconstruction process. In this anti-inflammatory phase (alternative activation or M2), macrophages prompt the synthesis of the extracellular matrix and cell growth (Dey et al., 2014).

Under physiological conditions, monocytes show the Ly6C⁻ and CD43⁺⁺ phenotype in mice and CD14⁻ and CD16⁺⁺ phenotype in humans (Ziegler-Heitbrock et al., 2010). Localized around blood vessels, these cells serve to monitor healthy tissues. When homeostasis is altered, they enter the affected tissues to initiate an early immune response. During the first 24 h of this response, the inflammatory loci are invaded first by neutrophils and then by monocytes carrying the Ly6C⁺⁺ CD43⁻ phenotype in mice and the CD14⁺⁺ CD16⁻ phenotype in humans. These monocytes trigger the pro- and anti-inflammatory activities mentioned earlier. The same macrophages that are polarized to pro-inflammatory action become anti-inflammatory a few days later (Arnold et al., 2007; Takeuchi and Akira, 2011). Interestingly, pro-inflammatory activity by agents such IFN- γ or LPS block macrophage proliferation (Xaus et al., 1999). However, during anti-inflammatory activation, tissue macrophages proliferate locally (Arnold et al., 2007) in a process mediated by IL-4 (Jenkins et al., 2011).

MAPKs AND MACROPHAGE BIOLOGY

MAPKs have been conserved over evolution; however, the subcellular compartmentalization and the kinetics of MAPK activation are cell-type specific, and these kinases orchestrate a range of cellular responses. One of the critical issues in macrophage biology, as in most cell types, is their capacity to respond to stimuli and to proliferate. Although M-CSF, GM-CSF, and IL-3 induce macrophage proliferation, M-CSF is the only specific growth factor for these cells. M-CSF interacts with its receptor-specific receptor CSF1R on the cell surface and induces the activation of signal transduction. The interaction is followed by receptor dimerization, which leads to the autophosphorylation of tyrosine residues in the intracellular domain and the recruitment of signaling molecules (Yu et al., 2012). Macrophage proliferation required the stimulation of the MAPK signaling pathway (Pixley and Stanley, 2004). These serine/threonine kinases are induced by external signals, and they play a critical role in regulating the growth, activation, differentiation, and apoptosis of various types of cells. MAPK activation calls for phosphorylation on the threonine and tyrosine residues situated in the activation loop. Upon phosphorylation, these kinases regulate transcription factors such as Ets-1, Elk/TCF, and AP-1, all of which are involved in immediate, early and late gene expression (Yoon and Seger, 2006), and also in protein expression by regulating the stability, transport or translocation of mRNA species that contain AU-rich elements (Wang and Liu, 2007). In response to growth factors, ERK-1/2, c-Jun NH2-terminal kinase 1 (JNK-1), and p38 are activated in macrophages. However, for the proliferation of these cells, only ERK-1/2 phosphorylation is required (Jaworowski et al., 1999; Valledor et al., 2000a; Sanchez-Tillo et al., 2007). The activation/phosphorylation of Ras is necessary to phosphorylate ERK-1/2 and the downstream targets such as the ribosomal S6 kinases and mitogen- and stress-activated kinases 1 and 2 (Murphy and Blenis, 2006). Activated ERK-1/2 translocate to the nucleus and, by means of phosphorylation, activate a number of transcription factors

that regulate genes involved in proliferation. Remarkably, M-CSF triggers other pathways independent of MAPKs to induce macrophage survival through the phosphorylation of phosphoinositide 3-kinase which prompts the translocation and activation of Akt (Comalada et al., 2004).

Lipopolysaccharide (LPS) is a powerful inflammatory signal present in the membrane of Gram-negative bacteria. It regulates the expression of many genes, including TNF- α , which causes septic shock when present in large amounts. ERK-1/2 becomes phosphorylated through the interaction of LPS with LPS binding protein (a serum protein), CD14, and TLR4 (membrane proteins in macrophages; Valledor et al., 2000b). The inhibition of this phosphorylation blocks the production of TNF- α and IL-1 β . This observation serves to demonstrate the role of these two molecules in macrophage activation (Valledor et al., 2000a).

IFN- γ , a major endogenous macrophage activator, not only requires Stat1 for signaling but also MAPKs. At early times (30 min), p38 is activated strongly, but ERK-1/2 and JNK-1 become phosphorylated between 2 and 5 h. Although p38 and ERK-1/2 regulate the expression of genes of the innate response, such as chemokines, TNF- α , and inducible NO synthase, the genes involved in antigen presentation are regulated by JNK-1 (Valledor et al., 2008b). These results emphasize the diversity of MAPK kinetics and activity in macrophages.

The kinetics of MAPK phosphorylation with respect to the proliferation or functional activation of macrophages may be associated with various factors, such as cell-surface receptor concentration and the different pathways used after ligand-receptor engagement (Murphy and Blenis, 2006). The induction of ERK-1/2 phosphorylation by M-CSF or LPS involves the same molecules, namely Ras, Raf, and MEK 1/2 (Casals-Casas et al., 2009). However, the early steps differ. While M-CSF receptor encloses a tyrosine kinase domain in the intra-cytoplasmic region, thus inducing rapid phosphorylation, LPS interacts with LPS-binding protein CD14, and finally TLR4 and requires more time to initiate the phosphorylation (Kolch, 2005).

THE MKP-1 PHOSPHATASE AND ITS REGULATION

MKP-1 phosphatase, also called dual specificity phosphatase 1 (DUSP1), is expressed ubiquitously. However, it is regulated in a cell-context manner and is able to dephosphorylate tyrosine or serine/threonine residues. MKPs dephosphorylate and consequently inactivate ERK-1/2 (Pouyssegur and Lenormand, 2003), p38, MAPKs (Kaiser et al., 2004), and JNKs (Sanchez-Perez et al., 1998). At least 11 MKP family members of these dual phosphatases have been found to differ in cellular specificity, subcellular localization, and substrate specificity (Farooq and Zhou, 2004).

MKP-1 is composed of two domains, one is bound the MAPK in the NH2-terminus and the other is the dual-specificity phosphatase domain located in the COOH terminus (Camps et al., 2000). This phosphatase localizes exclusively to the nucleus through a NH2-terminus LXXLL motif (Wu

et al., 2005). The catalytic activity is regulated through interaction with MAPKs mediated by a kinase interaction motif (Doddareddy et al., 2012).

In most cases, the induction of phosphatases is a negative response mechanism of the kinases that triggers the expression or activation of the phosphatases. In macrophages, MPK-1 induction yields conflicting results. In primary cultures of bone marrow-derived macrophages, LPS-induced MKP-1 expression is partially blocked by co-administration of inhibitors of ERK-1/2 and p38 (Ananieva et al., 2008) or when macrophage from a p38 KO mouse are used (Kim et al., 2008). In other studies using the same cell type, the inhibition of the MEK pathway required for M-CSF- or LPS-induced ERK-1/2 phosphorylation was not found to affect MKP-1 expression, thereby showing that this expression is independent of kinase activation (Valledor et al., 1999, 2000b).

The early steps in signal transduction to induce MKP-1 involves the phosphorylation of PKC ϵ (Valledor et al., 1999, 2000b). Interestingly, the expression of MKP-1 induced by either M-CSF or LPS also requires Raf-1 activation, as does the phosphorylation of ERK-1/2 (Sanchez-Tillo et al., 2006). Another mediator of MKP-1 stimulation is JNK-1, and inactivation of this kinase results in prolonged ERK phosphorylation (Sanchez-Tillo et al., 2007). Finally, inhibition of mTOR induces MKP-1 expression, thereby suggesting that the repression of immunity is mediated through MKP-1 (Rastogi et al., 2013).

MKP-1 is not expressed under basal conditions; however, in response to various stimuli MKP-1 is an immediate early gene that requires transcriptional activation, although its regulation is also mediated at several other levels. In the MKP-1 promoter, a region containing a CREB/AP-1 box is essential for the induction of transcription through a variety of different stimuli (Arthur et al., 2004; Ananieva et al., 2008; Cho et al., 2008; Kim et al., 2008; Casals-Casas et al., 2009). Upon activation by M-CSF or LPS, this box is occupied by phosphorylated CREB and c-Jun. It has been reported that the phosphorylation of CREB is mediated by p38 (Kamata et al., 2005) and by ERK-1/2. However, this phosphorylation is not direct but mediated through MSK1/2 activated by either ERK-1/2 or p38 (Ananieva et al., 2008; Kim et al., 2008). However, it has been reported that the inhibition of ERK does not block MKP-1 induction by M-CSF and LPS (Valledor et al., 1999, 2000b). Other studies have described that CREB phosphorylation is mediated by M-CSF-induced PI-3K (Kanagasundaram et al., 1999) or by LPS-activated PKA (Zhong et al., 1997). Vitamin D induces MKP-1, thus inhibiting pro-inflammatory cytokine production, and the binding of the vitamin D receptor at the vitamin D response element in the MKP-1 promoter is increased (Zhang et al., 2012). In the context of the proliferation vs. activation issue, IFN- γ , the major endogenous activator of macrophages, inhibits M-CSF-induced MKP-1 expression (Valledor et al., 2008a). The inhibition is mediated through STAT1, but no such repressive activity has been reported for this transcription factor. Transcription, in turn, calls for epigenetic modifications or chromatin remodeling on the MKP-1 promoter region. Through phosphorylation and acetylation of histone H3, chromatin is modified, thereby increasing the binding of RNA

polymerase II and inducing MKP-1 transcription (Li et al., 2001; **Figure 1**).

MKP-1 mRNA in macrophages has a very short half-life (Valledor et al., 1999). In other cellular models, the short mRNA half-life has been related to AU-rich 3' untranslated regions and the RNA binding proteins HuR and NF90 (Kuwano et al., 2008). It seems that NF90 represses the translation of mRNAs bearing the AU-rich signature sequence without affecting their half-lives (Kuwano et al., 2010). Tristetraprolin binds to these AU-rich sequences of MKP-1 and thus forms part of a negative feedback loop to limit the mRNA half-life (Emmons et al., 2008). In fact, tristetraprolin expression is suppressed by MKP-1 through inhibition of p38 (Huotari et al., 2012).

The activity of MKP-1 can be modified by stabilization of his on protein, which is determined by its ERK-mediated phosphorylation through a degradation pathway independent of polyubiquitination (Crowell et al., 2014). If Ser359 and Ser364 become phosphorylated, the protein is stabilized (Brondello et al., 1999); however, if phosphorylation occurs in Ser298 and Ser323, then proteosomal degradation is facilitated (Lin et al., 2003). We should mention that glucocorticoids not only induce the transcription of MKP-1 but also attenuate their proteasomal degradation (Kassel et al., 2001; **Figure 1**).


Metabolic stress in monocytes and macrophages induces Nox4, an inducible NADPH oxidase that promotes MKP-1-S-glutathionylation, resulting in MKP1 inactivation and subsequent proteasomal degradation (Kim et al., 2012). This observation shows that a redox-regulated mechanism links oxidative stress directly to metabolic disorders and macrophage hyperactivity.

The repressive cytokines IL-10 and TGF- β block most immune responses and increase LPS-dependent MKP-1 induction, thereby enhancing the degree and duration of the phosphatase and consequently dephosphorylating MAPKs with the consequent inhibition of inflammatory cytokine production (Jono et al., 2003; Hammer et al., 2005). Micro-RNA-101 (miR-101) and miR-210 transfections reduce LPS-dependent induction of MKP-1 and also prolong the phosphorylation of p38 and JNK (Zhu et al., 2010; Jin et al., 2014).

Glucocorticoids are one of the major families of inhibitors of the immune response used in clinical practice. Remarkably, these drugs increase the expression and decrease the degradation of MKP-1. Consequently, MKP-1 dephosphorylates MAPKs, thus decreasing the activity of these kinases, as well as reducing macrophage activation (Kassel et al., 2001; Shipp et al., 2010). However, the observation that MKP-1 KO mice with enhanced cytokine expression remain sensitive to glucocorticoids suggests that the drug targets molecules other than MKP-1 (Maier et al., 2007; Wang et al., 2008).

In response to LPS, MKP-1 is acetylated by p300/CBP on lysine 57 within its substrate-binding domain, thus increasing its interaction and dephosphorylation of p38 (Cao et al., 2008). This observation reveals another level of regulation of the functional activity mediated by histone deacetylase isoforms (HDAC1, HDAC2, and HDAC3; Jeong et al., 2014).

Therefore, the mechanisms regulating MKP-1 expression are critical in determining the duration of MAPK phosphorylation




and the length of the immune response (Manetsch et al., 2012; Tomida et al., 2015).

THE SWITCH BETWEEN PROLIFERATION AND ACTIVATION IS MEDIATED BY MKP-1

The activation and proliferation of macrophages are reciprocally exclusive processes (Xaus et al., 1999); however, both require ERK-1/2 phosphorylation, although with distinct kinetics. For macrophage proliferation, ERK-1/2 phosphorylation occurs with a prompt peak around 5 min, while for activation the phosphorylation takes place later, around 15 min (Valledor et al., 2000a; Suzu et al., 2007; **Figure 2**). The time course of ERK-1/2 dephosphorylation differs for growth factors and activating molecules, occurring 30 min after phosphorylation for M-CSF-induced proliferation and 90 min for LPS-induced functional activation. MAPK

phosphorylation is regulated by the dominant action of protein phosphatases, as evidenced by phosphorylation being reversible even in the continued presence of activating stimuli.

Both stimuli triggering proliferation and functional activation induce nuclear MKP-1 (Valledor et al., 1999, 2000b). The transcription of MKP-1 follows the same kinetics by M-CSF or LPS, producing early or late mRNA, depending on the stimuli. Regardless of whether macrophages are treated with M-CSF or LPS, ERK-1/2 dephosphorylation is associated only with the induction of MKP-1 (Valledor et al., 2008a; **Figure 2**). This induction requires the binding and phosphorylation of CREB and c-Jun, and the kinetics of these two molecules correlates to the different times courses of MKP-1 expression. The distinct kinetics of dephosphorylation suggests that the critical factor that allows LPS or IFN- γ to inhibit macrophage proliferation is related to the duration of ERK phosphorylation. This hypothesis was tested by inhibiting MKP-1 expression. However, Mkp-1 knockout (KO)



mice cannot be used for this purpose because compensatory mechanisms replace the role of MKP-1 in proliferation. In fact, in contrast to wild-type cells in which MKP-4 is not induced by M-CSF, in MKP-1 deficient macrophages, there is an early MKP-4 response to M-CSF. By using siRNA, it was observed that the inhibition of MKP-1 expression extends ERK-1/2 phosphorylation and blocks M-CSF-dependent proliferation (Valledor et al., 2008a).

The inhibition of proliferation induced by IFN- γ stimulation correlates with extended ERK-1/2 phosphorylation induced by M-CSF and is due to the inhibition of MKP-1 expression (Valledor et al., 2008a). Interestingly, this inhibition is mediated by STAT1. This observation would suggest that this transcription factor represses some genes.

The M-CSF-dependent induction of MKP-1 expression can be altered by other agents. MKP-1 expression can be inhibited by signaling through extracellular matrix proteins, such as decorin and fibrinogen (Xaus et al., 2001). As result of such inhibition, ERK phosphorylation is extended, and macrophage proliferation is inhibited (Figure 2).

OTHER ACTIVITIES REGULATED BY MKP-1

MKP-1 KO mice show extended MAPK phosphorylation that correlates with an increased expression of cytokines such as TNF- α , IL-6, and IL-12, all of which increase inflammation (mice exhibit kidney failure, severe hypotension, inflammatory infiltrates in the lung and other tissues, and impaired circulation compared with wild-type mice) and cause higher susceptibility to endotoxic shock (Chi et al., 2006; Hammer et al., 2006, 2010; Salojin et al., 2006; Zhao et al., 2006; Frazier et al., 2009; Rodriguez et al., 2010). In a microarray of 14,000 genes from *mkp-1*^{-/-} mouse spleen cells, 608 genes were found to be upregulated (Hammer et al., 2006). The absence of MKP-1 produces the longest p38 phosphorylation, which in turn increases C/EBP β phosphorylation, a critical process in several LPS-induced genes (Serrat et al., 2014). Another mechanism to by which to increase the gene expression of natural immunity in the absence of MKP-1 may be related to histones such as H3, which is a substrate of MKP-1 (Kinney et al., 2009). The loss of histone dephosphorylation may affect the transcription of several genes.

Interestingly, the half-lives of mRNAs of several cytokines, including IL-6, IL-10, and TNF- α , seem to be controlled by MKP-1 through translocation of RNA binding proteins from the nucleus to the cytosol (Yu et al., 2011). Recent data shows that MKP-1 modulates the activity of the cytokine mRNA by destabilizing the phosphorylation status of tristetraprolin, an RNA-destabilizing protein (Smallie et al., 2015).

In conclusion, in this review we have addressed the critical issue of the duration of ERK-1/2 kinase phosphorylation and its role in mediating macrophage proliferation or functional activation. The duration of the phosphorylation state is determined by the induction of MKP-1, thus making MKP-1 one of the critical regulators of macrophage biology.

AUTHOR CONTRIBUTIONS

JL: review literature, discuss with the other collaborators, made part of the original data and draw the graphics. LV, JT, and TV: review literature, discuss with the other collaborators and made part of the original data. AC: review literature, discuss with the other collaborators, made the original data and wrote the manuscript.

ACKNOWLEDGMENTS

This study was supported by a grant from the *Plan Nacional del Ministerio de Economía y Competividad* SAF2014-52887-R to AC and JL.

REFERENCES

- Ananieva, O., Darragh, J., Johansen, C., Carr, J. M., McIlrath, J., Park, J. M., et al. (2008). The kinases MSK1 and MSK2 act as negative regulators of Toll-like receptor signaling. *Nat. Immunol.* 9, 1028–1036. doi: 10.1038/ni.1644
- Arnold, L., Henry, A., Poron, F., Baba-Amer, Y., Van Rooijen, N., Plonquet, A., et al. (2007). Inflammatory monocytes recruited after skeletal muscle injury switch into antiinflammatory macrophages to support myogenesis. *J. Exp. Med.* 204, 1057–1069. doi: 10.1084/jem.20070075
- Arthur, J. S., Fong, A. L., Dwyer, J. M., Davare, M., Reese, E., Obrietan, K., et al. (2004). Mitogen- and stress-activated protein kinase 1 mediates cAMP response element-binding protein phosphorylation and activation by neurotrophins. *J. Neurosci.* 24, 4324–4332. doi: 10.1523/JNEUROSCI.5227-03.2004
- Brondello, J. M., Pouyssegur, J., and Mckenzie, F. R. (1999). Reduced MAP kinase phosphatase-1 degradation after p42/p44MAPK-dependent phosphorylation. *Science* 286, 2514–2517. doi: 10.1126/science.286.5449.2514
- Camps, M., Nichols, A., and Arkininstall, S. (2000). Dual specificity phosphatases: a gene family for control of MAP kinase function. *FASEB J.* 14, 6–16.
- Cao, W., Bao, C., Padalko, E., and Lowenstein, C. J. (2008). Acetylation of mitogen-activated protein kinase phosphatase-1 inhibits Toll-like receptor signaling. *J. Exp. Med.* 205, 1491–1503. doi: 10.1084/jem.20071728
- Casals-Casas, C., Alvarez, E., Serra, M., de la Torre, C., Farrera, C., Sanchez-Tillo, E., et al. (2009). CREB and AP-1 activation regulates MKP-1 induction by LPS or M-CSF and their kinetics correlate with macrophage activation versus proliferation. *Eur. J. Immunol.* 39, 1902–1913. doi: 10.1002/eji.200839037
- Chi, H., Barry, S. P., Roth, R. J., Wu, J. J., Jones, E. A., Bennett, A. M., et al. (2006). Dynamic regulation of pro- and anti-inflammatory cytokines by MAPK phosphatase 1 (MKP-1) in innate immune responses. *Proc. Natl. Acad. Sci. U.S.A.* 103, 2274–2279. doi: 10.1073/pnas.0510965103
- Cho, I. J., Woo, N. R., and Kim, S. G. (2008). The identification of C/EBPbeta as a transcription factor necessary for the induction of MAPK phosphatase-1 by toll-like receptor-4 ligand. *Arch. Biochem. Biophys.* 479, 88–96. doi: 10.1016/j.abb.2008.08.007
- Comalada, M., Xaus, J., Sanchez, E., Villedor, A. F., and Celada, A. (2004). Macrophage colony-stimulating factor-, granulocyte-macrophage colony-stimulating factor-, or IL-3-dependent survival of macrophages, but not proliferation, requires the expression of p21(Waf1) through the phosphatidylinositol 3-kinase/Akt pathway. *Eur. J. Immunol.* 34, 2257–2267. doi: 10.1002/eji.200425110
- Crowell, S., Wancket, L. M., Shakibi, Y., Xu, P., Xue, J., Samavati, L., et al. (2014). Post-translational regulation of mitogen-activated protein kinase phosphatase (MKP)-1 and MKP-2 in macrophages following lipopolysaccharide stimulation: the role of the C termini of the phosphatases in determining their stability. *J. Biol. Chem.* 289, 28753–28764. doi: 10.1074/jbc.M114.591925
- Dey, A., Allen, J., and Hankey-Giblin, P. A. (2014). Ontogeny and polarization of macrophages in inflammation: blood monocytes versus tissue macrophages. *Front. Immunol.* 5:683. doi: 10.3389/fimmu.2014.00683
- Doddareddy, M. R., Rawling, T., and Ammit, A. J. (2012). Targeting mitogen-activated protein kinase phosphatase-1 (MKP-1): structure-based design of MKP-1 inhibitors and upregulators. *Curr. Med. Chem.* 19, 163–173. doi: 10.2174/092986712803414196
- Edwards, J. R., and Mundy, G. R. (2011). Advances in osteoclast biology: old findings and new insights from mouse models. *Nat. Rev. Rheumatol.* 7, 235–243. doi: 10.1038/nrrheum.2011.23
- Emmons, J., Townley-Tilson, W. H., Deleault, K. M., Skinner, S. J., Gross, R. H., Whitfield, M. L., et al. (2008). Identification of TTP mRNA targets in human dendritic cells reveals TTP as a critical regulator of dendritic cell maturation. *RNA* 14, 888–902. doi: 10.1261/rna.748408
- Epelman, S., Lavine, K. J., and Randolph, G. J. (2014). Origin and functions of tissue macrophages. *Immunity* 41, 21–35. doi: 10.1016/j.immuni.2014.06.013
- Farooq, A., and Zhou, M. M. (2004). Structure and regulation of MAPK phosphatases. *Cell. Signal.* 16, 769–779. doi: 10.1016/j.cellsig.2003.12.008
- Frazier, W. J., Wang, X., Wancket, L. M., Li, X. A., Meng, X., Nelin, L. D., et al. (2009). Increased inflammation, impaired bacterial clearance, and metabolic disruption after gram-negative sepsis in Mkp-1-deficient mice. *J. Immunol.* 183, 7411–7419. doi: 10.4049/jimmunol.0804343
- Hammer, M., Echtenachter, B., Weighardt, H., Jozefowski, K., Rose-John, S., Mannel, D. N., et al. (2010). Increased inflammation and lethality of Dusp1^{-/-} mice in polymicrobial peritonitis models. *Immunology* 131, 395–404. doi: 10.1111/j.1365-2567.2010.03313.x
- Hammer, M., Mages, J., Dietrich, H., Schmitz, F., Striebel, F., Murray, P. J., et al. (2005). Control of dual-specificity phosphatase-1 expression in activated macrophages by IL-10. *Eur. J. Immunol.* 35, 2991–3001. doi: 10.1002/eji.200526192
- Hammer, M., Mages, J., Dietrich, H., Servatius, A., Howells, N., Cato, A. C., et al. (2006). Dual specificity phosphatase 1 (DUSP1) regulates a subset of LPS-induced genes and protects mice from lethal endotoxin shock. *J. Exp. Med.* 203, 15–20. doi: 10.1084/jem.20051753
- Huotari, N., Hommo, T., Taimi, V., Nieminen, R., Moilanen, E., and Korhonen, R. (2012). Regulation of tristetraprolin expression by mitogen-activated protein kinase phosphatase-1. *APMIS* 120, 988–999. doi: 10.1111/j.1600-0463.2012.02927.x
- Jaworowski, A., Wilson, N. J., Christy, E., Byrne, R., and Hamilton, J. A. (1999). Roles of the mitogen-activated protein kinase family in macrophage responses to colony stimulating factor-1 addition and withdrawal. *J. Biol. Chem.* 274, 15127–15133. doi: 10.1074/jbc.274.21.15127
- Jenkins, S. J., Ruckerl, D., Cook, P. C., Jones, L. H., Finkelman, F. D., Van Rooijen, N., et al. (2011). Local macrophage proliferation, rather than recruitment from the blood, is a signature of TH2 inflammation. *Science* 332, 1284–1288. doi: 10.1126/science.1204351
- Jeong, Y., Du, R., Zhu, X., Yin, S., Wang, J., Cui, H., et al. (2014). Histone deacetylase isoforms regulate innate immune responses by deacetylating mitogen-activated protein kinase phosphatase-1. *J. Leukoc. Biol.* 95, 651–659. doi: 10.1189/jlb.1013565
- Jin, C., Peng, X., Liu, F., Cheng, L., Lu, X., Yao, H., et al. (2014). MicroRNA-181 expression regulates specific post-transcriptional level of SAMHD1 expression *in vitro*. *Biochem. Biophys. Res. Commun.* 452, 760–767. doi: 10.1016/j.bbrc.2014.08.151
- Jono, H., Xu, H., Kai, H., Lim, D. J., Kim, Y. S., Feng, X. H., et al. (2003). Transforming growth factor-beta-Smad signaling pathway negatively regulates nontypeable Haemophilus influenzae-induced MUC5AC mucin transcription via mitogen-activated protein kinase (MAPK) phosphatase-1-dependent inhibition of p38 MAPK. *J. Biol. Chem.* 278, 27811–27819. doi: 10.1074/jbc.M301773200
- Kaiser, R. A., Bueno, O. F., Lips, D. J., Doevendans, P. A., Jones, F., Kimball, T. F., et al. (2004). Targeted inhibition of p38 mitogen-activated protein kinase antagonizes cardiac injury and cell death following ischemia-reperfusion *in vivo*. *J. Biol. Chem.* 279, 15524–15530. doi: 10.1074/jbc.M313717200
- Kamata, H., Honda, S., Maeda, S., Chang, L., Hirata, H., and Karin, M. (2005). Reactive oxygen species promote TNFalpha-induced death and sustained JNK activation by inhibiting MAP kinase phosphatases. *Cell* 120, 649–661. doi: 10.1016/j.cell.2004.12.041
- Kanagasundaram, V., Jaworowski, A., Byrne, R., and Hamilton, J. A. (1999). Separation and characterization of the activated pool of colony-stimulating factor 1 receptor forming distinct multimeric complexes with signalling molecules in macrophages. *Mol. Cell. Biol.* 19, 4079–4092. doi: 10.1128/MCB.19.6.4079
- Kassel, O., Sancono, A., Kratzschmar, J., Kreft, B., Stassen, M., and Cato, A. C. (2001). Glucocorticoids inhibit MAP kinase via increased expression and decreased degradation of MKP-1. *EMBO J.* 20, 7108–7116. doi: 10.1093/emboj/20.24.7108
- Kim, C., Sano, Y., Todorova, K., Carlson, B. A., Arpa, L., Celada, A., et al. (2008). The kinase p38 alpha serves cell type-specific inflammatory functions in skin injury and coordinates pro- and anti-inflammatory gene expression. *Nat. Immunol.* 9, 1019–1027. doi: 10.1038/ni.1640
- Kim, H. S., Ullevig, S. L., Zamora, D., Lee, C. F., and Asmis, R. (2012). Redox regulation of MAPK phosphatase 1 controls monocyte migration and macrophage recruitment. *Proc. Natl. Acad. Sci. U.S.A.* 109, E2803–E2812. doi: 10.1073/pnas.1212596109
- Kinney, C. M., Chandrasekharan, U. M., Yang, L., Shen, J., Kinter, M., Mcdermott, M. S., et al. (2009). Histone H3 as a novel substrate for MAP kinase phosphatase-1. *Am. J. Physiol. Cell Physiol.* 296, C242–C249. doi: 10.1152/ajpcell.00492.2008

- Kolch, W. (2005). Coordinating ERK/MAPK signalling through scaffolds and inhibitors. *Nat. Rev. Mol. Cell Biol.* 6, 827–837. doi: 10.1038/nrm1743
- Kuwano, Y., Kim, H. H., Abdelmohsen, K., Pullmann, R. Jr., Martindale, J. L., Yang, X., et al. (2008). MKP-1 mRNA stabilization and translational control by RNA-binding proteins HuR and NF90. *Mol. Cell. Biol.* 28, 4562–4575. doi: 10.1128/MCB.00165-08
- Kuwano, Y., Pullmann, R. Jr., Marasa, B. S., Abdelmohsen, K., Lee, E. K., Yang, X., et al. (2010). NF90 selectively represses the translation of target mRNAs bearing an AU-rich signature motif. *Nucleic Acids Res.* 38, 225–238. doi: 10.1093/nar/gkp861
- Li, J., Gorospe, M., Hutter, D., Barnes, J., Keyse, S. M., and Liu, Y. (2001). Transcriptional induction of MKP-1 in response to stress is associated with histone H3 phosphorylation-acetylation. *Mol. Cell. Biol.* 21, 8213–8224. doi: 10.1128/MCB.21.23.8213-8224.2001
- Lin, Y. W., Chuang, S. M., and Yang, J. L. (2003). ERK1/2 achieves sustained activation by stimulating MAPK phosphatase-1 degradation via the ubiquitin-proteasome pathway. *J. Biol. Chem.* 278, 21534–21541. doi: 10.1074/jbc.M301854200
- Maier, J. V., Brema, S., Tuckermann, J., Herzer, U., Klein, M., Stassen, M., et al. (2007). Dual specificity phosphatase 1 knockout mice show enhanced susceptibility to anaphylaxis but are sensitive to glucocorticoids. *Mol. Endocrinol.* 21, 2663–2671. doi: 10.1210/me.2007-0067
- Manetsch, M., Che, W., Seidel, P., Chen, Y., and Ammit, A. J. (2012). MKP-1: a negative feedback effector that represses MAPK-mediated pro-inflammatory signaling pathways and cytokine secretion in human airway smooth muscle cells. *Cell. Signal.* 24, 907–913. doi: 10.1016/j.cellsig.2011.12.013
- Murphy, L. O., and Blenis, J. (2006). MAPK signal specificity: the right place at the right time. *Trends Biochem. Sci.* 31, 268–275. doi: 10.1016/j.tibs.2006.03.009
- Pixley, F. J., and Stanley, E. R. (2004). CSF-1 regulation of the wandering macrophage: complexity in action. *Trends Cell Biol.* 14, 628–638. doi: 10.1016/j.tcb.2004.09.016
- Pouyssegur, J., and Lenormand, P. (2003). Fidelity and spatio-temporal control in MAP kinase (ERKs) signalling. *Eur. J. Biochem.* 270, 3291–3299. doi: 10.1046/j.1432-1033.2003.03707.x
- Rastogi, R., Jiang, Z., Ahmad, N., Rosati, R., Liu, Y., Beuret, L., et al. (2013). Rapamycin induces mitogen-activated protein (MAP) kinase phosphatase-1 (MKP-1) expression through activation of protein kinase B and mitogen-activated protein kinase kinase pathways. *J. Biol. Chem.* 288, 33966–33977. doi: 10.1074/jbc.M113.492702
- Rodriguez, N., Dietrich, H., Mossbrugger, I., Weintz, G., Scheller, J., Hammer, M., et al. (2010). Increased inflammation and impaired resistance to Chlamydia pneumoniae infection in Dusp1(−/−) mice: critical role of IL-6. *J. Leukoc. Biol.* 88, 579–587. doi: 10.1189/jlb.0210083
- Salojin, K. V., Owusu, I. B., Millerchip, K. A., Potter, M., Platt, K. A., and Oravec, T. (2006). Essential role of MAPK phosphatase-1 in the negative control of innate immune responses. *J. Immunol.* 176, 1899–1907. doi: 10.4049/jimmunol.176.3.1899
- Sanchez-Perez, I., Murguía, J. R., and Perona, R. (1998). Cisplatin induces a persistent activation of JNK that is related to cell death. *Oncogene* 16, 533–540. doi: 10.1038/sj.onc.1201578
- Sanchez-Tillo, E., Comalada, M., Farrera, C., Valledor, A. F., Lloberas, J., and Celada, A. (2006). Macrophage-colony-stimulating factor-induced proliferation and lipopolysaccharide-dependent activation of macrophages requires Raf-1 phosphorylation to induce mitogen kinase phosphatase-1 expression. *J. Immunol.* 176, 6594–6602. doi: 10.4049/jimmunol.176.11.6594
- Sanchez-Tillo, E., Comalada, M., Xaus, J., Farrera, C., Valledor, A. F., Caelles, C., et al. (2007). JNK1 Is required for the induction of Mkp1 expression in macrophages during proliferation and lipopolysaccharide-dependent activation. *J. Biol. Chem.* 282, 12566–12573. doi: 10.1074/jbc.M609662200
- Serrat, N., Sebastian, C., Pereira-Lopes, S., Valverde-Estrella, L., Lloberas, J., and Celada, A. (2014). The response of secondary genes to lipopolysaccharides in macrophages depends on histone deacetylase and phosphorylation of C/EBPβ. *J. Immunol.* 192, 418–426. doi: 10.4049/jimmunol.1203500
- Shipp, L. E., Lee, J. V., Yu, C. Y., Pufall, M., Zhang, P., Scott, D. K., et al. (2010). Transcriptional regulation of human dual specificity protein phosphatase 1 (DUSP1) gene by glucocorticoids. *PLoS ONE* 5:e13754. doi: 10.1371/journal.pone.0013754
- Smallie, T., Ross, E. A., Ammit, A. J., Cunliffe, H. E., Tang, T., Rosner, D. R., et al. (2015). Dual-specificity phosphatase 1 and tristetraprolin cooperate to regulate macrophage responses to lipopolysaccharide. *J. Immunol.* 195, 277–288. doi: 10.4049/jimmunol.1402830
- Suzu, S., Hiyoshi, M., Yoshidomi, Y., Harada, H., Takeya, M., Kimura, F., et al. (2007). M-CSF-mediated macrophage differentiation but not proliferation is correlated with increased and prolonged ERK activation. *J. Cell. Physiol.* 212, 519–525. doi: 10.1002/jcp.21045
- Takeuch, O., and Akira, S. (2011). Epigenetic control of macrophage polarization. *Eur. J. Immunol.* 41, 2490–2493. doi: 10.1002/eji.201141792
- Tomida, T., Takekawa, M., and Saito, H. (2015). Oscillation of p38 activity controls efficient pro-inflammatory gene expression. *Nat. Commun.* 6:8350. doi: 10.1038/ncomms9350
- Valledor, A. F., Arpa, L., Sanchez-Tillo, E., Comalada, M., Casals, C., Xaus, J., et al. (2008a). IFN-γ-mediated inhibition of MAPK phosphatase expression results in prolonged MAPK activity in response to M-CSF and inhibition of proliferation. *Blood* 112, 3274–3282. doi: 10.1182/blood-2007-11-123604
- Valledor, A. F., Comalada, M., Xaus, J., and Celada, A. (2000a). The differential time-course of extracellular-regulated kinase activity correlates with the macrophage response toward proliferation or activation. *J. Biol. Chem.* 275, 7403–7409. doi: 10.1074/jbc.275.10.7403
- Valledor, A. F., Sanchez-Tillo, E., Arpa, L., Park, J. M., Caelles, C., Lloberas, J., et al. (2008b). Selective roles of MAPKs during the macrophage response to IFN-γ. *J. Immunol.* 180, 4523–4529. doi: 10.4049/jimmunol.180.7.4523
- Valledor, A. F., Xaus, J., Comalada, M., Soler, C., and Celada, A. (2000b). Protein kinase C epsilon is required for the induction of mitogen-activated protein kinase phosphatase-1 in lipopolysaccharide-stimulated macrophages. *J. Immunol.* 164, 29–37. doi: 10.4049/jimmunol.164.1.29
- Valledor, A. F., Xaus, J., Marques, L., and Celada, A. (1999). Macrophage colony-stimulating factor induces the expression of mitogen-activated protein kinase phosphatase-1 through a protein kinase C-dependent pathway. *J. Immunol.* 163, 2452–2462.
- Wang, X., and Liu, Y. (2007). Regulation of innate immune response by MAP kinase phosphatase-1. *Cell. Signal.* 19, 1372–1382. doi: 10.1016/j.cellsig.2007.03.013
- Wang, X., Nelin, L. D., Kuhlman, J. R., Meng, X., Welty, S. E., and Liu, Y. (2008). The role of MAP kinase phosphatase-1 in the protective mechanism of dexamethasone against endotoxemia. *Life Sci.* 83, 671–680. doi: 10.1016/j.lfs.2008.09.003
- Wu, J. J., Zhang, L., and Bennett, A. M. (2005). The noncatalytic amino terminus of mitogen-activated protein kinase phosphatase 1 directs nuclear targeting and serum response element transcriptional regulation. *Mol. Cell. Biol.* 25, 4792–4803. doi: 10.1128/MCB.25.11.4792-4803.2005
- Xaus, J., Cardo, M., Valledor, A. F., Soler, C., Lloberas, J., and Celada, A. (1999). Interferon gamma induces the expression of p21 waf-1 and arrests macrophage cell cycle, preventing induction of apoptosis. *Immunity* 11, 103–113. doi: 10.1016/S1074-7613(00)80085-0
- Xaus, J., Comalada, M., Cardo, M., Valledor, A. F., and Celada, A. (2001). Decorin inhibits macrophage colony-stimulating factor proliferation of macrophages and enhances cell survival through induction of p27(Kip1) and p21(Waf1). *Blood* 98, 2124–2133. doi: 10.1182/blood.V98.7.2124
- Yoon, S., and Seger, R. (2006). The extracellular signal-regulated kinase: multiple substrates regulate diverse cellular functions. *Growth Factors* 24, 21–44. doi: 10.1080/02699050500284218
- Yu, H., Sun, Y., Haycraft, C., Palanisamy, V., and Kirkwood, K. L. (2011). MKP-1 regulates cytokine mRNA stability through selectively modulation subcellular translocation of AUF1. *Cytokine* 56, 245–255. doi: 10.1016/j.cyto.2011.06.006
- Yu, W., Chen, J., Xiong, Y., Pixley, F. J., Yeung, Y. G., and Stanley, E. R. (2012). Macrophage proliferation is regulated through CSF-1 receptor tyrosines 544, 559, and 807. *J. Biol. Chem.* 287, 13694–13704. doi: 10.1074/jbc.M112.355610
- Zhang, Y., Leung, D. Y., Richers, B. N., Liu, Y., Remigio, L. K., Riches, D. W., et al. (2012). Vitamin D inhibits monocyte/macrophage proinflammatory cytokine production by targeting MAPK phosphatase-1. *J. Immunol.* 188, 2127–2135. doi: 10.4049/jimmunol.1102412
- Zhao, Q., Wang, X., Nelin, L. D., Yao, Y., Matta, R., Manson, M. E., et al. (2006). MAP kinase phosphatase 1 controls innate immune responses and suppresses endotoxic shock. *J. Exp. Med.* 203, 131–140. doi: 10.1084/jem.20051794

- Zhong, H., Suyang, H., Erdjument-Bromage, H., Tempst, P., and Ghosh, S. (1997). The transcriptional activity of NF-kappaB is regulated by the IkappaB-associated PKAc subunit through a cyclic AMP-independent mechanism. *Cell* 89, 413–424. doi: 10.1016/S0092-8674(00)80222-6
- Zhu, Q. Y., Liu, Q., Chen, J. X., Lan, K., and Ge, B. X. (2010). MicroRNA-101 targets MAPK phosphatase-1 to regulate the activation of MAPKs in macrophages. *J. Immunol.* 185, 7435–7442. doi: 10.4049/jimmunol.1000798
- Ziegler-Heitbrock, L., Ancuta, P., Crowe, S., Dalod, M., Grau, V., Hart, D. N., et al. (2010). Nomenclature of monocytes and dendritic cells in blood. *Blood* 116, e74–e80. doi: 10.1182/blood-2010-02-258558

Conflict of Interest Statement: The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Copyright © 2016 Lloberas, Valverde-Estrella, Tur, Vico and Celada. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) or licensor are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.