

PDF hosted at the Radboud Repository of the Radboud University Nijmegen

The following full text is a publisher's version.

For additional information about this publication click this link.

<http://hdl.handle.net/2066/150832>

Please be advised that this information was generated on 2017-12-05 and may be subject to change.

Comparing Galactic Center MSSM dark matter solutions to the Reticulum II gamma-ray data

Abraham Achterberg,^a Melissa van Beekveld,^a Wim Beenakker,^{a,c}
Sascha Caron^{a,b} and Luc Hendriks^a

^aInstitute for Mathematics, Astrophysics and Particle Physics,
Faculty of Science, Mailbox 79, Radboud University Nijmegen,
P.O. Box 9010, NL-6500 GL Nijmegen, The Netherlands

^bNikhef, Science Park,
Amsterdam, The Netherlands

^cInstitute of Physics, University of Amsterdam,
Science Park 904, 1018 XE Amsterdam, The Netherlands

E-mail: a.achterberg@astro.ru.nl, mcbeekveld@gmail.com, wimb@hef.ru.nl,
scaron@cern.ch, luc.hendriks@gmail.com

Received September 15, 2015

Revised November 2, 2015

Accepted November 15, 2015

Published December 4, 2015

Abstract. Observations with the Fermi Large Area Telescope (LAT) indicate a possible small photon signal originating from the dwarf galaxy Reticulum II that exceeds the expected background between 2 GeV and 10 GeV. We have investigated two specific scenarios for annihilating WIMP dark matter within the phenomenological Minimal Supersymmetric Standard Model (pMSSM) framework as a possible source for these photons. We find that the same parameter ranges in pMSSM as reported by an earlier paper to be consistent with the Galactic Center excess, are also consistent with the excess observed in Reticulum II, resulting in a J -factor of $\log_{10}(J(\alpha_{\text{int}} = 0.5^\circ)) \simeq (20.3 - 20.5)_{-0.3}^{+0.2} \text{ GeV}^2 \text{ cm}^{-5}$. This J -factor is consistent with $\log_{10}(J(\alpha_{\text{int}} = 0.5^\circ)) = 19.6_{-0.7}^{+1.0} \text{ GeV}^2 \text{ cm}^{-5}$, which was derived using an optimized spherical Jeans analysis of kinematic data obtained from the Michigan/Magellan Fiber System (M2FS).

Keywords: dark matter theory, dark matter simulations, dwarfs galaxies

ArXiv ePrint: [1507.04644](https://arxiv.org/abs/1507.04644)



Contents

1	Introduction	1
2	Analysis set-up	2
3	Results	4
4	Summary and conclusions	6

1 Introduction

While the existence of dark matter (DM) is widely accepted and its cosmological abundance is measured, the fundamental nature of DM remains unknown. The most promising explanation is that DM is a neutral and weakly interacting particle, not described by the Standard Model (SM). A weakly interacting massive particle (WIMP) is a leading scenario for dark matter, as it can lead to the right abundance of DM originating from a thermal freeze out in the early universe. WIMPs can be searched for with several detection strategies, one of which being indirect detection via annihilation products (for example gamma rays) [1, 2].

Observations of our Galactic Center (GC) with the Fermi-LAT satellite over the past several years, show that there is a significant excess in gamma rays in the energy range of $1 \text{ GeV} \lesssim E_\gamma \lesssim 50 \text{ GeV}$ (see refs. [3–17] for details and interpretations). The source of these photons is under debate. Astrophysical explanations (like pulsars or cosmic ray outbursts) have been shown to face challenges, but it is not yet possible to rule out these explanations [18–24]. Another possibility for the origin of this γ -ray excess is annihilating DM. The GC is a promising target for searches of DM signals, as large-scale simulations of galaxy formation predict DM halos around galaxies such as the Milky way [25, 26]. This implies that an indirect DM signal could come from the Galactic Center [27].

Initially, the GC excess signal was reported to be compatible with a 30 GeV dark matter particle annihilating into $b\bar{b}$, or a 10 GeV dark matter particle annihilating to $\tau\bar{\tau}$ [5, 28]. The resulting b -quarks and τ -leptons can hadronize to neutral pions, which can decay to the observed γ -ray photons. Both processes should have an annihilation cross section close to $\langle\sigma v\rangle \simeq 1.3 \times 10^{-26} \text{ cm}^3\text{s}^{-1}$, the annihilation cross section expected for a thermal relic with a mass at the weak scale $E_w \simeq 100 \text{ GeV}$. These predictions have been explored within various contexts, for example within the Minimal Supersymmetric Standard Model (MSSM). In the framework of the MSSM, it is impossible to obtain such a scenario given LEP constraints [29].

However, a study on astrophysical background model systematics [13] allows for also higher WIMP masses and different annihilation channels to give a good fit to the GC excess [30]. In addition it is shown that uncertainties originating from high energy physics (especially partron showers) are also important to take into account [31]. Recently new annihilation scenarios within the phenomenological MSSM (pMSSM) and compatible with all the constraints are found [31, 32]. In the pMSSM the 105 Lagrangian parameters of the MSSM are reduced to 22, using phenomenological constraints on several parameters. In this model we assume that:

- Masses of the first and second generation sfermions are equal, separately in the lepton and quark sectors.

- All the soft SUSY-breaking parameters are real, so the only source for CP-violation is the CKM matrix in the standard model.
- Requiring minimal flavor violation, in the MSSM all sfermion mass matrices are assumed to be diagonal.

As a result of R-parity conservation in the MSSM, a lightest supersymmetric particle (LSP) exists. In most scenarios the LSP is the lightest neutralino, $\tilde{\chi}_1^0$, which is a combination of the neutral electroweak gaugino and higgsino fields. We select only models with a neutralino as LSP. The neutralino is a well-motivated DM candidate, as it is neutral, stable and can lead to a right DM relic density [33]. A 19-dimensional realization of the pMSSM is enough to encapsulate the phenomenology of the 22-parameter model. This is achieved by putting all trilinear couplings of the first and second generation to 0. The remaining parameters are 10 sfermion masses, 3 gaugino masses $M_{1,2,3}$, the ratio of the Higgs vacuum expectation values $\tan\beta$, the Higgsino mixing parameter μ , the mass m_A of the CP-odd Higgs-boson A^0 and 3 trilinear couplings $A_{b,t,\tau}$.

Dwarf galaxies (dSphs) in the vicinity of our galaxy provide an alternative to the Galactic Center for the search for γ -rays originating from annihilating dark matter. The amount of DM in these objects can be estimated by measuring the stellar velocities of member stars [34]. The resulting DM signal is proportional to the line-of-sight integral of the DM density distribution, a quantity known as the J -factor. Dwarf galaxies have a lower DM density (typical up to $\log_{10}(J) = 20 \text{ GeV}^2 \text{ cm}^{-5}$) than expected for the Galactic Center (for which the J -factor is predicted to be at least an order of magnitude larger). However, dSphs have less complicated backgrounds than the Galactic Center, as the latter suffers from systematic uncertainties associated with diffuse fore- and backgrounds [35]. These facts make dSphs promising targets for the detection of γ -rays originating from DM annihilation. Until recently, no dSphs with a significant γ -ray excess were found, setting strong limits on the cross section for DM annihilation. Recent imaging data from the Dark Energy Survey (DES) led to the discovery of new Milky Way satellites [36–38]. Among those, Reticulum II is of particular interest, as this object showed a gamma ray excess of a global significance level of 2.3–3.7 σ [39, 40] between 2 GeV and 10 GeV. The Fermi and DES collaborations found a less significant excess (p-value of 0.05 including a trials factor from testing multiple DM masses and channels, corresponding to a global significance of 1.65 σ), using an updated (Pass 8) data set [41]. The local significance they find is 2.4 σ ($p = 0.01$).

In this paper we will present a comparison of the Pass 7 data for Reticulum II in terms of annihilating neutralino dark matter. We use the pMSSM as a framework to provide this dark matter particle, using the results of ref. [31]. This study has no channel and mass trials factor since the DM mass and shape are fixed by the pMSSM model and the Galactic Center excess data. We will update this study to Pass 8 data in due course.

2 Analysis set-up

In ref. [40] events are included within 0.5° from Reticulum II. We adopt the same strategy and will refer to this as the region of interest (ROI). See figure 1 for the resulting photon spectrum, which is extracted from ref. [40]. The error bars indicate Poisson confidence level intervals (68%) on the number of counts in each bin. A logarithmic binning of 5 bins per decade between 0.2 and 300 GeV is used, resulting in a total of 15 bins. In this figure the last 2 bins are not shown, because of their large uncertainties. It is clear that between 2 and




Figure 1. Photon spectrum as extracted from ref. [40] of the Fermi-LAT data using events within 0.5° of Reticulum II (black points) with Poisson error bars. The solid gray line is the sum of background estimates of the Fermi Collaboration’s models for isotropic and Galactic diffuse emission at the location of Reticulum II. Between 2 GeV and 10 GeV, the photon spectrum rises above the background. This observed excess region is indicated by the blue box. The colored points are fluxes generated by DarkSUSY of the best fits for four scenarios: tt (blue), WW(1) (green), WW(2) (red) as extracted from [31] and bb (orange), added to the background of Reticulum II. The red box indicates the expected excess region, as determined by equation (3.2).

10 GeV, the spectrum of Reticulum II rises above the expected background. We will refer to this region as the observed excess region from now on. See ref. [40] for details on the analysis leading to this gamma-ray excess. From the ROI we derive the integration angle to be $\alpha_{\text{int}} = 0.5^\circ$, corresponding to $\Omega = 2\pi(1 - \cos \alpha_{\text{int}}) \simeq 2.4 \times 10^{-4}$ sr.

The observed gamma-ray flux from DM annihilation per unit solid angle at a photon energy E_γ is given by:

$$\frac{d\Phi_\gamma(E_\gamma)}{dE_\gamma d\Omega} = \frac{\langle \sigma v \rangle}{8\pi m_{\text{DM}}^2} \frac{dN_\gamma}{dE} \int_{\text{l.o.s.}} \rho_{\text{DM}}^2(l) dl, \quad (2.1)$$

where the integral over the DM density squared is along the line of sight (l.o.s.), $\langle \sigma v \rangle$ is the annihilation cross section weighted by the relative velocity, m_{DM} is the WIMP mass, and dN_γ/dE denotes the photon spectrum per annihilation.

The normalization factor $\int d\Omega \int_{\text{l.o.s.}} \rho_{\text{DM}}^2(l) dl$ is commonly referred to as the J -factor:

$$J = \int d\Omega \int_{\text{l.o.s.}} \rho_{\text{DM}}^2(l) dl \quad (2.2)$$

We use SUSPECT [42] as a spectrum generator and DarkSUSY 5.1.1 [43, 44] to compute the photon fluxes. In ref. [31], it is shown that there are three pMSSM parameter ranges that are consistent with the GC photon excess as originating from annihilating DM. That paper contains the details on how the possible solutions were found using a scan of the pMSSM

	M_1 (GeV)	M_2 (GeV)	μ (GeV)	$\tan \beta$
tt	171–189	190–1550	> 250	> 5
WW(1)	103–119	240–660	108–142	8–50
WW(2)	91–101	102–127	156–507	5–12

Table 1. Parameter ranges in pMSSM that correspond to the best solutions to the GC excess, as extracted from [31].

parameter space. The solutions presented in ref. [31] yield a value for the DM relic density that corresponds to $\Omega_{\text{DM}}h^2 \simeq 0.1$, close to the observed value $\Omega_{\text{DM}}h^2 = 0.12 \pm 0.0027$ [45], without being constrained a-priori to do so. In all cases the lightest neutralino is the DM candidate.

The first region of the pMSSM parameter space that was found to yield a good fit to the GC excess is one in which the lightest neutralinos annihilate mostly to a pair of top quarks. The lightest neutralinos have a very dominant bino component (99%), and their masses are at the kinematical threshold ($m_\chi \sim 174$ – 187 GeV). These models have $\Omega_{\text{DM}}h^2 = 0.08$ – 0.21 . We will refer to these models as tt.

The second solution is $\tilde{\chi}_1^0 \tilde{\chi}_1^0 \rightarrow W^+W^-$. In this case the lightest neutralino is a bino-higgsino mixture, with a mass range of $m_\chi \sim 84$ – 92 GeV. We will refer to these models as WW(1). This solution provided the best fit to the GC excess (p-value=0.45). These models correspond to a $\Omega_{\text{DM}}h^2$ in the range of 0.08–0.10.

The third solution yields a scenario in which the lightest neutralinos annihilate mostly to W^+W^- pairs. In this case, the composition of the lightest neutralinos is a mixture of bino, wino and higgsino, where the bino component is the most dominant one (90%). The mass of the lightest neutralinos is $m_\chi \sim 87$ – 97 GeV. The Galactic Center best fit points have an $\Omega_{\text{DM}}h^2$ of 0.07–0.18. We will refer to these models as WW(2).

In table 1 we show the exact parameter ranges in the pMSSM for each scenario. From each scenario we adopt the 50 best fit models (corresponding with the best χ^2 for the Galactic Center excess fit) for comparison with the Reticulum II signal.

In addition to these three scenarios, we create a 50 GeV neutralino and consider only the $b\bar{b}$ annihilation channel. We artificially put $\langle\sigma v\rangle \simeq 1.3 \times 10^{-26} \text{ cm}^3\text{s}^{-1}$, the upper limit for the annihilation cross section as found by ref. [46] and used in ref. [39] to explain the Reticulum II excess. We will refer to this model as bb.

3 Results

The resulting photon fluxes are fitted with the photon spectrum of Reticulum II, leaving the J -factor as a fitting parameter. Since the error bars are approximately symmetric and uncorrelated, we use the following χ^2 definition to determine the best fit:

$$\chi^2 = \sum_i \frac{(d_i - m_i)^2}{\sigma_i^2}, \quad (3.1)$$

where i is the energy bin number, running from 1 to 15, d_i is the Reticulum flux and m_i the DarkSUSY model flux. In figure 1 we show the resulting photon spectrum for the best fits. We use $(\text{minimal } \chi^2) + 1$ to define the 1σ range of the J -factor. In the following, we give the mean values of this J -factor error for all 50 best fit models within each scenario.




Figure 2. J -factor (note we do not use the logarithm) of the three best fits within tt (blue), $WW(1)$ (green), $WW(2)$ (red) and bb (orange) as a function of χ^2 (as defined in equation (3.1)).

We find that the tt scenarios yield the best results, with a minimal χ^2 ranging between 7.75 and 7.93. To determine the corresponding p-value we take a conservative approach and use 13 degrees of freedom (d.o.f.). The d.o.f. are counted as follows: there is a total of 15 bins, which gives 15 degrees of freedom. We subtract 1 d.o.f. for the background fit and 1 d.o.f. for the J -factor normalization. Using 13 d.o.f., the chi2-values correspond to a p-value of $\simeq 0.85$. The tt models give a J -factor of $\log_{10}(J(\alpha_{\text{int}})) = (20.33-20.56)_{-0.23}^{+0.15} \text{ GeV}^2 \text{ cm}^{-5}$. The numbers between brackets denote the range of the optimal fit J -factor for each model.

The $WW(1)$ scenario results in a J -factor of $\log_{10}(J(\alpha_{\text{int}})) = (20.31-20.35)_{-0.25}^{+0.16} \text{ GeV}^2 \text{ cm}^{-5}$. The chi2-values are slightly worse with a mean value of 8.75, corresponding to a p-value of 0.79 using 13 d.o.f.

The $WW(2)$ scenario gives a J -factor of $\log_{10}(J(\alpha_{\text{int}})) = (20.25-20.55)_{-0.25}^{+0.15} \text{ GeV}^2 \text{ cm}^{-5}$. These have a χ^2 between 8.35 and 8.66, corresponding to a p-value of 0.81.

The last scenario, bb , results in $\log_{10}(J(\alpha_{\text{int}})) = 19.57_{-0.35}^{+0.19}$ if we use an annihilation cross section of $\langle\sigma v\rangle \simeq 1.3 \times 10^{-26} \text{ cm}^3 \text{ s}^{-1}$. The bb scenario has $\chi^2 = 10.7$, which corresponds to a p-value of 0.63.

In figure 2 we show the J -factor as a function of χ^2 for one model in each scenario. Note that we do not plot the logarithm of the J -factor in this figure.

In table 2 we summarize the best p-values of the total signal and the p-values for the observed excess region for each scenario (indicated by *Reticulum II data* and *Observed excess region*). In order to provide for an unbiased statistical test of the signal shapes, we also include p-values for an expected excess region and for a background only model. We define the expected excess region as the range of bins where:

$$\min \{p_i(\text{signal} + \text{background} | \text{background})\} \quad (3.2)$$

is satisfied, where a minimum number of bins of 4 is demanded, because the observed excess region consists of 4 bins. The signal refers to the photon flux as generated by DarkSUSY. In

	Reticulum II data	Observed excess region	Expected excess region
tt	0.85	0.36	0.53
WW(1)	0.79	0.27	0.36
WW(2)	0.81	0.31	0.40
bb	0.63	0.09	0.24
background	0.37	0.01	0.03

Table 2. The p-values for the proposed four scenarios corresponding to the Reticulum II measured data, the observed excess region from 2 to 10 GeV (as indicated in figure 1 by the blue box) and the expected excess region (as indicated in figure 1 by the red box). In the last row we include p-values for the background only model to explain the Reticulum II measured data, the observed excess region and the expected excess region.

equation (3.2), the label i denotes a certain continuous bin range and p denotes the p-value. χ^2 is calculated according to equation (3.1) and the number of d.o.f. is equal to the number of bins - 1 (note that the J -factor normalization is now set, so we gain one extra d.o.f.). This range of bins corresponds to the situation where the best discrepancy between signal as generated by DarkSUSY and background can be made. We adopt the relative Poisson distributed errors on the Reticulum II measured data as an error on the background. The expected excess region is for all four models the same and indicated by the red box in figure 1. This region differs from the observed excess region by a few bins.

As shown in table 2, the tt scenario yields the best fit results for the observed excess region ($p=0.36$) and expected excess region ($p=0.53$). The bb scenario yields the worst fit ($p=0.09$ for the observed excess region, $p=0.24$ for the expected excess region).

Our conclusion is that the DM annihilation models, as proposed in ref. [31] which are consistent with the gamma ray excess in the Galactic Center as reported by Fermi-LAT, provide also a good fit to the observed gamma ray excess in Reticulum II. We find here that all these models predict a J -factor between $\log_{10}(J(\alpha_{\text{int}})) = 20.0 \text{ GeV}^2 \text{cm}^{-5}$ and $\log_{10}(J(\alpha_{\text{int}})) = 20.7 \text{ GeV}^2 \text{cm}^{-5}$ (including 1σ error). The bb model results in a J -factor of $\log_{10}(J(\alpha_{\text{int}})) = 19.57^{+0.19}_{-0.35}$. The GC models lie within the 1σ region of $\log_{10}(J(\alpha_{\text{int}})) = 19.6^{+1.0}_{-0.7} \text{ GeV}^2 \text{cm}^{-5}$ for an integration angle of $\alpha_{\text{int}} = 0.5^\circ$ as reported by ref. [47]. This derivation is done using an optimized spherical Jeans analysis of kinematic data obtained from the Michigan/Magellan Fiber System, an analysis which is independent of our approach. Ref. [48] reported a lower J -factor of $\log_{10}(J(\alpha_{\text{int}} = 0.5^\circ)) = 18.9 \pm 0.6 \text{ GeV}^2 \text{cm}^{-5}$ (1σ uncertainty), using the same spectroscopic data. This difference could originate from a different choice of dark matter priors (such as the dark matter local density and halo profile). Our models are consistent within 2σ with this J -factor.

4 Summary and conclusions

We found that the same models that are consistent with the Galactic Center gamma ray excess, shown in ref. [31], are also consistent with the small excess observed in Reticulum II, as reported by ref. [40]. The resulting J -factor is $\log_{10}(J(\alpha_{\text{int}} = 0.5^\circ)) \simeq (20.3 - 20.5)^{+0.2}_{-0.3} \text{ GeV}^2 \text{cm}^{-5}$, which is consistent with another determination of the J -factor using kinematic data obtained from the Michigan/Magellan Fiber System, resulting in a J -factor of $\log_{10}(J(\alpha_{\text{int}} = 0.5^\circ)) = 19.6^{+1.0}_{-0.7} \text{ GeV}^2 \text{cm}^{-5}$ as reported by ref. [47]. An improvement on the uncertainty of the J -factor will be important for interpreting these results.

Acknowledgments

We want to thank Alex Geringer-Sameth in providing the Reticulum II data and helping with problems.

References

- [1] G. Jungman, M. Kamionkowski and K. Griest, *Supersymmetric dark matter*, *Phys. Rept.* **267** (1996) 195 [[hep-ph/9506380](#)] [[INSPIRE](#)].
- [2] G. Bertone, D. Hooper and J. Silk, *Particle dark matter: Evidence, candidates and constraints*, *Phys. Rept.* **405** (2005) 279 [[hep-ph/0404175](#)] [[INSPIRE](#)].
- [3] L. Goodenough and D. Hooper, *Possible Evidence For Dark Matter Annihilation In The Inner Milky Way From The Fermi Gamma Ray Space Telescope*, [arXiv:0910.2998](#) [[INSPIRE](#)].
- [4] FERMI/LAT collaboration, V. Vitale and A. Morselli, *Indirect Search for Dark Matter from the center of the Milky Way with the Fermi-Large Area Telescope*, [arXiv:0912.3828](#).
- [5] D. Hooper and L. Goodenough, *Dark Matter Annihilation in The Galactic Center As Seen by the Fermi Gamma Ray Space Telescope*, *Phys. Lett. B* **697** (2011) 412 [[arXiv:1010.2752](#)] [[INSPIRE](#)].
- [6] D. Hooper and T. Linden, *On The Origin Of The Gamma Rays From The Galactic Center*, *Phys. Rev. D* **84** (2011) 123005 [[arXiv:1110.0006](#)] [[INSPIRE](#)].
- [7] K.N. Abazajian and M. Kaplinghat, *Detection of a Gamma-Ray Source in the Galactic Center Consistent with Extended Emission from Dark Matter Annihilation and Concentrated Astrophysical Emission*, *Phys. Rev. D* **86** (2012) 083511 [[arXiv:1207.6047](#)] [[INSPIRE](#)].
- [8] C. Gordon and O. Macias, *Dark Matter and Pulsar Model Constraints from Galactic Center Fermi-LAT Gamma Ray Observations*, *Phys. Rev. D* **88** (2013) 083521 [[arXiv:1306.5725](#)] [[INSPIRE](#)].
- [9] O. Macias and C. Gordon, *Contribution of cosmic rays interacting with molecular clouds to the Galactic Center gamma-ray excess*, *Phys. Rev. D* **89** (2014) 063515 [[arXiv:1312.6671](#)] [[INSPIRE](#)].
- [10] K.N. Abazajian, N. Canac, S. Horiuchi and M. Kaplinghat, *Astrophysical and Dark Matter Interpretations of Extended Gamma-Ray Emission from the Galactic Center*, *Phys. Rev. D* **90** (2014) 023526 [[arXiv:1402.4090](#)] [[INSPIRE](#)].
- [11] T. Daylan, D.P. Finkbeiner, D. Hooper, T. Linden, S.K.N. Portillo, N.L. Rodd et al., *The Characterization of the Gamma-Ray Signal from the Central Milky Way: A Compelling Case for Annihilating Dark Matter*, [arXiv:1402.6703](#) [[INSPIRE](#)].
- [12] B. Zhou, Y.-F. Liang, X. Huang, X. Li, Y.-Z. Fan, L. Feng et al., *GeV excess in the Milky Way: the role of diffuse galactic gamma-ray emission templates*, *Phys. Rev. D* **91** (2015) 123010 [[arXiv:1406.6948](#)] [[INSPIRE](#)].
- [13] F. Calore, I. Cholis and C. Weniger, *Background model systematics for the Fermi GeV excess*, *JCAP* **03** (2015) 038 [[arXiv:1409.0042](#)] [[INSPIRE](#)].
- [14] S. Murgia, *Observation of the high energy gamma-ray emission towards the Galactic center*, Talk given on *Fifth Fermi Symposium*, Nagoya, 20–24 October 2014.
- [15] D. Gaggero, M. Taoso, A. Urbano, M. Valli and P. Ullio, *Towards a realistic astrophysical interpretation of the Galactic center excess*, [arXiv:1507.06129](#) [[INSPIRE](#)].
- [16] I. Cholis, C. Evoli, F. Calore, T. Linden, C. Weniger and D. Hooper, *The Galactic Center GeV Excess from a Series of Leptonic Cosmic-Ray Outbursts*, [arXiv:1506.05119](#) [[INSPIRE](#)].

- [17] R.M. O’Leary, M.D. Kistler, M. Kerr and J. Dexter, *Young Pulsars and the Galactic Center GeV Gamma-ray Excess*, [arXiv:1504.02477](#) [INSPIRE].
- [18] I. Cholis, D. Hooper and T. Linden, *Challenges in Explaining the Galactic Center Gamma-Ray Excess with Millisecond Pulsars*, *JCAP* **06** (2015) 043 [[arXiv:1407.5625](#)] [INSPIRE].
- [19] D. Hooper, I. Cholis, T. Linden, J. Siegal-Gaskins and T. Slatyer, *Pulsars Cannot Account for the Inner Galaxy’s GeV Excess*, *Phys. Rev. D* **88** (2013) 083009 [[arXiv:1305.0830](#)] [INSPIRE].
- [20] J. Petrović, P.D. Serpico and G. Zaharijas, *Millisecond pulsars and the Galactic Center gamma-ray excess: the importance of luminosity function and secondary emission*, *JCAP* **02** (2015) 023 [[arXiv:1411.2980](#)] [INSPIRE].
- [21] J. Petrović, P.D. Serpico and G. Zaharijas, *Galactic Center gamma-ray “excess” from an active past of the Galactic Centre?*, *JCAP* **10** (2014) 052 [[arXiv:1405.7928](#)] [INSPIRE].
- [22] E. Carlson and S. Profumo, *Cosmic Ray Protons in the Inner Galaxy and the Galactic Center Gamma-Ray Excess*, *Phys. Rev. D* **90** (2014) 023015 [[arXiv:1405.7685](#)] [INSPIRE].
- [23] R. Bartels, S. Krishnamurthy and C. Weniger, *Strong support for the millisecond pulsar origin of the Galactic center GeV excess*, [arXiv:1506.05104](#) [INSPIRE].
- [24] S.K. Lee, M. Lisanti, B.R. Safdi, T.R. Slatyer and W. Xue, *Evidence for Unresolved Gamma-Ray Point Sources in the Inner Galaxy*, [arXiv:1506.05124](#) [INSPIRE].
- [25] J.F. Navarro, C.S. Frenk and S.D.M. White, *A Universal density profile from hierarchical clustering*, *Astrophys. J.* **490** (1997) 493 [[astro-ph/9611107](#)] [INSPIRE].
- [26] M. Boylan-Kolchin, V. Springel, S.D.M. White, A. Jenkins and G. Lemson, *Resolving Cosmic Structure Formation with the Millennium-II Simulation*, *Mon. Not. Roy. Astron. Soc.* **398** (2009) 1150 [[arXiv:0903.3041](#)] [INSPIRE].
- [27] I.V. Moskalenko, T.A. Porter and A.W. Strong, *Attenuation of the gamma rays by the milky way interstellar radiation field*, *Astrophys. J.* **640** (2006) L155 [[astro-ph/0511149](#)] [INSPIRE].
- [28] D. Hooper and T. Linden, *On The Origin Of The Gamma Rays From The Galactic Center*, *Phys. Rev. D* **84** (2011) 123005 [[arXiv:1110.0006](#)] [INSPIRE].
- [29] M. Cahill-Rowley, J. Gainer, J. Hewett and T. Rizzo, *Towards a Supersymmetric Description of the Fermi Galactic Center Excess*, *JHEP* **02** (2015) 057 [[arXiv:1409.1573](#)] [INSPIRE].
- [30] P. Agrawal, B. Batell, P.J. Fox and R. Harnik, *WIMPs at the Galactic Center*, *JCAP* **05** (2015) 011 [[arXiv:1411.2592](#)] [INSPIRE].
- [31] A. Achterberg, S. Amoroso, S. Caron, L. Hendriks, R. Ruiz de Austri and C. Weniger, *A description of the Galactic Center excess in the Minimal Supersymmetric Standard Model*, *JCAP* **08** (2015) 006 [[arXiv:1502.05703](#)] [INSPIRE].
- [32] G. Bertone, F. Calore, S. Caron, R.R. de Austri, J.S. Kim, R. Trotta et al., *Global analysis of the pMSSM in light of the Fermi GeV excess: prospects for the LHC Run-II and astroparticle experiments*, [arXiv:1507.07008](#) [INSPIRE].
- [33] M. van Beekveld, W. Beenakker, S. Caron, R. Castelijn, M. Lanfermann and A. Strubig, *Higgs, di-Higgs and tri-Higgs production via SUSY processes at the LHC with 14 TeV*, *JHEP* **05** (2015) 044 [[arXiv:1501.02145](#)] [INSPIRE].
- [34] V. Bonnivard et al., *Dark matter annihilation and decay in dwarf spheroidal galaxies: The classical and ultrafaint dSphs*, *Mon. Not. Roy. Astron. Soc.* **453** (2015) 849 [[arXiv:1504.02048](#)] [INSPIRE].
- [35] L.E. Strigari, *Galactic Searches for Dark Matter*, *Phys. Rept.* **531** (2013) 1 [[arXiv:1211.7090](#)] [INSPIRE].
- [36] DARK ENERGY SURVEY collaboration, T. Abbott et al., *The dark energy survey*, [astro-ph/0510346](#) [INSPIRE].

- [37] S.E. Kopusov, V. Belokurov, G. Torrealba and N.W. Evans, *Beasts of the Southern Wild: Discovery of nine Ultra Faint satellites in the vicinity of the Magellanic Clouds*, *Astrophys. J.* **805** (2015) 130 [[arXiv:1503.02079](#)] [[INSPIRE](#)].
- [38] DES collaboration, K. Bechtol et al., *Eight New Milky Way Companions Discovered in First-Year Dark Energy Survey Data*, *Astrophys. J.* **807** (2015) 50 [[arXiv:1503.02584](#)] [[INSPIRE](#)].
- [39] D. Hooper and T. Linden, *On The Gamma-Ray Emission From Reticulum II and Other Dwarf Galaxies*, *JCAP* **09** (2015) 016 [[arXiv:1503.06209](#)] [[INSPIRE](#)].
- [40] A. Geringer-Sameth, M.G. Walker, S.M. Koushiappas, S.E. Kopusov, V. Belokurov, G. Torrealba et al., *Indication of Gamma-ray Emission from the Newly Discovered Dwarf Galaxy Reticulum II*, *Phys. Rev. Lett.* **115** (2015) 081101 [[arXiv:1503.02320](#)] [[INSPIRE](#)].
- [41] DES, FERMI-LAT collaboration, A. Drlica-Wagner et al., *Search for Gamma-Ray Emission from DES Dwarf Spheroidal Galaxy Candidates with Fermi-LAT Data*, *Astrophys. J.* **809** (2015) L4 [[arXiv:1503.02632](#)] [[INSPIRE](#)].
- [42] A. Djouadi, J.-L. Kneur and G. Moultaka, *SuSpect: A Fortran code for the supersymmetric and Higgs particle spectrum in the MSSM*, *Comput. Phys. Commun.* **176** (2007) 426 [[hep-ph/0211331](#)] [[INSPIRE](#)].
- [43] P. Gondolo, J. Edsjo, P. Ullio, L. Bergstrom, M. Schelke and E.A. Baltz, *DarkSUSY: Computing supersymmetric dark matter properties numerically*, *JCAP* **07** (2004) 008 [[astro-ph/0406204](#)] [[INSPIRE](#)].
- [44] P. Gondolo, et al., <http://www.darksusy.org>.
- [45] PLANCK collaboration, P.A.R. Ade et al., *Planck 2015 results. XIII. Cosmological parameters*, [arXiv:1502.01589](#) [[INSPIRE](#)].
- [46] FERMI-LAT collaboration, M. Ackermann et al., *Searching for Dark Matter Annihilation from Milky Way Dwarf Spheroidal Galaxies with Six Years of Fermi-LAT Data*, [arXiv:1503.02641](#) [[INSPIRE](#)].
- [47] V. Bonnavard, C. Combet, D. Maurin, A. Geringer-Sameth, S.M. Koushiappas, M.G. Walker et al., *Dark matter annihilation and decay profiles for the Reticulum II dwarf spheroidal galaxy*, *Astrophys. J.* **808** (2015) L36 [[arXiv:1504.03309](#)] [[INSPIRE](#)].
- [48] DES collaboration, J.D. Simon et al., *Stellar Kinematics and Metallicities in the Ultra-Faint Dwarf Galaxy Reticulum II*, *Astrophys. J.* **808** (2015) 95 [[arXiv:1504.02889](#)] [[INSPIRE](#)].