

PDF hosted at the Radboud Repository of the Radboud University Nijmegen

The following full text is a publisher's version.

For additional information about this publication click this link.

<http://hdl.handle.net/2066/153062>

Please be advised that this information was generated on 2017-12-05 and may be subject to change.



POSTER PRESENTATION

Open Access

Immune infiltrates impact on the prediction of prognosis and response to immunotherapy of melanoma patients

Angela Vasaturo^{1*}, Dagmar Verweij¹, Lucie Heinzerling², Jolanda de Vries¹, Willeke Blokx³, Carl G Figdor¹

From Melanoma Bridge Meeting 2014
Naples, Italy. 03-06 December 2014

Background

Melanoma is a highly malignant melanocyte-derived tumor and its incidence is increasing at outstanding rate. Despite specific therapies have been explored for many years, no effective therapeutic options have been developed. Vaccination strategies, including Dendritic Cells (DC) based immunotherapy, are consistently increasing the proportion of cancer patients with anti-vaccine immune responses although the number of patients with increased overall survival is still limited. The efficacy of the immunotherapy is mainly dependent on tumor microenvironment –immune system interactions.

Our aim is to evaluate the host immune response to melanoma by quantifying the density and location of T cells in primary tumors of patients treated with DC immunotherapy and correlate them with clinical variables such as overall survival (OS).

Materials and methods

We collected 60 FFPE primary tumors from melanoma patients treated with DC immunotherapy. Serial sections (4 micron in thickness) were stained with CD3, CD8 and CD45RO antibodies. Haematoxylin was used as a counterstain and Nova Red for the immunohistochemical stain. All the slides were digitalized and an automated quantitative analysis was performed in order to evaluate the density and location of two lymphocyte populations, cytotoxic (CD8) and memory (CD45RO) T cells. Of all samples the clinical outcome of the patient is known.

Results

The immunohistochemical analysis of primary melanoma using a small set of patients resulted in significant differences between short (OS<12 months) and long survivors (OS>24 months). A high degree of T cells infiltration was seen in the tumor area of both patients, responding and non-responding to DC immunotherapy. However, the location but not the density of TILs was significantly different in the two cohorts of patients and showed a strong correlation with clinical response to DC vaccination.

Conclusions

Immune cells within melanoma tumors may have a prognostic value and clinical significance as a predictor of patient outcome and response to DC immunotherapy in melanoma patients.

Authors' details

¹Department of Tumor Immunology, Radboud Institute for Molecular Life Sciences, Nijmegen, Netherlands. ²Department of Dermatology, University Hospital of Erlangen, erlangen, Germany. ³Department of Pathology, Radboud University Medical Center, Nijmegen, The Netherlands.

Published: 15 January 2015

doi:10.1186/1479-5876-13-S1-P12

Cite this article as: Vasaturo et al.: Immune infiltrates impact on the prediction of prognosis and response to immunotherapy of melanoma patients. *Journal of Translational Medicine* 2015 **13**(Suppl 1):P12.

¹Department of Tumor Immunology, Radboud Institute for Molecular Life Sciences, Nijmegen, Netherlands

Full list of author information is available at the end of the article