

## PROCEEDINGS OF THE I INTERNATIONAL MEETING IN EXERCISE PHYSIOLOGY

**Comparative analysis of cardiopulmonary responses of healthy sedentary men and men after acute myocardial infarction #73**

Vandeni C. Kunz<sup>1</sup>, Karina B. Serra<sup>2</sup>; Erica N. Borges<sup>2</sup>, Ana Cristina S. Rebelo<sup>1</sup>, Roberta S. Zuttin<sup>2</sup>, Nayara Y. Tamburus<sup>2</sup>, Raquel B. Souza<sup>2</sup>, Marlene A. Moreno<sup>2</sup>, Paulo E. Serra<sup>3</sup>, Luiz A. Gubolino<sup>3</sup>, Nicolau A. Merino<sup>3</sup>, Luciano D. Dantas<sup>4</sup>, Ester da Silva<sup>1,2</sup>.

<sup>1</sup>Department of Physiotherapy, Federal University of São Carlos São Carlos, SP, Brazil; <sup>2</sup>Cardiovascular Physiotherapy and Functional Tests Research Laboratory, Faculty of Health Sciences, Methodist University of Piracicaba, Piracicaba, SP, Brazil; <sup>3</sup>Department of Cardiology, Hospital dos Fornecedores de Cana de Piracicaba, Piracicaba, SP, Brazil; <sup>4</sup>Department of Cardiology, Santa Casa de Misericórdia de Limeira, Limeira, SP, Brazil.

E-mail: [ester.silva@pq.cnpq.br](mailto:ester.silva@pq.cnpq.br)

The aim of this study was to assess the aerobic capacity of men after acute myocardial infarction (G-AMI) and of healthy sedentary men (G-C) at the anaerobic threshold (AT) and at the peak of the exercise (up to physical exhaustion) during a ramp-type ergospirometric test (R-ET). Methods: 22 volunteers divided into two groups: G-AMI (n=10), 55.6 ± 9.7 years old, undergoing β-blocking therapy (atenolol, dosage 46 ± 9.4 mg/day), Killip classification grade I, and G-C (n=12), 53.3 ± 3.2 years old. The G-C subjects took no type of medication. Experimental protocol: R-ET, with power increments of 10W/min for the G-AMI and of 15W/min for the G-C. The evaluated variables were: heart rate (HR), ventilatory and metabolic, recorded breath-by-breath using an ergospirometer (CPX/D MedGrafics - Breeze, St. Paul, Minnesota, USA). Statistical Analysis: Mann-Whitney tests α = 5%. The power, cardiovascular, ventilatory and metabolic variables obtained from the R-ET at the peak of the exercise, for G-AMI and G-C, presented a statistically significant difference (p<0.05), showing higher values for G-C, except the VO<sub>2</sub> in L/min. However, the variables showed similar values at the AT (p>0.05). Conclusion: The lower aerobic capacity at the peak of the exercise can be attributed to the use of β-blocking therapy and to prolonged bed rest.

**Key words:** acute myocardial infarction; heart rate; men.