

### Relationships between peak oxygen uptake and arterial function: a preliminary study.

Pearman, M., DeVallance, E., Fournier, S., Bonner, D., Warden, B.E., Gharib, W., Donley, D., Chantler, P. West Virginia University, Morgantown, WV

[mpearman2@mix.wvu.edu](mailto:mpearman2@mix.wvu.edu), [Edevalla@mix.wvu.edu](mailto:Edevalla@mix.wvu.edu), [sfournier@hsc.wvu.edu](mailto:sfournier@hsc.wvu.edu), [dbonner@hsc.wvu.edu](mailto:dbonner@hsc.wvu.edu), [bwarden@hsc.wvu.edu](mailto:bwarden@hsc.wvu.edu), [wgharib@hsc.wvu.edu](mailto:wgharib@hsc.wvu.edu), [ddonley@hsc.wvu.edu](mailto:ddonley@hsc.wvu.edu), [pchantler@hsc.wvu.edu](mailto:pchantler@hsc.wvu.edu)

**INTRODUCTION:** Aerobic fitness, as measured by  $VO_{2peak}$ , is a well-validated predictor of morbidity and mortality in healthy individuals and those with cardiovascular disease. Resting arterial function, as indicated by carotid to femoral pulse wave velocity (cfPWV: arterial stiffness), and arterial structure as indicated by intima-media thickness (IMT), and carotid diameter (Dc), are correlated to cardiovascular risk. As alterations in arterial health likely impact aerobic capacity, identifying arterial parameters that predict cardiovascular fitness are necessary; however, this relationship has not been fully investigated. **PURPOSE:** This preliminarily cross-sectional study examined the relationship between relative peak  $VO_2$  and resting arterial function of 51 subjects without overt cardiovascular disease (CVD) (mean age  $44.9 \pm 11.6$ ). **METHODS:**  $VO_{2peak}$  was assessed using a staged graded exercise test on a semi-recumbent bicycle until volitional fatigue. cfPWV was measured with Applanation tonometry; IMT and Dc were measured with B mode ultrasound. To account for individual differences in CVD risk each individual was assigned a metabolic risk score based on their age, sex, blood pressure, BMI, triglycerides, HDL, and glucose. **RESULTS:** Univariate regression models indicated that  $VO_{2peak}$  is significantly related to cfPWV [ $r = -0.43$ ,  $p = 0.002$ ], IMT [ $r = -0.46$ ,  $p = 0.001$ ], and Dc [ $r = -0.30$ ,  $p = 0.035$ ]. For multivariate analysis adjusted for age, sex, and metabolic risk scores,  $VO_{2peak}$  remained significantly related to cfPWV [ $r = -0.29$ ,  $p = 0.036$ ], IMT [ $r = -0.32$ ,  $p = 0.012$ ], and Dc [ $r = -0.25$ ,  $p = 0.037$ ]. **CONCLUSION:** Preliminary data suggest that peak oxygen uptake is independently correlated with arterial health in populations free of overt CVD. These results suggest that improving peak oxygen uptake may have a favorable effect on arterial function and vice versa.

Research funded in part by the American Heart Association 11CRP7370056 and National Heart, Lung, Blood Institute T32- HL090610