



INTERNATIONAL SOCIETY OF EXERCISE AND IMMUNOLOGY  
**11th ISEI SYMPOSIUM, NEWCASTLE, AUSTRALIA, 9-12 September 2013**  
 Exercise Immunology: Prescriptions for Health

**ABSTRACT FORM**

This form should be read in conjunction with the **ISEI ABSTRACT GUIDELINES**.

These 2 pages have been provided separately to assist you with the submission of your abstracts in Word format to the Conference Organising Committee and online to the **International Journal of Exercise Science (IJES)**.

You **MUST** complete this entire Abstract Form to provide us with details needed to assign your abstract to the correct theme.

Also advise us if you are applying to be considered for one of the Early Career Researcher Awards (Best Poster and Best Oral Presentation).

Send this entire form by E-mail to the ISEI Organising Committee, but **ALSO** follow the guidelines for submission of the Abstract (single page only) to the **IJES**.

**ABSTRACT SUBMISSION – DEADLINE 10 May 2013**

<b>Title</b> (up to 30 words, Arial, 11 pt, single line spaced, in sentence case. Like this:	Contribution of <i>Lactobacillus casei</i> to the recovery from chemically induced skeletal muscle damage under chronic stress
<b>Authors</b> (Underline the <u>presenting author</u> )	<u>K Matsuo</u> , I Tabata & R Nagatomi Must also be exactly as submitted on-line to <i>IJES</i>
<b>Department, Institution, Country</b>	1. Center of General Education, Tohoku Institute of Technology, Japan 2. Division of Biomedical Engineering for Health and Welfare, Tohoku University Graduate School of Medicine, Japan 3. Faculty of Sports and Health Science, Ritsumeikan University, Japan
<b>Address Corresponding Author</b>	35-1 Yagiyama, Kasumimachi, Taihaku-ku, Sendai, Miyagi, 982-8577 JAPAN k-matsuo@tohtech.ac.jp
<b>Select Your Abstract Session Theme Category</b>	Exercise prescriptions for a healthy gut - 4
<b>Preferred Presentation Form</b>	<input type="checkbox"/> Oral communication <input checked="" type="checkbox"/> Poster
<b>Is the presenter eligible for the Early Career Researcher Awards (poster and oral awards)?</b>  (Previous winners are ineligible for same category)	<input type="checkbox"/> Yes <input checked="" type="checkbox"/> No  Eligible persons are those studying for a higher degree – MSc, MPhil or PhD – or who have completed their PhD within the last 3 years as at 10th September 2013



**ISEI Abstract – Please do not change any fonts, font sizing, or spacing!  
Replace this wording with your selected “Session theme number – 4”**

---

## **Contribution of *Lactobacillus casei* to the recovery from chemically induced skeletal muscle damage under chronic stress**

MATSUO K 1,2, TABATA I 3, and NAGATOMI R 2.

1 Center of General Education; Human Science; Tohoku Institute of Technology; Sendai, JAPAN.

2 Department of Medicine and Science in Sports and Exercise; Tohoku University, Sendai, JAPAN.

3 Faculty of Sports and Health Science, Ritsumeikan University, Kusatsu, JAPAN.

---

### ABSTRACT

**Background:** Regeneration of damaged skeletal muscle requires sufficient supply of nutrients. Fully functional intestine and colon assure sufficient supply of nutrients. Gut commensal bacteria are known to support intestinal function. Previously lactobacillus treatment of elite athletes was shown to be effective in attenuating fatigue and impaired performance. We hypothesized that *Lactobacillus casei* (*L.casei*) administration may facilitate recovery of damaged skeletal muscle when the gut function is suppressed under chronic stress in which muscle regeneration is compromised.

**Objective:** To investigate the contribution of *L.casei* under chronic stress to the recovery of damaged skeletal muscle in young and older adult mice.

**Methods:** *L.casei* was given orally at a dose of  $10^8$  /day for 7days to 10 weeks old (young) and 45-55 weeks old (older adult) male C57BL/6J mice. Vehicle control mice received an equivalent volume of water for 7 days. On the eighth day, cardiotoxin (CTX) was injected to gastrocnemius muscle to induce muscle damage. Both groups were assigned 2 hours repeated-restraint stress everyday (chronic stress). On days 3, 5, 7, 10, 14 and 20 after CTX injection, mice were sacrificed. Excised gastrocnemius muscle was subjected to weight measurement and immunohistochemical analyses.

**Results:** There were significant differences in both the recovery of muscle weight and the regeneration process of gastrocnemius muscle examined immunohistochemically between control and *L.casei* treated young and older adult groups. Especially, the expression of developmental MHC (dMHC), a marker of premature regeneration, was positive up to 3 days in older adult groups. The delay in the recovery of muscle weight was obvious in older adult mice regardless of the treatment. However, while the expression of dMHC was prolonged up to day 7 in the vehicle control, dMHC expression was notable only up to day 3 or day 5 in the *L.casei* treated. Therefore, in older adult mice *L.casei* treatment under chronic stress may have facilitated the maturation process of regenerating skeletal muscle.

**Conclusion:** Our result suggests that *L. casei* favor under chronic stress favors the recovery of skeletal muscle from muscle damage. The maintenance of gut function by *L. casei* treatment may have facilitated the maturation process of regenerating skeletal muscle.

**Acknowledgments:** This study was supported by Yakult Bio-Science Foundation, the Uehara Memorial Foundation, Ritsumeikan University Research Grant and Grant-in-Aid for Young Scientific Research (B), the Ministry of Education, Culture, Sports science and technology.