

## **Energy Expenditure following Acute Cold Exposure**

Steven Stelly, Danielle Bravo, Natalie Hines, Loren Koehler, Michael Levi and Donovan Fogt

Exercise Biochemistry & Metabolism Laboratory; Department of Kinesiology, Health & Nutrition; The University of Texas at San Antonio; San Antonio, TX

---

*Category: Masters*

*Advisor / Mentor: Fogt, Donovan (donovan.fogt@utsa.edu)*

### ABSTRACT

Whole body energy expenditure (i.e., RMR) increases during acute cold exposure. Whether this increase in energy expenditure persists in the post-cold term in humans due to non-shivering thermogenesis has not yet been evaluated. Therefore we tested the hypothesis that RMR would be different directly following acute cold exposure. RMR ( $\text{kcal} \cdot \text{min}^{-1}$ ) was assessed via indirect calorimetry on eight (8) males and seven (7) females at six time points: prior to cold exposure (T1), at the end of 30 minutes of cold exposure (T2), immediately post-cold-post-shivering (T3), at 35 minutes post-cold (T4), at 75 minutes post-cold (T5), and at 115 minutes post-cold (T6). This RMR data for the aforementioned time points was analyzed using paired, dependent t-tests and one way ANOVA; the significance level was placed at  $p < 0.05$ . The RMR data for each time point was as follows: T1 ( $1.19 \pm 0.21$ ), T2 ( $2.30 \pm 0.94$ ), T3 ( $1.37 \pm 0.25$ ), T4 ( $1.12 \pm 0.19$ ), T5 ( $1.14 \pm 0.22$ ), and T6 ( $1.14 \pm 0.22$ ). The analysis of the RMR data showed a significant difference between the cold (T2) data and all other time points (T1, T3, T4, T5, and T6). Additionally there was a significant difference between the pre-cold (T1) and the immediately post-cold (T3) data. However there was no significant difference between the pre-cold (T1) data when compared with the remaining post-cold data (T4 – T6). These results suggest that the human body is capable of returning RMR to baseline levels relatively immediately following the cessation of acute cold exposure.