

Relationship Among Waist Girth, Body Mass Index, Total Cholesterol in College-Aged Males

Miltenberger, M, Papalia, Z, DiLeo, T, Bopp, C. Department of Kinesiology, Pennsylvania State University, University Park, PA

ACSM risk stratification includes fasting glucose and lipid measures. Assessment of these risk factors often requires specialized equipment that may not be available in all settings. **PURPOSE:** To determine what relationship, if any, exists between waist girth (WG) and body mass index (BMI) in individuals with elevated total cholesterol (tCHO) ≥ 200 mg/dL. A secondary purpose was to determine if the addition of cholesterol testing to our screening procedures changed risk status in our participants. **METHODS:** A blood sample of 40 μ L was obtained from the participants' distal phalange using a single use lancet. The sample was placed in a lipid/glucose cassette and analyzed by a commercially available analyzer. BMI was calculated from height and weight collected using standard procedures. WG was measured with a standard tension tape measure. Correlations between variables was assessed with Pearson product moment correlation. **RESULTS:** The males ($N=666$) tested for the study had an average age of 21.92 years. 9.9% of the test subjects ($n=66$) had a tCHO value ≥ 200 mg/dL. A strong linear correlation was observed between Wg and BMI ($r=0.64$). No relationship was observed between tCHO and BMI ($r=0.16$) or WG($r=.23$). Among our subjects with elevated cholesterol, tCHO on average was 257mg/dL, WG on average was 37 inches, and a BMI average of 26.25 kg/m². **CONCLUSION:** Adding tCHO testing to the screening procedures identifies additional participants not captured by the remaining risk factors. Despite the cost, fasting cholesterol assessments should be performed to get a complete risk factor profile on all college aged subjects.