

Supporting Information

Joint experimental and theoretical study on vibrational excitation cross sections for electron collisions with diacetylene

Roman Čurík^{1*}, Ivana Paidarová¹, Michael Allan², and Petr Čársky^{1*}

¹J. Heyrovský Institute of Physical Chemistry, Academy of Sciences of the Czech Republic, v.v.i, Dolejškova 3, 18223 Prague 8, Czech Republic

²Department of Chemistry, University of Fribourg, Chemin du Musée 9, CH-1700 Fribourg, Switzerland

*Corresponding authors; e-mail: roman.curik@jh-inst.cas.cz, carsky@jh-inst.cas.cz

Contents

A. Results of calculations

S1. Calculated energy dependences of vibrationally inelastic differential cross sections for nine normal modes of diacetylene at 45°.


S2. Calculated energy dependences of vibrationally inelastic differential cross sections for nine normal modes of diacetylene at 135°.

B. Experimental results


S3. Overview of energy-loss spectra recorded at the incident energy of 1 eV at six different scattering angles.

S4. Overview of the cross sections recorded at 135° for excitation functions. The curves were recorded at energy-losses of 0, 27, 78, and 155 meV.


S5. Continuation of figure S4 for higher-lying vibrational states.




S1. Calculated energy dependences of vibrationally inelastic differential cross sections for nine normal modes of diacetylene at 45°.




S2. Calculated energy dependences of vibrationally inelastic differential cross sections for nine normal modes of diacetylene at 135° .



S3. Overview of energy-loss spectra recorded at the incident energy of 1 eV, within the ${}^2\Pi_u$ resonance, at 6 different scattering angles. The spectra at 0° , 90° and 180° were already published in ref 1 and are reproduced here for a global comparison. The spectra illustrate the rich excitation of overtone and combination vibrations mentioned in the main text, which is due to the narrow autodetachment width of the ${}^2\Pi_u$ resonance. An interesting point is the increasing density of high overtones with low-frequency, observed at high energy losses (i.e., low scattered electron energies) and angles of 45° and higher. It is a manifestation of the ‘unspecific’ vibrational excitation, related to IVR. Another interesting point is the preference for exciting double quanta of the bending vibrations $\nu_{6,8}$, particularly clear in combination with ν_2 , i.e., $\nu_2 + \nu_{6,8}$ is weak at 90° and 135° and practically missing at the other angles, whereas $\nu_2 + 2\nu_{6,8}$ is strong.¹



S4. Overview of the cross sections recorded at 135° . The red lines are excitation functions; they are normalized to the individual absolute measurements, indicated by yellow circles. The curves recorded at energy-losses of $\Delta E = 27$ meV and $\Delta E = 78$ meV have threshold peaks, presumably a consequence of the IR activity of the ν_9 (weak) and ν_8 (strong) normal modes. The cross section for the overtone at $\Delta E = 155$ meV, only weakly IR active, has only a weak threshold peak. Note that the $^2\Pi_u$ resonance at 1 eV has a deep boomerang structure in the overtone excitation at $\Delta E = 155$ meV, but no boomerang structure in the excitation of the fundamental at $\Delta E = 78$ meV. This reflects the entirely different excitation mechanism, revealed also by the dramatically different angular distributions.¹ The $^2\Pi_g$ resonance at 5.3 eV is much weaker, relatively to the $^2\Pi_u$ resonance at 1 eV, in the overtone excitation than in the fundamental excitation, reflecting its much larger autodetachment width. This is already indicated by the absence of boomerang structure.



S5. Continuation of S4, for higher-lying vibrational states. Note that the 6.5 eV resonance excites overtones such as $2\nu_2$ more weakly than the longer-lived 1.0 eV resonance. The stronger excitation of double quanta of $2\nu_{6,8}$ is also seen.

1. M. Allan, O. May, J. Fedor, B. C. Ibănescu and L. Andric, *Phys. Rev. A* **2011**, 83, 052701.