

X-ray two-photon absorption with high fluence XFEL pulses

J. Hoszowska*¹, J. Szlachetko^{†a}, J.-Cl. Dousse*, W. Błachucki*, Y. Kayser[‡], Ch. Milne[‡],
M. Pajek^a, S. Boutet^b, M. Messerschmidt^b, G. Williams^b, and C.T. Chantler^c

* Department of Physics, University of Fribourg, CH-1700 Fribourg, Switzerland

[†] Paul Scherrer Institut, CH-5232 Villigen PSI, Switzerland

^a Institute of Physics, Jan Kochanowski University, 25-406 Kielce, Poland

^b Linac Coherent Light Source (LCLS), SLAC National Accelerator Laboratory, 94025 California, USA

^c School of Physics, University of Melbourne, Parkville, Victoria 3010, Australia

Synopsis. We report on nonlinear interaction of solid Fe with intense femtosecond hard x-ray free-electron laser (XFEL) pulses. The experiment was performed at the CXI end-station of the Linac Coherent Light Source (LCLS) by means of high-resolution x-ray emission spectroscopy. The focused x-ray beam provided extreme fluence of $\sim 10^5$ photons/ \AA^2 . Two-photon absorption leading to K-shell hollow atom formation and to single K-shell ionization of solid Fe was investigated.

X-ray free electron laser (XFEL) facilities, with unprecedentedly high peak power densities reaching $\sim 10^{20}$ W/cm², have paved the way to study nonlinear phenomena in the x-ray regime [1-6]. In this work we explored nonlinear interaction of high-fluence hard x-ray femtosecond pulses with solid Fe. Single and double K-shell electron ionization processes resulting from two-photon absorption were observed.

The experiment was carried out at the CXI end-station of the Linac Coherent Light Source (Menlo Park, USA) XFEL by means of the high energy resolution x-ray emission technique. The XFEL beam of $\sim 5 \times 10^{11}$ x-rays/pulse and pulse energy of 0.6 mJ was focused on a metallic Fe sample. The ultra-focused x-ray beam provided extreme fluence of $\sim 10^5$ photons/ \AA^2 . Moving the sample out of the focus along the beam allowed varying the fluence. For the Fe $K\alpha$ ($K^{-1} \rightarrow L^{-1}$) and $K\alpha^h$ ($K^{-2} \rightarrow K^{-1}L^{-1}$) radiative transitions measurements the bent crystal von Hamos x-ray spectrometer of PSI [7] installed at CXI and equipped with the CSPAD detector developed at SLAC was employed. The K x-ray emission spectra were collected at photon beam energies below the Fe K-shell single- and double-ionization thresholds for the two-photon single ionization and double ionization processes, respectively.

For illustration, the probability of double K-hole formation *via* sequential absorption of two photons *versus* x-ray fluence is shown in figure 1. We observe a ~ 60 -fold increase in the production probability of Fe hollow-atoms as compared to single-photon double ionization mediated by K-shell electron-electron correlations [8]. The cross-sections for double-K-hole formation and two-photon single K-shell ionization were derived from the x-ray fluence dependence of the measured x-ray emission intensities. For the two-photon single ionization process a square dependence of the $K\alpha$ signal was found.

This is the first observation of K-shell double core-hole creation following sequential photon absorption, and two-photon single K-shell ionization for metallic Fe.




Figure 1. Probability of double K-hole formation *via* sequential absorption of two photons for Fe as a function of x-ray fluence. The data were derived from the $K\alpha^h$ ($K^{-2} \rightarrow K^{-1}L^{-1}$) to $K\alpha$ ($K^{-1} \rightarrow L^{-1}$) intensity ratios. The x-ray pulse energy was 7.6 keV and the duration 30 fs.

References

- [1] L. Young *et al* 2010 *Nature* **466** 56
- [2] G. Doumy *et al* 2011 *Phys. Rev. Lett.* **106** 083002
- [3] B. Rudek *et al* 2012 *Nat. Photon.* **6** 858
- [4] K. Tamasaku *et al* 2013 *Phys. Rev. Lett.* **111** 043001
- [5] K. Tamasaku *et al* 2014 *Nat. Photon.* **8** 313
- [6] H. Yoneda *et al* 2014 *Nat. Commun.* **5** 5080
- [7] J. Szlachetko *et al* 2012 *Rev. Sci. Instrum.* **83** 103105
- [8] J. Hoszowska *et al* 2009 *Phys. Rev. Lett.* **102** 073006

¹ E-mail: joanna.hoszowska@unifr.ch

