



THE UNIVERSITY *of* EDINBURGH

Edinburgh Research Explorer

Aging and the mucosal immune system in the intestine

Citation for published version:

Mabbott, N, Kobayashi, A, Sehgal, A, Bradford, B, Pattison, M & Donaldson, D 2015, 'Aging and the mucosal immune system in the intestine' *Biogerontology*, vol. 16, no. 2 Sp. Iss. S1, pp. 133-145. DOI: 10.1007/s10522-014-9498-z

Digital Object Identifier (DOI):

[10.1007/s10522-014-9498-z](https://doi.org/10.1007/s10522-014-9498-z)

Link:

[Link to publication record in Edinburgh Research Explorer](#)

Document Version:

Peer reviewed version

Published In:

Biogerontology

Publisher Rights Statement:

This is a PDF file of an unedited manuscript that has been accepted for publication. The publisher version is available at: <http://link.springer.com/article/10.1007%2Fs10522-014-9498-z>

General rights

Copyright for the publications made accessible via the Edinburgh Research Explorer is retained by the author(s) and / or other copyright owners and it is a condition of accessing these publications that users recognise and abide by the legal requirements associated with these rights.

Take down policy

The University of Edinburgh has made every reasonable effort to ensure that Edinburgh Research Explorer content complies with UK legislation. If you believe that the public display of this file breaches copyright please contact openaccess@ed.ac.uk providing details, and we will remove access to the work immediately and investigate your claim.



Aging and the mucosal immune system in the intestine

Running title: Ageing and the mucosal immune system

Neil A. Mabbott^{1#}, Atsushi Kobayashi², Anuj Sehgal¹, Barry M. Bradford¹, Mari Pattison¹ & David S. Donaldson¹

¹The Roslin Institute & Royal (Dick) School of Veterinary Sciences, University of Edinburgh, United Kingdom

10

²Division of Neurological Science, Tohoku University Graduate School of Medicine
2-1 Seiryō-machi, Aoba-ku, Sendai 980-8575, JAPAN

#Correspondence: Dr. Neil A. Mabbott: The Roslin Institute & Royal (Dick) School of Veterinary Sciences, University of Edinburgh, Easter Bush, Midlothian EH25 9RG, United Kingdom

Tel: +44 (0)131 651 9100 Fax: +44 (0)131 651 9105

e-mail: neil.mabbott@roslin.ed.ac.uk

20

Keywords: Aging; intestine; mucosal immune system; M cells; IgA

The authors declare no competing financial conflict of interest.

Please cite this article as: Mabbott, N, Kobayashi, A, Sehgal, A, Bradford, B, Pattison, M & Donaldson, D, 'Aging and the mucosal immune system in the intestine' *Biogerontology* (2014), <http://dx.doi.org/10.1007/s10522-014-9498-z>

30

Abstract

Bacterial and viral infections of the gastrointestinal tract are more common in the elderly and represent a major cause of morbidity and mortality. The mucosal immune system provides the first line of defence against pathogens acquired by ingestion and inhalation, but its function is adversely affected in the elderly. This aging-related decline in the immune function is termed *immunosenescence* and is associated with diminished abilities to generate protective immunity, reduced vaccine efficacy, increased incidence of cancer, inflammation and autoimmunity, and the impaired ability to generate tolerance to harmless antigens. In this review we describe our current understanding of the effects immunosenescence has on the innate and adaptive arms of the mucosal immune system in the intestine. Current estimates suggest that by the year 2050 up to 40% of the UK population will be over 65 years old, bringing with it important health challenges. A thorough understanding of the mechanisms that contribute to the development of immunosenescence is therefore crucial to help identify novel approaches to improve mucosal immunity in the elderly.

50

Introduction

The mucosal immune system of the gastrointestinal tract provides the first line of defence against pathogens acquired by ingestion. In addition to pathogenic organisms, the gastrointestinal tract is continuously exposed to large amounts of commensal microorganisms and food proteins. The mucosal immune system must discriminate between these harmless antigens and generate tolerance towards them, whilst retaining the ability to generate protective immune responses against pathogens. Both responses are generated in the gut-associated lymphoid tissues (GALT) comprising chiefly of the appendix, tonsils, Peyer's patches, colonic patches, caecal patches and isolated lymphoid follicles (ILF). In addition, immune cells and lymph draining from the intestine accumulates in the mesenteric lymph nodes (MLN). The highly organized microarchitecture of the GALT enables antigen, antigen presenting cells and rare antigen-specific lymphocytes to interact and mount effective immune responses. These lymphoid structures, together with the diffusely distributed effector cells within the epithelium, the underlying lamina propria, protect the intestine against infection from pathogens and their toxins (Mowat 2003).

The mucosal immune response in the intestine is significantly compromised in the elderly (termed *immunosencecence*). This decline with age affects both innate and adaptive immunity and is associated with diminished antigen-specific antibody titres in the intestinal lumen and a decreased ability to generate tolerance to harmless antigens (Schmucker et al. 2003; Koga et al. 2000; Kato et al. 2003; Fujihashi and McGhee 2004; Dunn-Walters et al. 2003). This effect coincides with age-related increases in the incidence and severity of gastrointestinal infections, tumours and

inflammatory diseases, coupled with decreases in the efficacy of vaccinations (Fagiolo et al. 1993; Fujihashi et al. 2000; Fujihashi and McGhee 2004). Although many studies have addressed the age-related changes to systemic immune responses, particularly the effects of aging on T cell responses (which have been reviewed elsewhere (Akbar and Henson 2011; Henson and Akbar 2010)), much less
80 is comparatively known of the mechanisms underlying the decline in intestinal immune function. Age-related involution of Peyer's patches is frequently discussed in the literature (for example (St. Rose et al. 2006; Fujihashi and Kiyono 2009)) but these tissues persist throughout the lifespan of many species (Liebler-Tenorio and Pabst 2006; Zidan and Pabst 2008). Furthermore, the number of ILF, individual B-cell follicles found throughout the intestines, significantly increases in the intestines of aged mice (McDonald et al. 2011).

The UK elderly population continues to expand due to a combination of factors such as increasing life expectancy and reduced birth rates. At the time of writing data
90 suggested that ~10 million people in the UK were ≥ 65 years old, and this number is predicted to increase to ~15.5 million by 2030, and double to ~19 million by 2050 (Commons 2013). This significant demographic change brings with it important health challenges. Morbidity and mortality from gastrointestinal tract pathogens are substantially increased in the elderly, and the treatment of aging-related conditions absorbs the largest and growing share of the UK health care budget. A thorough understanding of the mechanisms that lead to mucosal immunosenescence is therefore crucial for the development of vaccines and therapeutic strategies to enhance mucosal immunity in the elderly.

100 The induction and maintenance of an antigen-specific mucosal immune response in the intestine is considered to be a multistep process occurring via the following key stages:

Step 1: Transport of gut luminal antigen across the intestinal epithelium

Step 2: Antigen presentation by mononuclear phagocytes to T cells within the GALT and MLN

Step 3: Polarization of T cells along effector or tolerance pathways

Step 4: Antibody isotype switching, differentiation and subsequent migration of antigen-specific IgA⁺ B cell immunoblasts to the intestinal lamina propria

110 **Step 5:** Local IgA production by plasma cells in the lamina propria

Step 6: Transport of IgA across the intestinal epithelium and secretion into the lumen

With the above steps in mind, this review summarizes our current understanding of how the function of the mucosal immune system in the intestine is affected by aging, particularly the effects on the innate immune system and IgA production by B cells.

Effects on the transport of lumen antigen across the intestinal epithelium

M cells: The lumen of the mammalian lower intestine contains an enormous burden of commensal microorganisms which is estimated to comprise $\sim 10^{12}$ bacteria/g of contents (Whitman et al. 1998). Access of these microorganisms to the underlying host tissues is restricted by a single layer of intestinal epithelial cells bound by tight-junctions. However, to initiate a mucosal immune response against potential pathogens, antigens must first be acquired from the gut lumen. The epithelium overlying the B-cell follicles of the GALT, termed the follicle-associated epithelium

120

(FAE), is specialized to enable the transportation of antigen across the intestinal epithelium into the GALT (a process termed, *transcytosis*). Under steady-state conditions ~10 % of the cells within FAE are M cells (Lorenz and Newberry 2004; Kanaya et al. 2012; Tahoun et al. 2012). Villi typically lack significant numbers of M cells, but studies suggest their density may increase in response to certain
130 inflammatory stimuli (Jang et al. 2004; Terahara et al. 2008). M cells in the FAE, in contrast to the neighbouring enterocytes, have a reduced glycocalyx, irregular brush border and reduced microvilli and are specialized for the transcytosis of particulate luminal antigen and microorganisms across the FAE (Krahenbuhl and Neutra 2000; Knoop et al. 2009; Nakato et al. 2009; Hase et al. 2009; Nakato et al. 2012; Sato et al. 2013; Kim et al. 2011). The transcytosed antigens exit into the basolateral pocket beneath the M-cell enabling the luminal antigens to be sampled and processed by lymphocytes and mononuclear phagocytes (MNP) (Lelouard et al. 2012; Wang et al. 2011). In the absence of M cells, antigen-specific T-cell responses in the Peyer's patches of mice orally-infected with *Salmonella enterica* serovar Typhimurium are
140 reduced (Hase et al. 2009; Kanaya et al. 2012). This indicates that antigen transcytosis by M cells is an important initial step in the induction of some mucosal immune responses (Lelouard et al. 2012; Wang et al. 2011; Hase et al. 2009; Kanaya et al. 2012). However, some pathogens have evolved to exploit M cells to invade host tissues from the gut lumen (Wolf et al. 1981; Amerongen et al. 1991; Fujimura et al. 2004; Takahashi et al. 2008; Westphal et al. 2008; Donaldson et al. 2012; Tahoun et al. 2012).

Using an immunosenescent mouse model (\geq 18 months old C57BL/6 mice) we have shown that there is a dramatic decline in the density of mature M cells in the Peyer's

150 patches of aged mice which impeded the transcytosis of particulate antigen across the FAE (Kobayashi et al. 2013) (**Fig. 1**). To gain an understanding of the mechanism underlying this aging-related decline, we also examined effects on the factors which control the induction and M-cell differentiation and maturation. The production of receptor activator of NF- κ B ligand (RANKL) by the network of stromal cells beneath the FAE stimulates the differentiation of receptor activator of NF- κ B (RANK)-expressing enterocytes into M cells (Knoop et al. 2009). The subsequent functional maturation of M cells is then mediated by their intrinsic expression of the ETS (E26 transformation-specific) family transcription factor Spi-B (Kanaya et al. 2012). Our data show that RANKL expression by the underlying stromal cells
160 beneath the FAE, or the initial induction of M cell differentiation, were unaffected in aged mice. Instead, specific impairments in the expression of Spi-B and the downstream functional development of immature M cells into functionally mature M cells were observed in aged mice.

In the GALT, the chemokine CCL20 is specifically expressed by the FAE where it mediates the chemoattraction of CCR6-expressing lymphocytes and leukocytes towards the intestinal epithelium. CCL20-CCR6-stimulation has also been suggested to influence M-cell maturation as their density is reduced in CCR6-deficient mice (Westphal et al. 2008; Ebisawa et al. 2011; Kobayashi et al. 2013).
170 The expression of CCL20 by the FAE was reduced in aged mice, consistent with data from an independent study on ILF in aged mice (McDonald et al. 2011). As a consequence of the reduced CCL20 expression, the chemoattraction of certain populations of MNP and lymphocytes, including those with apparent "M-cell-

inducing” potential (Ebisawa et al. 2011), towards the FAE were similarly reduced (Kobayashi et al. 2013).

180 These data suggest that the effects of aging on Spi-B and CCL20 expression appear to dramatically impede the functional maturation of M cells. Antigen-specific mucosal immune responses are markedly diminished in Spi-B-deficient (Kanaya et al. 2012) and CCR6-deficient mice (Cook et al. 2000). The density of mature M cells in the FAE of these mouse strains is also dramatically reduced (Kanaya et al. 2012; Ebisawa et al. 2011), suggesting that the effects of aging on M cell status may likewise contribute to the impaired Ag-specific mucosal immune responses observed in aged mice. These data also imply that the efficacy of M-cell-targeted vaccines may be significantly reduced in the elderly (Nochi et al. 2007; Misumi et al. 2009). However, despite this apparent deficiency, it is plausible that the effects of ageing on M cell maturation may reduce susceptibility to some pathogens that appear to exploit M cells to cross the gut epithelium and infect the host (Amerongen et al. 1991; Jones et al. 1994; Neutra et al. 1996; Sansonetti and Phalipon 1999; Donaldson et al. 190 2012). Susceptibility to oral prion infection is reduced in aged mice (Brown et al. 2009). Since M cells and Peyer’s patches are important sites of prion uptake from the gut lumen after oral exposure (Donaldson et al. 2012; Glaysher and Mabbott 2007), the reduced density of M cells in the FAE of aged mice (Kobayashi et al. 2013) may potentially impede their ability to initially cross the gut epithelium and infect the host.

Whether similar effects of aging on M cells are observed in ILF or the multi-follicular patch-like structures in the large intestine remains to be determined. However, the

demonstration that CCL20 expression by the FAE overlying ILF is reduced in aged mice (McDonald et al. 2011) suggests M-cell density may likewise be impeded aged ILF. A thorough analysis of the aging-related factors that underpin the dramatic decline in the functional maturation of M cells will help identify novel approaches to stimulate M cell differentiation and improve mucosal immunity in the elderly.

Goblet cells: Goblet cells are characteristically considered to act as mucin secreting cells in the gut epithelium. However, in the steady-state goblet cells may also provide passages for the delivery of low molecular weight soluble antigens to CD103⁺ MNP in the lamina propria (McDole et al. 2012). The precise immunological consequences of antigen sampling via these goblet-cell-associated antigen passages (GAPs) are uncertain, but the apparent preferential delivery of antigens to MNP with tolerogenic properties suggests an important role in regulating intestinal immune homeostasis. Goblet cell dysfunction has been associated with the development of intestinal inflammation (Dvorak and Dickersin 1980; van der Sluis et al. 2006), but whether aging influences their density, function and/or antigen sampling via GAPs is uncertain. Increased goblet cell density has been reported in the intestines of 12 month old rats (Aller et al. 2006), but no differences were observed in the FAE of aged mice (Kobayashi et al. 2013) (**Fig. 1**).

Epithelial cell barrier integrity: An intact intestinal epithelium provides an important protective barrier between the contents of the gut lumen and host tissues. Intercellular tight-junctions at the apical lateral membrane link neighbouring epithelial cells together and help determine paracellular permeability. Data from two independent studies suggest the expression of molecular components

of tight-junctions including zonula occludens-1 (ZO-1), junctional adhesion molecules (JAMs) and occludins is decreased in the intestinal epithelia of aged rats (Ren et al. 2013) and baboons (Tran and Greenwood-Van Meerveld 2013). These studies suggest that aging is associated with significant intestinal barrier dysfunction. The precise implications for mucosal immunity are uncertain. Disruption to tight junctions in the intestinal epithelium may lead to increased paracellular permeability to lumenal
230 antigens and/or proinflammatory stimuli such as bacterial endotoxins. Whether this apparent dysregulation in intestinal permeability contributes to the low-level chronic inflammation often observed in the elderly (termed *inflamm-aging*) remains to be determined (Franceschi et al. 2000).

Effects on MNP

MNP comprise a heterogeneous population of macrophages and classical dendritic cells (Mabbott et al. 2010; Bradford et al. 2011; Hume et al. 2013) and are abundant throughout the intestine, residing within the villous cores and in distinct anatomical locations within the GALT. At these sites MNP are strategically positioned to aid the
240 clearance of potential pathogens by phagocytosing and destroying them, or processing and presenting antigens on their surfaces to stimulate lymphocytes. For example, the sub-epithelium dome (SED) region of the GALT beneath the FAE (**Fig. 2**) contains a dense collection of MNP which readily acquires pathogens and antigens after transcytosis through M cells. Immunohistological data suggested CD11c⁺ MNP were reduced in the Peyer's patches of one year old mice (Kato et al. 2003), and a significant decline in the number of macrophages in the lamina propria was also noted in the intestines of aged dogs (Kleinschmidt et al. 2007). However, our own analysis of Peyer's patches from aged mice (~18 months old) using a highly

sensitive immunohistochemical method of detection does not concur with the Kato
250 study. Whereas the specific association of apparent M-cell-inducing
CD11c⁺CD45R⁺(B220) cells (Ebisawa et al. 2011) within the FAE was significantly
reduced in aged mice (coincident with the reduced expression of CCL20 (Kobayashi
et al. 2013)), the density of CD11c⁺ MNP in the SED was not adversely affected
(**Fig. 3**). Whether aging affects phagocytosis or antigen presentation by mucosal
MNP or their expression of MHC, co-stimulatory molecules or toll-like receptors,
influencing the nature of the subsequent mucosal immune response (inflammatory
vs. tolerogenic) is uncertain? One study has reported that CD11c⁺ MNP from the
MLN of 9 and 12 month old mice had a suboptimal ability to evoke an antigen-
specific T-cell response after oral infection with the intracellular parasite
260 *Encephalitozoon cuniculi* (Moretto et al. 2008). Interestingly, splenic CD11c⁺ MNP
from these animals displayed intact responses, indicating that aging may have
differing effects on mucosal and systemic immune responses.

Although the incidence is rare in the steady-state, certain inflammatory stimuli
appear to recruit MNP from the lamina propria to the gut epithelium where they insert
their dendrites through the tight junctions between enterocytes to directly sample the
luminal contents (Rescigno et al. 2001; Vallon-Eberhard et al. 2006; Niess et al.
2005; Farache et al. 2013). Other studies suggest that following challenge with non-
invasive *Salmonella*, CD11c⁺CX₃CR1⁺MHCII⁺CD11b⁻CD8 α ⁻ MNP may even migrate
270 into the intestinal lumen (Nicoletti et al. 2009; Argues et al. 2009). However, it is
unclear whether aging impedes the ability of MNP to directly sample the luminal
contents of the gut.

Conversely the effects of aging on other immune cell populations such as T cells may indirectly affect the functions of MNP in the gut. Alternatively activated macrophages, typified by their high expression of arginase-1, chitinase 3-like 3 (Ym1), restin-like- α (FIZZ-1) and the mannose receptor, are important effector cells in the clearance of intestinal parasites and the modulation of tissue repair (Jenkins and Allen 2010). A Th2-polarized immune response dominated by expression of high levels of IL-4, IL-10, IL-13 and IL-21 is considered important for the induction of alternatively activated macrophages. Classically activated macrophages, in contrast, are stimulated by the Th1 cytokine IFN- γ and characteristically express high levels of inducible nitric oxide synthase which acts to kill intracellular pathogens via the production of reactive nitrogen intermediates from L-arginine. The elevated background levels of the pro-inflammatory cytokines IFN- γ , TNF α and IL-6 in the intestines of naïve aged mice appear to restrain the induction of a strong Th2 response following intestinal helminth infection and the differentiation of alternatively activated macrophages (Sugawara et al. 2011).

290 **Effects on secretory IgA responses**

The production of antigen-specific secretory immunoglobulin A (IgA) dominates the humoral mucosal immune response and provides an important, high-affinity, first line of defence in the intestine by preventing specific pathogens such as *Salmonella typhimurium*, *Shigella flexneri*, reovirus and pathogen-derived toxins from crossing the intestinal epithelium (Michetti et al. 1992; Lycke et al. 1999; Silvey et al. 2001; Macpherson et al. 2008; Boullier et al. 2009; Mantis et al. 2011). In addition to effects on pathogenic microorganisms, secretory IgA also helps regulate the composition of the commensal gut microbiota (Mantis et al. 2011).

300 A diverse B-cell repertoire is important to ensure provision of a wide range of antibodies against potential pathogens. This diversity is maintained by rearrangements in the immunoglobulin gene which produces B cells with unique immunoglobulin genes and different antigen specificities. High affinity antibodies are produced by B cells as a consequence of affinity maturation which occurs within the germinal centres of lymphoid tissues. In the intestine, immunoglobulin isotype class-switching to IgA appears to occur only in the organized lymphoid structures of the GALT such as Peyer's patches and ILF (Shikina et al. 2004; Barone et al. 2011; Knoop and Newberry 2012) which contain all the necessary cellular components required to generate IgA-committed B cells including B cell follicles with germinal

310 centres, T cells and networks of follicular dendritic cells (FDCs). When a B-cell is activated by antigen it enters the germinal centre, proliferates and mutations are accumulated within the immunoglobulin gene (termed *hypermutation*). These mutations attempt to improve the antigen-binding affinity of the antibody. Thus, the newly encoded antibody is expressed on the B-cell surface and negative selection occurs whereby the cell undergoes apoptosis unless it receives rescue signals based on the antigen-affinity of the new antibody. If the B-cell receives the appropriate rescue signals, its immunoglobulin gene undergoes further rounds of mutation and selection, ultimately developing into an antibody secreting plasma cell or memory B cell.

320

There are conflicting reports in the literature over whether the overall quantity of IgA produced by the aged mucosal immune system is affected. Some data suggest aging has no effect on the amount of antibodies produced, whereas our own data

(alongside others) reveal a significant increase (**Fig. 4**) (Arranz et al. 1992; Senda et al. 1988; McDonald et al. 2011; Ebersole et al. 1985; Koga et al. 2000). However, antigen-specific IgA responses are adversely affected in the elderly and have a lower affinity (Taylor et al. 1992; Thoreux et al. 2000; Koga et al. 2000; Banerjee et al. 2002; Ademokun et al. 2010; Dunn-Walters et al. 2003). These contrasting effects suggest that the production of IgA becomes dysregulated as a consequence of aging

330 (McDonald et al. 2011). The hypermutation rate is similar in young and old B cells, but aged cells have increased accumulations of mutations (Linder et al. 2012; Dunn-Walters et al. 2003; Koga et al. 2000; Banerjee et al. 2002). Furthermore, within the Peyer's patches of elderly humans the process of Ig gene selection within the GC is also decreased (Banerjee et al. 2002). The stromal FDC within the B cell follicles play key roles in the induction of antibody hypermutation and maintenance of B cell memory (Kosco-Vilbois 2003; Haberman and Shlomchik 2003). In the spleens from aged mice FDC status is severely compromised (Aydar et al. 2004; Brown et al. 2009; Brown et al. 2012). This defect in FDC status has been shown to result in impaired immune-complex retention, germinal centre formation and antibody

340 production. Whether FDC in the GALT are similarly affected by aging is uncertain (Kato et al. 2003).

The IGHV-IGHD-IGHJ region of the immunoglobulin heavy chain gene is known as the CDR3 region and displays large diversity through recombination of different V, D and J segments. As a consequence, the diversity of the B-cell repertoire can be readily assessed by comparing immunoglobulin heavy chain CDR3 sequences and the level of hypermutations within them. In aged mice the intestinal IgA repertoire diversity has been shown to be increased (Linder et al. 2012) and skewed to one

more reflective of the systemic B-cell pool (McDonald et al. 2011). Some very old
350 humans, however, display a dramatically reduced diversity in their peripheral blood
B-cells which is associated with frailty (Gibson et al. 2009).

In addition to T-cell-dependent GC reactions, alternative modes of plasma cell
generation have been described including T cell-independent processes and
immunoglobulin isotype class-switching by peritoneal B-1 B cells in the lamina
propria (Cerruti et al. 2011; Suzuki et al. 2010). Further studies are required to
determine how aging specifically impacts on the induction of T-cell-dependent and T-
cell-independent IgA responses, but in aged mice and elderly humans current
evidence suggests most IgA-secreting plasma cells appear to be generated by T-cell
360 help (Barone et al. 2011; Gibbons and Spencer 2011; Linder et al. 2012).

Once high-affinity IgA-committed B cells are selected they then circulate through the
bloodstream and lymphatics to seed the lamina propria with plasma cell precursors
which synthesize dimeric IgA for secretion through the epithelium into the gut lumen
(Macpherson et al. 2008). The subsequent emigration of IgA⁺ B cell plasmablasts
from the Peyer's patches and their homing to the intestinal lamina propria has also
been reported to be reduced in aged rodents (Schmucker et al. 2003; Thoreux et al.
2000). Similarly, adoptive transfer experiments showed that the migration of aged
MLN lymphocytes to the lamina propria of young recipients was impaired (Thoreux et
370 al. 2000). Whether the decline in aged plasmablast emigration coincides with their
reduced expression of integrins is uncertain (Schmucker et al. 2003; Ogino et al.
2004). No differences in the expression of the polymeric immunoglobulin receptors
on the basolateral plasma membrane of enterocytes in the aged small intestine have

been observed, implying that the secretion of IgA into the gut lumen is not affected by aging (Taylor et al. 1992; Daniels et al. 1998).

In addition to the secretion of IgA into the gut lumen to facilitate immune-exclusion, the transcytosis of secretory IgA-containing immune-complexes back across the epithelium by M cells may also help modulate the mucosal immune response (Rey et al. 2004; Kadaoui and Corthesy 2007; Rochereau et al. 2013). The transcytosed IgA-containing immune-complexes appear to be acquired by the underlying MNP and, as a consequence, are unable to disseminate beyond the MLNs. This selective delivery of potential gut pathogens to MNP by IgA may enable antigen recognition under neutralizing and non-inflammatory conditions without causing widespread tissue dissemination. The reduced functional maturation of M cells in the aged intestine suggests that the transcytosis of IgA-containing immune complexes would be impaired by aging (Kobayashi et al. 2013). Whether there is increased retention of IgA in the aged gut lumen (**Fig. 4**)(McDonald et al. 2011), for example as a consequence of the effects of aging on M-cell function, remains to be determined.

390

Effects on the gut microbiota

The mammalian intestine is host to a huge number and variety of indigenous bacteria. The composition of this microbiota has a profound effect on the host. In addition to providing nutritional benefit (vitamin synthesis, metabolism of dietary components etc.) these commensals help stimulate the maturation of the immune system and the effective generation of adaptive immune responses (Macpherson and Harris 2004; Bouskra et al. 2008). Disturbances to this balance have been implicated in the development of intestinal pathology. Therefore, much attention has

400 been focussed on understanding how aging-related effects on the intestinal microbiota may influence host immunity. Although a polyclonal IgA repertoire can develop in the absence of the gut microbiota, the CDR3 sequence repertoire of germ-free mice contains fewer hypermutations, and their intestines harbour fewer plasma cells (Linder et al. 2012). The exterior and interior surfaces of the FAE of Peyer's patches are colonized by distinct bacterial species which can also influence the development and maturation of the mucosal immune system. Segmented filamentous bacteria are potent activators of the mucosal immune system and preferentially attach to the exterior surface of the FAE, whereas *Alcaligenes* inhabits the interior of Peyer's patches (Obata et al. 2010). Whether the distribution of these bacteria in the GALT is significantly affected by aging is uncertain, but the gut
410 microbiota composition of young individuals has been shown to differ significantly from that of the elderly (Biagi et al. 2010; Claesson et al. 2012).

Following weaning the human gut microbiota is considered to develop an "adult-like" composition at around two years of age (Favier et al. 2002), remaining relatively stable throughout adult life. While acute variations may occur, for example as a consequence of gastrointestinal infection, antibiotic treatment or dietary changes (David et al. 2013), this stability is dramatically reduced in old age. Data have shown that bifidobacteria significantly decrease in the elderly, whereas lactobacilli, coliforms, enterococci and *Clostridium perfringens* are increased (Hopkins et al.
420 2001; Mueller et al. 2006; Tiihonen et al. 2008). The precise aging-related mechanisms behind these changes to the gut microbiota are uncertain, but it is clear that they correlate significantly with effects on health status. For example, in one study suggested the microbiota of centenarians was enriched in facultative

anaerobes which may contribute towards the increased inflammatory status in these individuals (Biagi et al. 2010). The microbial fermentation product butyrate has been shown to induce the differentiation of colonic regulatory T cells and ameliorate experimental colitis (Maslowski et al. 2009; Furusawa et al. 2013; Smith et al. 2013). Increased levels of systemic IL-6 and IL-8 in the elderly, coincided with an enrichment of Proteobacteria and a decrease in the amount of some butyrate
430 producing bacteria (Biagi et al. 2010). Cross-talk between the host immune system and the microbiota influences the development of the mucosal immune system, which in turn regulates the microbiota (Round and Mazmanian 2009). Thus, further studies are necessary to determine whether these age-related differences in the composition of the gut microbiota are the cause of the increased inflammation in the elderly, or a consequence of the systemic inflammatory status. A detailed understanding of the mechanisms underlying host-intestinal microbiota cross-talk will help determine whether nutritional manipulation of the gut microbiota in the elderly may help preserve mucosal immunity (Schiffrin et al. 2009).

440 **Concluding remarks**

The aging-associated increase in the incidence of inflammation, autoimmunity, cancer and susceptibility to gastrointestinal infections coincides with a significant decline in host immunity and the efficacy of vaccinations. As discussed above, this immunosenescence has multiple effects both within the innate and adaptive arms of the mucosal immune system in the intestine. Despite significant progress in our understanding of the effects of aging on the mucosal immune system in the intestine, significant gaps remain. A thorough analysis of the cellular and molecular mechanisms that underpin the dramatic aging-related decline in immune

homeostasis will help identify the factors that enhance susceptibility to orally-
450 acquired pathogens and reduce the efficacy of mucosal vaccinations, and identify
novel approaches to improve mucosal immunity in the elderly.

Acknowledgements

This work was supported by project (BB/J014672/1, BB/K021257/1 & BB/L007452/1)
and Institute Strategic Programme Grant funding from the Biotechnology and
Biological Sciences Research Council.

References

- 460 Ademokun A, Wu Y-C, Martin V, Mitra R, Sack U, Baxendale H, Kipling D, Dunn-Walters DK (2010) Vaccination-induced changes in human B-cell repertoire and pneumococcal IgM and IgA antibody at different ages. *Aging Cell* 10:922-930
- Akbar AN, Henson SM (2011) Are senescence and exhaustion intertwined or unrelated processes that comprise immunity? *Nature Rev Immunol* 11:289-295
- Aller MA, Corcuera MT, Gomez-Aguado F, Sanchez-Patan F, Alonso MJ, Arias J (2006) Age-related changes in goblet cell density in the rat small bowel. Comment on "Age-related changes in liver structure and function: Implications for disease?" by DL Schmucker. *Exp Gerontol* 41:564
- 470 Amerongen HM, Weltzin R, Farnet CM, Michetti P, Haseltine WA, Neutra MR (1991) Transepithelial transport of HIV-1 by intestinal M cells: a mechanism for transmission of AIDS. *J Acquir Immune Defic Syndr* 4:760-765
- Argues JL, Hautefort I, Ivory K, Bertelli E, Regoli M, Clare S, Hinton JC, Nicoletti C (2009) Salmonella induces flagellin- and MyD88-dependent migration of bacteria-capturing dendritic cells into the gut lumen. *Gastroenterology* 137:579-587
- Arranz E, O'Mahony S, Barton JR, Ferguson A (1992) Immunosenescence and mucosal immunity: significant effects of old age on secretory IgA
- 480 Aydar Y, Balogh P, Tew JG, Szakal AK (2004) Follicular dendritic cells in aging, a 'bottle-neck' in the humoral response. *Ageing Research Reviews* 3:15-29
- Banerjee M, Mehr R, Belevsky A, Spencer J, Dunn-Walters DK (2002) Age- and tissue-specific differences in human germinal center B cell selection revealed by analysis of IgV_H gene hypermutation and lineage trees. *Eur J Immunol* 32:1947-1957
- Barone F, Vossenkamper A, Boursier L, Su W, Watson A, John S, Dunn-Walters DK, Fields P, Wijetilleka S, Edgeworth JD, Spencer J (2011) IgA-producing plasma cells originate from germinal centres that are induced by B-cell
- 490 Biagi E, Nylund L, Candela M, Ostan R, Bucci L, Pini E, Nikkila J, Monti D, Satokari R, Franceschi C, Brigidi P, De Vos W (2010) Through ageing, and beyond: gut microbiota and inflammatory status in seniors and centenarians. *PLoS One* 5:e10667
- Boullier S, Tanguy M, Kadaoui KA, Caubet C, Sansonetti P, Corthesy B, Phalipon A (2009) Secretory IgA-mediated neutralization of *Shigella flexneri* prevents intestinal tissue destruction by down-regulating inflammatory circuits. *J Immunol* 183:5879-5885
- Bouskra D, Brezillon C, Berard M, Werts C, Varona C, Boneca IG, Eberl G (2008)
- 500 Lymphoid tissue genesis induced by commensals through NOD1 regulates intestinal homeostasis. *Nature* 456:507-510
- Bradford BM, Sester D, Hume DA, Mabbott NA (2011) Defining the anatomical localisation of subsets of the murine mononuclear phagocyte system using integrin alpha X (ITGAX) and colony stimulating factor 1 receptor (CSF1-R) expression fails to discriminate dendritic cells from macrophages. *Immunobiology* 216:1228-1237

- 510 Brown KL, Gossner A, Mok S, Mabbott NA (2012) The effects of host age on the transport of complement-bound complexes to the spleen and the pathogenesis of intravenous scrapie infection. *J Virol* 86:1228-1237. doi:10.1128/JVI.05581-11
- Brown KL, Wathne GJ, Sales J, Bruce ME, Mabbott NA (2009) The effects of host age on follicular dendritic cell status dramatically impair scrapie agent neuroinvasion in aged mice. *J Immunol* 183:5199-5207. doi:10.4049/jimmunol.0802695
- Cerruti A, Chen K, Chorny A (2011) Immunoglobulin responses at the mucosal interface. *Annu Rev Immunol* 29:273-293
- 520 Claesson MJ, Jeffrey IB, Conde S, Power SE, O'Connor EM, Cusack S, Harris HM, Coakley M, Lakshminarayanan B, O'Sullivan O, Fitzgerald GF, Deane J, O'Connor M, Harnedy N, O'Mahony D, van Sinderen D, Wallace M, Brennan L, Stanton C, Marchesi JR, Fitzgerald AP, Shanahan F, Hill C, Ross RP, O'Toole PW (2012) Gut microbiota composition correlates with diet and health in the elderly. *Nature* 488:178-184
- Commons Ho (2013) The ageing population. <http://www.parliament.uk/business/publications/research/key-issues-for-the-new-parliament/value-for-money-in-public-services/the-ageing-population/>
- 530 Cook DN, Prosser DM, Forster R, Zhang J, Kuklin NA, Abbondanzo SJ, Niu X-D, Chen S-C, Manfra DJ, Wiekowski MT, Sullivan LM, Smith SR, Greenberg HB, Narula SK, Lipp M, Lira SA (2000) CCR6 mediates dendritic cell localization, lymphocyte homeostasis, and immune responses in mucosal tissue. *Immunity* 12:495-503
- Daniels CK, Schmucker D, Bazin H, Jones AL (1998) Immunoglobulin A receptor of rat small intestinal enterocytes is unaffected by ageing. *Gastroenterology* 94:1432-1440
- David LA, Maurice CF, Carmody RN, Gootenberg DB, Button JE, Wolfe BE, Ling AV, Devlin S, Varma Y, Fischbach MA, Biddinger SB, Dutton RJ, Turnbaugh PJ (2013) Diet rapidly and reproducibly alters the human gut microbiome. *Nature* in press
- 540 Donaldson DS, Kobayashi A, Ohno H, Yagita H, Williams IR, Mabbott NA (2012) M cell depletion blocks oral prion disease pathogenesis. *Mucosal Immunol* 5:216-225
- Dunn-Walters DK, Banerjee M, Mehr R (2003) Effects of age in antibody affinity maturation. *Biochemical Society Transactions* 31:447-448
- Dvorak AM, Dickersin GR (1980) Crohn's disease: transmission electron microscopic studies. I. Barrier function. Possible changes related to alterations of cell coat, mucous coat, epithelial cells, and Paneth cells. *Hum Pathol* 11:561-571
- Ebersole JL, Smith DJ, Taubman MA (1985) Secretory immune responses in ageing rats. I. Immunoglobulin levels. *Immunology* 56:345-350
- 550 Ebisawa M, Hase K, Takahashi D, Kitamura D, Knoop KA, Williams IR, Ohno H (2011) CCR6hiCD11cint B cells promote M-cell differentiation in Peyer's patch. *Int Immunol* 23:261-269
- Fagiolo U, Amadori A, Cozzi E, Bendo R, Lama M, Douglas A, Palu G (1993) Humoral and cellular immune responses to influenza virus vaccination in aged humans. *Aging* 5:451-458
- Farache J, Koren I, Milo I, Gurevich I, Kim K-W, Zigmund E, Furtado GC, Lira SA, Shakhar G (2013) Luminal bacteria recruit CD103+ dendritic cells into the

- intestinal epithelium to sample bacterial antigens for presentation. *Immunity* 38:581-595
- 560 Favier CF, Vaughan EE, De Vos WM, Akkermans AD (2002) Molecular monitoring of succession of bacterial communities in human neonates. *Appl Environ Microbiol* 68:291-226
- Franceschi C, Bonafe M, Valensin S, Olivieri F, De Luca M, Ottaviani E, De Benedictis G (2000) Inflamm-aging. An evolutionary perspective on immunosenescence. *Ann N Y Acad Sci* 908:244-254
- Fujihashi K, Kiyono H (2009) Mucosal immunosenescence: new developments and vaccines to control infectious diseases. *Trends Immunol* 30:334:343
- Fujihashi K, Koga T, McGhee JR (2000) Mucosal vaccination and immune responses in the elderly. *Vaccine* 18:1675-1680
- 570 Fujihashi K, McGhee JR (2004) Mucosal immunity and tolerance in the elderly. *Mech Ageing Dev* 125:889-898
- Fujimura Y, Takeda M, Ikai H, Haruma K, Akisada T, Harada T, Sakai T, Ohuchi M (2004) The role of M cells of human nasopharyngeal lymphoid tissue in influenza virus sampling. *Virchows Arch* 444:36-42
- Furusawa Y, Obata Y, Fukuda S, Endo TA, Nakato G, Takahashi D, Nakanishi Y, Uetake c, Kato K, Kato T, Takahashi M, Fukuda NN, Murukami S, Miyauchi E, Hino S, Atarashi K, Onawa S, Fujimura Y, Lockett T, Clarke JM, Topping DL, Tomita M, Hori S, Ohara O, Morita T, Koseki H, Kikuchi J, Honda K, Hase K, Ohno H (2013) Commensal microbe-derived butyrate induces the differentiation of colonic regulatory T cells. *Nature* 504:446-450
- 580 Gibbons DL, Spencer J (2011) Mouse and human intestinal immunity: Same ballpark, different players; different rules, same score. *Mucosal Immunol* 4:148-157
- Gibson KL, Wu Y-C, Barnett Y, Duggan O, Vaughan R, Kondeatis E, Nilsson B-O, Wikby A, Kipling D, Dunn-Walters DK (2009) B-cell diversity decreases in old age and is correlated with poor health status. *Aging Cell* 8:18-25
- Glaysner BR, Mabbott NA (2007) Role of the GALT in scrapie agent neuroinvasion from the intestine. *J Immunol* 178:3757-3766
- Haberman AM, Shlomchik MJ (2003) Reassessing the function of immune-complex retention by follicular dendritic cells. *Nature Rev Immunol* 3:757-764
- 590 Hase K, Kawano K, Nochi T, Pontes GS, Fukuda S, Ebisawa M, Kadokura K, Tobe T, Fujimura Y, Kawano S, Yabashi A, Waguri S, Nakato G, Kimura S, Murakami T, Imura M, Hamura K, Fukuoka S-I, Lowe AW, Itoh K, Kiyono H, Ohno H (2009) Uptake through glycoprotein 2 of FimH⁺ bacteria by M cells initiates mucosal immune responses. *Nature* 462:226-231
- Henson SM, Akbar AN (2010) Memory T-cell homeostasis and senescence during aging. *Adv Exp Med Biol* 684:189-197
- Hopkins MJ, Sharp R, Macfarlane GT (2001) Age and disease related changes in intestinal bacterial populations assessed by cell culture, 16S rRNA abundance, and community cellular fatty acid profiles. *Gut* 48:198-205
- 600 Hume DA, Mabbott N, Raza S, Freeman TC (2013) Can DCs be distinguished from macrophages by molecular signatures. *Nat Immunol* 14:187-189
- Jang MH, Kweon MN, Iwatani K, Yamamoto M, Terahara K, Sasakawa C, Suzuki T, Nochi T, Yokota Y, Rennert PD, Hiroi T, Tamagawa H, Iijima H, Kunisawa J, Yuki Y, Kiyono H (2004) Intestinal villous M cells: An antigen entry site in the mucosal epithelium. *Proc Natl Acad Sci USA* 101:6110-6115

- Jenkins SJ, Allen JE (2010) Similarity and diversity in macrophage activation by nematodes, trematodes, and cestodes. *J Biomed Biotechnol* 2010:262609
- Jones B, D., Ghorri N, Falkow S (1994) Salmonella typhimurium initiates murine infection by penetrating and destroying the specialized epithelial M cells of the Peyer's patches. *J Exp Med* 180:15-23
- 610 Kadaoui KA, Corthesy B (2007) Secretory IgA mediates bacterial translocation to dendritic cells in mouse Peyer's patches with restriction to mucosal compartment. *J Immunol* 179:7751-7757.
- Kanaya T, Hase K, Takahashi D, Fukuda S, Hoshino K, Sasaki I, Hemmi H, Knoop KA, Kumar N, Sato M, Katsuno T, Yokosuka O, Toyooka K, Nakai K, Sakamoto A, Kitahara Y, Jinnohara T, McSorley SJ, Kaisho T, Williams IR, Ohno H (2012) The Ets transcription factor Spi-B is essential for the differentiation of intestinal microfold cells. *Nat Immunol* 13:729-736
- Kato H, Fujihashi K, Kato R, Dohi T, Fujihashi K, Hagiwara Y, Kataoka K, Kobayashi R, McGhee JR (2003) Lack of oral tolerance in ageing is due to sequential loss of Peyer's patch cell interactions. *Int Immunol* 15:145-158
- 620 Kim SH, Jung DI, Yang IY, Kim J, Lee KY, Nochi T, Kiyono H, Jang YS (2011) M cells expressing the complement C5a receptor are efficient targets for mucosal vaccine delivery. *Eur J Immunol* 41:3219-3229
- Kleinschmidt S, Meneses F, Nolte I, Hewicker-Trautwein M (2007) Distribution of mast cell subtypes and immune cell populations in canine intestines: Evidence for age-related decline in T cells and macrophages and increase of IgA-positive plasma cells. *Res Vet Sci* 84:41-48
- Knoop KA, Kumar N, Butler BR, Sakthivel SK, Taylor RT, Nochi T, Akiba H, Yagita H, Kiyono H, Williams IR (2009) RANKL is necessary and sufficient to initiate development of antigen-sampling M cells in the intestinal epithelium. *J Immunol* 183:5738-5747. doi:10.4049/j.immunol.0901563
- 630 Knoop KA, Newberry RD (2012) Isolated lymphoid follicles are dynamic reservoirs for the induction of intestinal IgA. *Frontiers Immunol* 3:84
- Kobayashi A, Donaldson DS, Erridge C, Kanaya T, Williams IR, Ohno H, Mahajan A, Mabbott NA (2013) The functional maturation of M cells is dramatically reduced in the Peyer's patches of aged mice. *Mucosal Immunol* 6:1027-1037
- Koga T, McGhee JR, Kato H, Kato R, Kiyono H, Fujihashi K (2000) Evidence for early aging in the mucosal immune system. *J Immunol* 165:5352-5359
- 640 Kosco-Vilbois MH (2003) Are follicular dendritic cells really good for nothing? *Nature Rev Immunol* 3:764-769
- Krahenbuhl JP, Neutra MR (2000) Epithelial M cells: differentiation and function. *Annu Rev Cell Dev Biol* 16:301-332
- Lelouard H, Fallet M, De Bovis B, Meresse S, Gorvel JP (2012) Peyer's patch dendritic cells sample antigens by extending dendrites through M cell-specific transcellular pores. *Gastroenterology* 142:592-601
- Liebler-Tenorio EM, Pabst P (2006) MALT structure and function in farm animals. *Vet Res* 37:257-280
- Linder C, Wahl B, Fohse L, Suerbaum S, Macpherson AJ, Prinz I, Pabst O (2012) Age, microbiota, and T cells shape diverse individual IgA repertoires in the intestine. *J Exp Med* 209:365-377
- 650 Lorenz RG, Newberry RD (2004) Isolated lymphoid follicles can function as sites for induction of mucosal immune responses. *Ann N Y Acad Sci* 1029:44-57

- Lycke N, Erlandsson L, Ekman L, Schon K, Leanderson T (1999) Lack of J chain inhibits the transport of gut IgA and abrogates the development of intestinal antitoxic protection. *J Immunol* 163:913-919
- Mabbott NA, Baillie JC, Hume DA, Freeman TC (2010) Meta-analysis of co-expressed gene signatures in mouse leukocyte populations. *Immunobiology* 215:724-736
- 660 Macpherson AJ, Harris NL (2004) Interactions between commensal intestinal bacteria and the immune system. *Nature Rev Immunol* 4:478-485
- Macpherson AJ, McCoy KD, Johansen F-E, Brandtzaeg P (2008) The immune geography of IgA induction and function. *Mucosal Immunol* 1:11-22
- Mantis NJ, Rol N, Corthesy B (2011) Secretory IgA's complex roles in immunity and mucosal homeostasis in the gut. *Mucosal Immunol* 4:603-611
- Maslowski KM, Vieira AT, Ng A, Kranich J, Sierro F, Yu D, Schilter HC, Rolph MS, Mackay F, Artis D, Xavier RJ, Teixeira MM, Mackay F (2009) Regulation of inflammatory responses by gut microbiota and chemoattractant receptor GPR43. *Nature* 461:1282-1286
- 670 McDole JR, Wheeler LW, McDonald KG, Wang B, Konjufca V, Knoop KA, Newberry RD, Miller MJ (2012) Goblet cells deliver luminal antigen to CD103⁺ dendritic cells in the small intestine. *Nature* 483:345-349
- McDonald KG, Leach MR, Huang C, Wang C, Newberry RD (2011) Aging impacts isolated lymphoid follicle development and function. *Immun Aging* 8:1
- Michetti P, Mahan MJ, Slauch JM, Mekalanos JJ, Neutra MR (1992) Monoclonal secretory immunoglobulin A protects mice against oral challenge with the invasive pathogen *Salmonella typhimurium*. *Infect Immun*, 60:1786-1792
- Misumi S, Masuyama M, Takamune N, Nakayama D, Mitsumata R, Matsumoto H, Urata N, Takahashi Y, Muneoka A, Sukamoto T, Fukuzaki K, Shoji S (2009)
- 680 Targeted delivery of immunogen to primate m cells with tetragalloyl lysine dendrimer. *J Immunol* 182:6061-6070
- Moretto MM, Lawlor EM, Khan IA (2008) Aging mice exhibit a functional defect in mucosal dendritic cell response against an intracellular pathogen. *J Immunol* 181:7977-7984
- Mowat AM (2003) Anatomical basis of tolerance and immunity to intestinal pathogens. *Nature Rev Immunol* 3:331-341
- Mueller S, Saunier K, Hanisch C, Norin E, Alm L, Midtvedt T, Cresci A, Silvi S, Orpianesi C, Verdenelli MC, Clavel T, Koebnick C, Zunft HJ, Dore J, Blaut M (2006) Differences in fecal microbiota in different European study populations
- 690 in relation to age, gender, and country: a cross-sectional study. *Appl Environ Microbiol* 72:1027-1033
- Nakato G, Fukuda S, Hase K, Goitsuka R, Cooper MD, Ohno H (2009) New approach for M-cell-specific molecules by screening comprehensive transcriptome analysis. *DNA Res* 16:227-235
- Nakato G, Hase K, Suzuki M, Kimura M, Ato M, Hanazato M, Tobiume M, Horiuchi M, Atarashi R, Nishida N, Watarai H, Imaoka K, Ohno H (2012) Cutting edge: *Brucella abortus* exploits a cellular prion protein on intestinal M cells as an invasive receptor. *J Immunol* 189:1540-1544
- Neutra MR, Frey A, Kraehenbuhl J-P (1996) Epithelial M Cells: Gateways for
- 700 Mucosal Infection and Immunization. *Cell* 86:345-348
- Nicoletti C, Regoli M, Bertelli E (2009) Dendritic cells in the gut: to sample and to exclude? *Mucosal Immunol* 2:462

- Niess JH, Brand S, Gu X, Landsman L, Jung S, McCormick BA, Vyas JM, Boes M, Ploegh HL, Fox JG, al. e (2005) CX₃CR1-mediated dendritic cell access to the intestinal lumen and bacterial clearance. *Science* 307:254-258
- Nochi T, Yuki Y, Matsumura A, Mejima M, Terahara K, Kim D-Y, Fukuyama S, Iwatsuki-Horimoto K, Kawaoka Y, Kohda T, Kozaki S, Igrashi O, Kiyono H (2007) A novel M cell-specific carbohydrate-targeted mucosal vaccine effectively induces antigen-specific immune responses. *J Exp Med* 204:2789-2796
- 710 Obata T, Goto Y, Kunisawa Y, Sato S, Sakamoto M, Setoyama H, Matsuki T, Nonaka K, Shibata N, Gohda M, Kagiya Y, Nochi T, Yuki Y, Fukuyama Y, Mukai A, Shinzaki S, Fujihashi K, Sasakawa C, Iijima H, Goto M, Umesaki Y, Benno Y, Kiyono H (2010) Indigenous opportunistic bacteria inhabit mammalian gut-associated lymphoid tissues and share a mucosal antibody-mediated symbiosis. *Proc Natl Acad Sci USA* 107:7419-7424
- Ogino T, Miura S, Komoto S, Hara Y, Hokari R, Tsuzuki Y, Watanabe C, Koseki S, Nagata H, Hachimura S, Kaminogawa S, Ishii H (2004) Senescence-associated decline of lymphocyte migration in gut-associated lymphoid tissues of rat small intestine. *Mech Ageing Dev* 125
- 720 Ren W, Wu K, Li X, Luo M, Liu H, Zhang S, Hu Y (2013) Age-related changes in small intestinal mucosa epithelium architecture and epithelial tight junction in rat models. *Aging Clin Exp Res* in press
- Rescigno M, Urbano M, Valzasina B, Francolini M, Rotta G, Bonasio R, Granucci F, Kraehenbuhl J-P, Ricciardi-Castagnoli P (2001) Dendritic cells express tight junction proteins and penetrate gut epithelial monolayers to sample bacteria. *Nat Immunol* 2:361-367
- Rey J, Garin N, Spertini F, Corthesy B (2004) Targeting of secretory IgA to Peyer's patch dendritic cells and T cells after transport by intestinal M cells. *J Immunol* 172:3026-3033
- 730 Rochereau N, Drocourt D, Perouzel E, Pavot V, Redelinguys P, Brown GD, Tiraby G, Roblin X, Verrier B, Genin C, Corthesy B, Paul S (2013) Dectin-1 is essential for reverse transcytosis of glycosylated SIgA-antigen complexes by intestinal M cells. *PLoS Biol* 11:e1001658
- Round JL, Mazmanian SK (2009) The gut microbiota shapes intestinal immune responses during health and disease. *Nature Rev Immunol* 9:313-323
- Sansonetti PJ, Phalipon A (1999) M cells as ports of entry for enteroinvasive pathogens: mechanisms of interaction, consequences for the disease process. *Seminars Immunol* 11:193-203
- 740 Sasmono RT, Oceandy D, Pollard JW, Tong W, Pavli P, Wainwright BJ, Ostrowski MC, Himes SR, Hume DA (2003) A macrophage colony-stimulating factor receptor-green fluorescent protein transgene is expressed throughout the mononuclear phagocyte system of the mouse. *Blood* 101:1155-1163
- Sato S, Kaneto S, Shibata N, Takahashi Y, Okura H, Yuki Y, Kunisawa J, Kiyono H (2013) Transcription factor Spi-B-dependent and -independent pathways for the development of Peyer's patch M cells. *Mucosal Immunol* 6:838-846
- Schiffrin EJ, Parlesak A, Bode C, Bode JC, van't Hof MA, Grathwohl D, Guigoz Y (2009) Probiotic yoghurt in the elderly with intestinal bacterial overgrowth: endotoxaemia and innate immune functions. *Br J Nutr* 101:961-966
- 750 Schmucker D, Owen R, Outenreath R, Thoreux K (2003) Basis for the age-related decline in intestinal mucosal immunity. *Clinical and Developmental Immunology* 10:167-172

- Senda S, Cheng E, Kawanishi H (1988) Aging-associated changes in murine intestinal immunoglobulin A and M secretions. *Scand J Immunol* 27:157-164
- Shikina T, Hiroi T, Iwatani K, Jang MH, Fukuyama S, Tamura M, Kubo T, Ishikawa H, Kiyono H (2004) IgA class switch occurs in the organized nasopharynx- and gut-associated lymphoid tissue, but not in the diffuse lamina propria or airways and gut. *J Immunol* 172:6259-6264
- 760 Silvey KJ, Hutchings AB, Vajdy M, Petzke MM, Neutra MR (2001) Role of immunoglobulin A in protection against reovirus entry into murine Peyer's patches. *J Virol* 75:10870-10879
- Smith PM, Howitt MR, Panikov N, Michaud M, Gallini CA, Bohlooly-Y M, Glickman JN, Garrett WS (2013) The microbial metabolites, short-chain fatty acids, regulate colonic Treg cell homeostasis. *Science* 341:569-573
- St. Rose S, Hunter N, Matthews D, Foster J, Chase-Topping ME, Kruuk LEB, Shaw DJ, Rhind SM, Will RG, Woolhouse MEJ (2006) Comparative evidence for a link between Peyer's patch development and susceptibility to transmissible spongiform encephalopathies. *BMC Infectious Diseases* 6:5
- 770 Sugawara Y, Azuma N, Onodera S, Tsunoka Y, Morimoto M (2011) Th2 immune responses and alternatively activated macrophages (AAMacs) in helminth infection in aged mice. *J Vet Med Sci* 73:511-516
- Suzuki K, Maruya M, Kawamoto S, Fagarasan S (2010) Roles of B-1 and B-2 cells in innate and acquired IgA-mediated immunity. *Immunol Rev* 237:180-190
- Tahoun A, Mahajan S, Paxton E, Malterer G, Donaldson DS, Wang D, Tan A, Gillespie TL, O'Shea M, Rose A, Shaw DJ, Gally DL, Lengeling A, Mabbott NA, Haas J, Mahajan A (2012) Salmonella transforms follicle-associated epithelial cells into M cells to promote intestinal invasion. *Cell Host & Microbe* 12:645-666
- 780 Takahashi Y, Misumi S, Muneoka A, Masuyama M, Tokado H, Fukuzaki K, Takamune N, Shoji S (2008) Nonhuman primate intestinal villous M-like cells: an effective poliovirus entry site. *Biochem Biophys Res Commun* 368:501-507
- Taylor DL, Daniels CK, Schmucker D (1992) Ageing compromises gastrointestinal mucosal immune response in the rhesus monkey. *Immunology* 75:614-618
- Terahara K, Yoshida M, Igarashi O, Nochi T, Soares Pontes G, Hase K, Ohno H, Kurokawa S, Mejima M, Takayama N, Yuki Y, Lowe AW, Kiyono H (2008) Comprehensive gene expression profiling of Peyer's patch M cells, villous M-like cells, and intestinal epithelial cells. *J Immunol* 180:7840-7846
- Thoreux K, Owen RL, Schmucker D (2000) Intestinal lymphocyte number, migration and antibody secretion in young and old rats. *Immunology* 101:161-167
- 790 Tiihonen K, Tynkkynen S, Ouwehand A, Ahlroos T, Rautonen N (2008) The effect of ageing with and without non-steroidal anti-inflammatory drugs on gastrointestinal microbiology and immunology. *Br J Nutr* 100:130-137
- Tran L, Greenwood-Van Meerveld B (2013) Age-associated remodeling of the intestinal epithelial barrier. *J Gerontol A Biol Sci Med Sci* 68:1045-1056
- Vallon-Eberhard A, Landsman L, Yogev N, Verrier B, Jung S (2006) Transepithelial pathogen uptake into the small intestinal lamina propria. *J Immunol* 176:2465-2469
- 800 van der Sluis M, de Koning BA, De Bruijn AC, Velcich A, Meijerink JP, Van Goudoever JB, Buller HA, Dekker J, Van Seuningen I, Renes IB, Einerhand AW (2006) Muc2-deficient mice spontaneously develop colitis, indicating that MUC2 is critical for colonic protection. *Gastroenterology* 131:117-129

- Wang J, Gusti V, Saraswati A, Lo DD (2011) Convergent and divergent development among M cell lineages in mouse mucosal epithelium. *J Immunol* 187:5277-5285
- Westphal S, Lugering A, von Wedel J, von Eiff C, Maaser C, Spahn T, Heusipp G, Schmidt MA, Herbst H, Williams IR, Domschke W, Kucharzik T (2008) Resistance of chemokine receptor 6-deficient mice to *Yersinia enterocolitica* infection: evidence on defective M-cell formation *in vivo*. *Am J Pathol* 172:671-680
- 810 Whitman WB, Coleman DC, Wiebe WJ (1998) Prokaryotes: the unseen majority. *Proc Natl Acad Sci USA* 95:6578-6583
- Wolf JL, Rubin DH, Finberg R, Kauffman RS, Sharpe AH, Trier JS, Fields BN (1981) Intestinal M cells: a pathway for entry of reovirus into the host. *Science* 212:471-472
- Zidan M, Pabst R (2008) Unique microanatomy of ileal Peyer's patches of the one humped camel (*Cameuls dromodarius*) is not age dependent. *Anat Rec* 291:1023-1028

820

Figure 1 M-cell density is significantly reduced in the FAE of aged Peyer's

patches. (a) Z-stack image from whole-mount image of a Peyer's patch from a young mouse showing an M-cell with characteristic basolateral pocket (GP2⁺ cell; green) and UEA-1-binding goblet cell (red). Tissue counterstained to detect f-actin (blue). (b) M-cell density is dramatically reduced in the FAE of Peyer's patches from aged mice. Whole-mount images of Peyer's patches from each mouse group are shown. The positions of the X-Z and Y-Z projections of the FAE are indicated by the broken line in the main X-Y images. Closed arrows indicate GP2⁺ M cells (green).

830 Open arrow-heads indicate GP2⁻ UEA-1-binding goblet cells (red). (c) Morphometric analysis indicated that the density of M cells in the FAE of aged mice was significantly reduced. (d) Aging did not affect the density of goblet cells in the FAE. Data are derived from 3-5 Peyer's patches from at least 4 mice from each group (Kobayashi et al. 2013).

Figure 2 Mononuclear phagocytes (MNP) are abundant in the intestine.

Fluorescent confocal microscopy analysis of Peyer's patches in the intestines of a Tg(Csf1r-EGFP)1Hume "MacGreen" mouse in which the Csf1r promoter drives expression of enhanced green fluorescent protein (EGFP) in all MNP populations

840 (Sasmono et al. 2003). (a) By whole-mount imaging, abundant MNP expressing EGFP are detected in the lamina propria of villi and beneath the follicle-associated epithelium (FAE) of the Peyer's patches. (b) Analysis of tissue sections shows abundant MNP in the lamina propria of villi and in the sub-epithelial dome (SED) of the Peyer's patches.

Figure 3 Effects of aging on the density of CD11c⁺ cells in Peyer's patches. (a) IHC analysis of the distribution of CD11c⁺ cells (red) in the Peyer's patches of young and aged mice. Morphometric analysis suggested there was no difference in the density of CD11c⁺ cells in the sub-epithelial dome (SED) in Peyer's patches from each mouse group. (b) IHC analysis of the distribution of CD11c⁺ (red) and CD45R⁺ (B220, green) cells in FAE of Peyer's patches from young and aged mice. Arrows show CD11c⁺CD45R⁺ cells with apparent "M-cell-inducing" potential (Ebisawa et al. 2011). Morphometric analysis suggested the density of these CD11c⁺CD45R⁺ cells was significantly reduced in the FAE of aged mice. Data are derived from 3-5 Peyer's patches from at least 4 mice from each group (Kobayashi et al. 2013).

Figure 4 Total faecal IgA levels are elevated in aged mice. The concentration of total IgA in the faeces of aged mice is significantly elevated when compared to young mice (McDonald et al. 2011). Figure shows authors' own data from the analysis of individual faecal pellets from at least 4 mice from each group.

Figure 1
[Click here to download high resolution image](#)




Figure 3
[Click here to download high resolution image](#)




Figure 2
[Click here to download high resolution image](#)






Figure 4
[Click here to download high resolution image](#)



Authors responses to reviewers comments

[Click here to download Supplementary Material: Mabbott-BGEN-D-13-00135-AuthorResponses.pdf](#)

[Click here to view linked References](#)