



THE UNIVERSITY *of* EDINBURGH

Edinburgh Research Explorer

## Reassortant Avian Influenza A(H9N2) viruses in chickens in retail poultry shops, Pakistan, 2009-2010

**Citation for published version:**

Chaudhry, M, Angot, A, Rashid, HB, Cattoli, G, Hussain, M, Trovò, G, Drago, A, Valastro, V, Thrusfield, M, Welburn, S, Eisler, MC & Capua, I 2015, 'Reassortant Avian Influenza A(H9N2) viruses in chickens in retail poultry shops, Pakistan, 2009-2010' Emerging Infectious Diseases, vol. 21, no. 4, pp. 673-6. DOI: 10.3201/eid2104.141570

**Digital Object Identifier (DOI):**

[10.3201/eid2104.141570](https://doi.org/10.3201/eid2104.141570)

**Link:**

[Link to publication record in Edinburgh Research Explorer](#)

**Document Version:**

Publisher's PDF, also known as Version of record

**Published In:**

Emerging Infectious Diseases

**Publisher Rights Statement:**

All materials published in Emerging Infectious Diseases are in the public domain and can be used without permission

**General rights**

Copyright for the publications made accessible via the Edinburgh Research Explorer is retained by the author(s) and / or other copyright owners and it is a condition of accessing these publications that users recognise and abide by the legal requirements associated with these rights.

**Take down policy**

The University of Edinburgh has made every reasonable effort to ensure that Edinburgh Research Explorer content complies with UK legislation. If you believe that the public display of this file breaches copyright please contact [openaccess@ed.ac.uk](mailto:openaccess@ed.ac.uk) providing details, and we will remove access to the work immediately and investigate your claim.



# Reassortant Avian Influenza A(H9N2) Viruses in Chickens in Retail Poultry Shops, Pakistan, 2009–2010

Mamoona Chaudhry, Angélique Angot,  
Hamad B. Rashid, Giovanni Cattoli,  
Manzoor Hussain, Giulia Trovò,  
Alessandra Drago, Viviana Valastro,  
Michael Thrusfield, Sue Welburn,  
Mark C. Eisler, Ilaria Capua

Phylogenetic analysis of influenza viruses collected during December 2009–February 2010 from chickens in live poultry retail shops in Lahore, Pakistan, showed influenza A(H9N2) lineage polymerase and nonstructural genes generate through inter- and intrasubtypic reassortments. Many amino acid signatures observed were characteristic of human isolates; hence, their circulation could enhance inter- or intrasubtypic reassortment.

The first outbreak of illness caused by avian influenza A(H9N2) virus in Pakistan was reported in 1998; isolates showed a close relationship to subtype H9N2 avian influenza viruses (AIVs) circulating in Hong Kong, China during 1997 that were grouped within the G1 lineage (1). In recent years, H9N2 genes have reassorted extensively, generating novel genotypes on the Indian subcontinent. Widespread co-circulation of H9N2 with other AIVs (e.g., highly pathogenic AIVs H5N1 and H7N3) could instigate the generation of novel variant and reassorted viruses, possibly with increased zoonotic potential (2).

No information was available about the genetic make-up of AIVs circulating in live poultry retail shops (LPRSs) in Pakistan. This study was conducted to genetically characterize AIVs in LPRSs in Lahore District, Pakistan.

## The Study

We conducted a cross-sectional survey of LPRSs in Lahore (Figure), which is the capital of the Punjab Province in Pakistan. In each of 280 LPRSs, we collected tracheal


swab samples from 5 randomly selected chickens and pooled them into 1 composite sample, totaling 280 pooled samples. The samples were characterized at the World Organisation for Animal Health/Food and Agriculture Organization of the United Nations National Reference Laboratory for Avian Influenza in Padua, Italy.

Of the 280 samples, 10 tested positive for H9N2 subtype by real-time reverse transcription PCR (rRT-PCR) (3) and virus isolation test (United Nations [www.oie.int/fileadmin/Home/eng/Health\\_standards/tahm/2.03.04\\_AI.pdf](http://www.oie.int/fileadmin/Home/eng/Health_standards/tahm/2.03.04_AI.pdf)) (Table 1). Results were negative for H5 and H7 subtypes. Two of the 280 samples were positive for Newcastle disease virus. Each of the 10 H9N2 isolates was characterized by sequencing for 6 gene segments (hemagglutinin [HA]; neuraminidase [NA], nonstructural [NS], matrix [M], and polymerase basic 1 and 2 [PB1, PB2]); 2 isolates (A/chicken/Pakistan/10RS3039-283-11/2010 and A/chicken/Pakistan/10RS3039-289-186/2010) were sequenced for all 8 gene segments (HA, NA, NS, M, PB1, PB2, polymerase [PA], and nucleoprotein [NP]), as described (4). Sequences for these viruses were deposited into GenBank (accession nos. KF975457–KF975503, and KP223678–KP223693). The NS gene sequence of A/chicken/Pakistan/10RS3039-284-11/2010 could not be deposited due to poor data quality.

We generated neighbor-joining phylogenetic trees for all gene segments using the distance-based method in MEGA version 5.2.2 (<http://www.megasoftware.net/>). We calculated bootstrap values based on 1,000 replicates of alignment (5). HA and NA genes of viruses in this study tightly clustered within the G1 lineage along with H9N2 viruses from Pakistan, India, Iran, Israel, Saudi Arabia, and Bangladesh, suggesting derivation from a common ancestor: A/quail/Hong Kong/G1/97 (online Technical Appendix Figure panels A, B <http://wwwnc.cdc.gov/EID/article/21/4/14-1570-Techapp1.pdf>). M and NP genes clustered within the G1 lineage (online Technical Appendix Figure panels C, D) along with an influenza (H7N3) isolate from Pakistan (A/chicken/Karachi/NARC-100/2004; nucleotide identity >96% for M gene and 95% for NP gene). Two polymerase complex genes, PB1 and PA, and the NS gene did not cluster within G1 lineage or any other established Eurasian lineages: these gene sequences made a separate well-supported cluster with highly similar H9N2 viruses that circulated in Pakistan (2005–2008), Iran, India, and Bangladesh; these sequences had a high sequence

Author affiliations: University of Veterinary and Animal Sciences, Lahore, Pakistan (M. Chaudhry, H.B. Rashid); University of Edinburgh, Edinburgh, Scotland, UK (M. Chaudhry, M. Thrusfield, S. Welburn); Istituto Zooprofilattico Sperimentale delle Venezie, Viale dell'Università, Legnaro, Padua, Italy (A. Angot, G. Cattoli, G. Trovò, A. Drago, V. Valastro, I. Capua); Food and Agriculture Organization of the United Nations, Islamabad, Pakistan (M. Hussain); University of Bristol, Bristol, UK (M.C. Eisler)

DOI: <http://dx.doi.org/10.3201/eid2104.141570>



**Figure.** Location of live poultry retail shops (X) in 5 towns in Lahore, Pakistan, where avian influenza A(H9N2) virus isolates were identified in chickens, 2009–2010. Inset shows location of Lahore in Punjab Province.

identity (>95%) with A/chicken/Karachi/NARC-100/04 (H7N3) (2,6,7) (online Technical Appendix Figure panels E, F, G). The PB2 segment clustered separately with H9N2 viruses from the subcontinent and Middle East (online Technical Appendix Figure panel H), except A/chicken/Karachi/NARC-100/04. These results indicate a separate Indian subcontinental lineage of H9N2 viruses has emerged (2).

On the basis of these analyses, we could conclude that internal genes PB1, PA, and NS of these viruses originated by intersubtypic (between different HA subtypes) reassortment events from local H7N3 viruses circulating in Pakistan (nucleotide identity >95%). This suggests that intersubtypic reassortment events continuously result from mixing of AIV subtypes in domestic poultry and wild birds in Pakistan (8), and that PB2, M, and NP genes were acquired by intrasubtypic reassortment between H9N2 viruses of G1 lineage circulating within Pakistan. These results support

speculation that the currently circulating H9N2 lineage is a reassortment of G1 lineage from Hong Kong and the highly pathogenic H7N3 virus that circulated in Pakistan and can be assigned to genetic group B (8).

We aligned amino acid sequences of current viruses and compared them to representative H9N2 lineages. When compared to the prototype G1 viruses, H9N2 viruses isolated from LPRSs showed that they have evolved to acquire mammalian host-specific mutations throughout the genome (online Technical Appendix Table 1). Of these mutations, certain amino acid substitutions throughout the viral genome have become fixed (9). All LPRS isolates possessed the K-S-S-R motif at the cleavage site. The presence of lysine at position 4 was observed in H9 isolates from the Indian subcontinent (2,9), but it is uncommon elsewhere. Amino acid signature changes of human influenza viruses were also observed in internal gene segments (Table 2).

**Table 1.** Influenza A(H9N2) viruses isolated from oropharyngeal swab samples from chickens in live poultry retail shops of Lahore, Pakistan, 2010

Isolates	Collection date	Towns of Lahore District*
A/chicken/Pakistan/10RS3039-283-11/2010	Jan 23	Gulberg Town
A/chicken/Pakistan/10RS3039-284-48/2010	Jan 25	Data Gunj Bakhsh Town
A/chicken/Pakistan/10RS3039-285-63/2010	Jan 26	Data Gunj Bakhsh Town
A/chicken/Pakistan/10RS3039-286-65/2010	Jan 26	Data Gunj Bakhsh Town
A/chicken/Pakistan/10RS3039-287-98/2010	Jan 27	Ravi Town
A/chicken/Pakistan/10RS3039-288-102/2010	Jan 27	Ravi Town
A/chicken/Pakistan/10RS3039-289-186/2010	Feb 11	Samanabad Town
A/chicken/Pakistan/10RS3039-199-199/2010	Feb 11	Samanabad Town
A/chicken/Pakistan/10RS3039-290-230/2010	Feb 13	Allama Iqbal Town
A/chicken/Pakistan/10RS3039-291-266/2010	Feb 14	Allama Iqbal Town

\*Lahore District contains 9 towns and 1 cantonment area.

**Table 2.** Amino acid signature changes observed in M1, NP, PB1 and NS1 proteins of influenza A(H9N2) viruses isolated from chickens in live poultry retail shops, Lahore, Pakistan, 2009–2010\*

Protein	Amino acid position	Predicted aa		Viruses with detected mutation	Mutation
		Avian	Human		
M1	15	V	I	All 10 isolates analyzed in this study	V15I
NP	372	E	D	Only 2 viruses were analyzed (A/chicken/ Pakistan/10RS3039-283-11/2010 A/chicken/ Pakistan/10RS3039-289-186/2010)	E372D
PB1	13	L	P	All 10 isolates analyzed in this study	L13P
NS1	149	V	A	All 10 isolates analyzed in this study	V149A
	227	E	R or K (H1N1 1918)		E227K

\*M, matrix; NP, nucleoprotein; PB, basic polymerase; NS, nonstructured.

Each of the 10 isolates had the Q<sup>226</sup>L substitution (H3 numbering) in the receptor-binding site of HA, correlating to a shift in affinity from avian-type sialic acid receptor to human-type (10). We identified 3 representative substitutions: E/T<sup>190</sup>A in 9 viruses and E/T<sup>190</sup>V in 1 virus, and Q<sup>227</sup>I in all 10 viruses (H3 numbering). The outcomes of these substitutions have not been investigated; further study is needed (2). Glycosylation sites at positions 551, 218, and 206 were absent in the study viruses, suggesting a frequent alteration in sequences from this region (2,6,9) and possibly signifying the selected adaptation of H9N2 to poultry (10). We did not find an R<sup>292</sup>K substitution, which is associated with resistance to the sialidase inhibitors oseltamivir and zanamivir, in the NA proteins of any of the 10 LPRS virus isolates. HB sites were also well conserved with few substitutions (K<sup>367</sup>E in 5 viruses, K<sup>367</sup>G in 1 virus, and S<sup>372</sup>A and W<sup>403</sup>R in all 10 viruses). An additional glycosylation site was present at position 44, which is believed to enhance virulence caused by altered antigenicity or sialidase activity (11) (online Technical Appendix Table 1).

Many residues in nucleoprotein and polymerase are considered determinants of the host range of AIVs and increase virulence or replication in the mammalian host (2,10,12). The analyses of internal genes showed that these viruses also contained mammalian host-specific markers (2,8) that have become permanent in M protein (M1, V<sup>15</sup>I, T<sup>37</sup>A; M2, E<sup>16</sup>G, L<sup>55</sup>F); in PB1 protein (L<sup>13</sup>P); and in NS1 protein (V<sup>149</sup>A) of all LPRS isolates and in NP (E<sup>372</sup>D) of the 2 isolates we sequenced (online Technical Appendix Table 2). In M2 protein, no substitutions linked to resistant amantadine were seen. All viruses contained an uncommon K-S-E-I sequence as a PDZ ligand motif in the NS protein. Residue isoleucine at the C-terminus of PDZ ligand motif has been reported as a rare substitution (12). NS also harbored E<sup>227</sup>K mutation in the C-terminal, which has been demonstrated to modulate pathogenicity of AIVs (13) and appeared to be a rare amino acid signature, although in this study it was observed in all 10 virus isolates from chickens in LPRSs.

## Conclusions

Our analysis confirmed that continuous gene reassortment has occurred among influenza A(H9N2) viruses since their

emergence in poultry in Pakistan. Because H9N2 viruses infect multiple species, they may donate genes to emerging human pathogens; it has been observed that H7N9 acquired internal genes from the avian H9N2 virus (14). In wet markets, availability of freshly slaughtered poultry, live poultry transportation, and mixed trading of domestic animals provide a favorable environment for gene reassortment, mutation, and interspecies transmission of AIVs (15). Continuous circulation of these viruses in LPRSs increases the chances of their evolution into new genotypes. Close contact of humans and poultry in LPRSs with no biosecurity barriers increases the risk for emergence of novel influenza viruses with zoonotic or human pandemic potential. Continued surveillance in LPRSs is essential to better understand the public health risk posed by H9N2 AIVs.

## Acknowledgments

We thank the staff members of the live poultry retail shops in which the study subjects were housed for data and sample collection.

The European Union project Training and technology transfer of avian influenza diagnostics and disease management skills (FLUTRAIN) provided financial support for training of MC at IZSve, Italy. The Higher Education Commission of Pakistan provided a PhD scholarship to MC through the Overseas Scholarship Scheme (10%- Batch-2).

We acknowledge the memory of Isabel Minguez Tudela (1955–2011), of the European Commission, who supported this collaboration as a contribution to improving veterinary public health.


Dr. Chaudhry is an academician in the Department of Epidemiology and Public Health, University of Veterinary and Animal Sciences, Lahore. Her research interests include epidemiology of infectious diseases of public health concern.

## References

1. Cameron KR, Gregory V, Banks J, Brown IH, Alexander DJ, Hay AJ, et al. H9N2 subtype influenza A viruses in poultry in Pakistan are closely related to the H9N2 viruses responsible for human infection in Hong Kong. *Virology*. 2000;278:36–41. <http://dx.doi.org/10.1006/viro.2000.0585>

2. Iqbal M, Yaqub T, Reddy K, McCauley JW. Novel genotypes of H9N2 influenza A viruses isolated from poultry in Pakistan containing NS genes similar to highly pathogenic H7N3 and H5N1 viruses. *PLoS ONE*. 2009;4:e5788. <http://dx.doi.org/10.1371/journal.pone.0005788>
3. Monne I, Ormelli S, Salviato A, De Battisti C, Bettini F, Salomoni A, et al. Development and validation of a one-step real-time PCR assay for simultaneous detection of subtype H5, H7, and H9 avian influenza viruses. *J Clin Microbiol*. 2008; 46:1769–73. <http://dx.doi.org/10.1128/JCM.02204-07>
4. Cattoli G, Monne I, Fusaro A, Joannis TM, Lombin LH, Aly MM, et al. Highly pathogenic avian influenza virus subtype H5N1 in Africa: a comprehensive phylogenetic analysis and molecular characterization of isolates. *PLoS ONE*. 2009;4:e4842. <http://dx.doi.org/10.1371/journal.pone.0004842>
5. Tamura K, Dudley J, Nei M, Kumar S. MEGA4: Molecular Evolutionary Genetics Analysis (MEGA) software version 4.0. *Mol Biol Evol*. 2007;24:1596–9. <http://dx.doi.org/10.1093/molbev/msm092>
6. Aamir UB, Wernery U, Ilyushina N, Webster RG. Characterization of avian H9N2 influenza viruses from United Arab Emirates 2000 to 2003. *Virology*. 2007;361:45–55. <http://dx.doi.org/10.1016/j.virol.2006.10.037>
7. Abbas MA, Spackman E, Swayne DE, Ahmed Z, Sarmento L, Siddique N, et al. Sequence and phylogenetic analysis of H7N3 avian influenza viruses isolated from poultry in Pakistan 1995–2004. *Virology*. 2010;7:137. <http://dx.doi.org/10.1186/1743-422X-7-137>
8. Fusaro A, Monne I, Salviato A, Valastro V, Schivo A, Amarin NM, et al. Phylogeography and evolutionary history of reassortant H9N2 viruses with potential human health implications. *J Virol*. 2011;85:8413–21. <http://dx.doi.org/10.1128/JVI.00219-11>
9. Shanmuganatham K, Feeroz MM, Jones-Engel L, Smith GJ, Fourment M, Walker D, et al. Antigenic and molecular characterization of avian influenza A(H9N2) viruses, Bangladesh. *Emerg Infect Dis*. 2013;19:1393–1402. <http://dx.doi.org/10.3201/eid1909.130336>
10. Baigent SJ, McCauley JW. Influenza type A in humans, mammals and birds: determinants of virus virulence, host-range and interspecies transmission. *Bioessays*. 2003;25:657–71. <http://dx.doi.org/10.1002/bies.10303>
11. Hulse DJ, Webster RG, Russell RJ, Perez DR. Molecular determinants within the surface proteins involved in the pathogenicity of H5N1 influenza viruses in chickens. *J Virol*. 2004;78:9954–64. <http://dx.doi.org/10.1128/JVI.78.18.9954-9964.2004>
12. Finkelstein DB, Mukatira S, Mehta PK, Obenauer JC, Su X, Webster RG, et al. Persistent host markers in pandemic and H5N1 influenza viruses. *J Virol*. 2007;81:10292–9. <http://dx.doi.org/10.1128/JVI.00921-07>
13. Obenauer JC, Denson J, Mehta PK, Su X, Mukatira S, Finkelstein DB, et al. Large-scale sequence analysis of avian influenza isolates. *Science*. 2006;311:1576–80. <http://dx.doi.org/10.1126/science.1121586>
14. Gao R, Cao B, Hu Y, Feng Z, Wang D, Hu W, et al. Human infection with a novel avian-origin influenza A (H7N9) virus. *N Engl J Med*. 2013;368:1888–97. <http://dx.doi.org/10.1056/NEJMoa1304459>
15. Bao CJ, Cui LB, Zhou MH, Hong L, Gao GF, Wang H. Live-animal markets and influenza A (H7N9) virus infection. *N Engl J Med*. 2013;368:2337–9. <http://dx.doi.org/10.1056/NEJMc1306100>

Address for correspondence: Mamoona Chaudhry, Department of Epidemiology and Public Health, University of Veterinary and Animal Sciences, Abdul Qadir Jilani Road, Lahore, Pakistan; email: mamoona.chaudhry@gmail.com



## February 2014: High-Consequence Pathogens

Including:

- Poxvirus Viability and Signatures in Historical Relics
- Novel Paramyxovirus Associated with Severe Acute Febrile Disease, South Sudan and Uganda, 2012
- Subtyping *Cryptosporidium ubiquitum*, a Zoonotic Pathogen Emerging in Humans
- Genomic Variability of Monkeypox Virus among Humans, Democratic Republic of the Congo

<http://wwwnc.cdc.gov/eid/content/20/2/contents.htm>