



# THE UNIVERSITY *of* EDINBURGH

## Edinburgh Research Explorer

### Experimental evolution gone wild

**Citation for published version:**

Scheinin, M, Riebesell, U, Rynearson, TA, Lohbeck, KT & Collins, S 2015, 'Experimental evolution gone wild' *Journal of the Royal Society Interface*, vol. 12, no. 106, 20150056. DOI: 10.1098/rsif.2015.0056

**Digital Object Identifier (DOI):**

[10.1098/rsif.2015.0056](https://doi.org/10.1098/rsif.2015.0056)

**Link:**

[Link to publication record in Edinburgh Research Explorer](#)

**Document Version:**

Publisher's PDF, also known as Version of record

**Published In:**

*Journal of the Royal Society Interface*

**Publisher Rights Statement:**

© 2015 The Authors. Published by the Royal Society under the terms of the Creative Commons Attribution License <http://creativecommons.org/licenses/by/4.0/>, which permits unrestricted use, provided the original author and source are credited.

**General rights**

Copyright for the publications made accessible via the Edinburgh Research Explorer is retained by the author(s) and / or other copyright owners and it is a condition of accessing these publications that users recognise and abide by the legal requirements associated with these rights.

**Take down policy**

The University of Edinburgh has made every reasonable effort to ensure that Edinburgh Research Explorer content complies with UK legislation. If you believe that the public display of this file breaches copyright please contact [openaccess@ed.ac.uk](mailto:openaccess@ed.ac.uk) providing details, and we will remove access to the work immediately and investigate your claim.



CrossMark  
click for updates

**Cite this article:** Scheinin M, Riebesell U, Rynearson TA, Lohbeck KT, Collins S. 2015 Experimental evolution gone wild. *J. R. Soc. Interface* **12**: 20150056.  
<http://dx.doi.org/10.1098/rsif.2015.0056>

Received: 22 January 2015

Accepted: 6 March 2015

**Subject Areas:**

biogeochemistry, environmental science, systems biology

**Keywords:***Skeletonema*, diatom evolution, *in situ* mesocosms, carbon dioxide, ocean acidification, experimental evolution**Author for correspondence:**

S. Collins

e-mail: [s.collins@ed.ac.uk](mailto:s.collins@ed.ac.uk)

Electronic supplementary material is available at <http://dx.doi.org/10.1098/rsif.2015.0056> or via <http://rsif.royalsocietypublishing.org>.

## Experimental evolution gone wild

M. Scheinin<sup>1,2</sup>, U. Riebesell<sup>1</sup>, T. A. Rynearson<sup>3</sup>, K. T. Lohbeck<sup>1</sup> and S. Collins<sup>4</sup><sup>1</sup>GEOMAR Helmholtz Centre for Ocean Research Kiel, Düsternbrooker Weg 20, Kiel, Germany<sup>2</sup>Tvärminne Zoological Station, University of Helsinki, J. A. Palménin tie 260, Hanko, Finland<sup>3</sup>Graduate School of Oceanography, University of Rhode Island, Narragansett, RI, USA<sup>4</sup>Institute of Evolutionary Biology, University of Edinburgh, Ashworth Laboratories, The King's Building's, West Mains Road, Edinburgh, UK

Because of their large population sizes and rapid cell division rates, marine microbes have, or can generate, ample variation to fuel evolution over a few weeks or months, and subsequently have the potential to evolve in response to global change. Here we measure evolution in the marine diatom *Skeletonema marinoi* evolved in a natural plankton community in CO<sub>2</sub>-enriched mesocosms deployed *in situ*. Mesocosm enclosures are typically used to study how the species composition and biogeochemistry of marine communities respond to environmental shifts, but have not been used for experimental evolution to date. Using this approach, we detect a large evolutionary response to CO<sub>2</sub> enrichment in a focal marine diatom, where population growth rate increased by 1.3-fold in high CO<sub>2</sub>-evolved lineages. This study opens an exciting new possibility of carrying out *in situ* evolution experiments to understand how marine microbial communities evolve in response to environmental change.

**1. Introduction**

Experimental evolution is a method that uses replicate populations, in controlled environments, to measure evolution in real time [1]. The power of this approach is that first, it produces generalizable results that further our understanding of how natural selection and evolution work—it allows us to uncover the rules that evolution plays by. This is partly because analyses focus on fitness, which is what natural selection acts on, regardless of particular genetic, epigenetic and phenotypic changes that underlie fitness shifts. Second, the experimenter manipulates the environment and uses replicate populations, so environmental changes can be linked causally to evolutionary responses [2]. However, experimental evolution has largely been confined to laboratory populations, and there are few experimental evolution studies on natural microbial populations *in situ*. Here, we show that marine mesocosms can be used for microbial evolution experiments by measuring evolution in a marine diatom in CO<sub>2</sub>-enriched marine mesocosms. This provides a link between laboratory evolution experiments and natural populations by using enclosures that are tractable, controllable and offer replication, but which also keep focal species in the context of a more natural community and habitat than is possible in the laboratory.

One limitation of experimental evolution stems from the same characters that give it its power: such replication, control and tractability can usually only be achieved under laboratory conditions, leading to a trade-off between uncovering general evolutionary mechanisms and understanding how they apply in complex natural environments, which in turn limits our understanding of how natural populations evolve in response to particular environmental drivers [2,3]. To make predictions about evolution in natural populations, it is vital that we link laboratory experiments to field studies. We propose that the most obvious way to do this is by conducting evolution experiments *in situ*. This requires the following criteria to be met: a starting population needs to be divided among independent replicate control and treatment environments. In addition, to measure evolution directly, rather than infer it from population genetics, the focal organisms need to reproduce quickly enough (or experience

enough selective mortality) to measure heritable changes in fitness or genotype frequencies over an experiment. Previous evolution experiments in natural populations have taken advantage of natural replicate selection environments, such as stream systems where fish populations can be transplanted, for example to study predator/prey evolution [4]. However, this relies on finding fortuitous replicate environments (streams). We show that mesocosm enclosures fulfil the above criteria and can be coopted for microbial experimental evolution.

Marine mesocosms are commonly used to study community-level responses to environmental changes such as CO<sub>2</sub> enrichment [5,6]. They are analogous to laboratory evolution experiments in that replicate mesocosms enclose random samples of the same aquatic community, which are subjected to an environmental change (e.g. CO<sub>2</sub> enrichment) or not (control mesocosms). Because the mesocosms are closed, biotic or abiotic changes within them can be causally linked to the environmental manipulation. Here, we measure the evolutionary response to CO<sub>2</sub> enrichment in a focal species of marine diatom during a mesocosm experiment.

## 2. Material and methods

### 2.1. Mesocosm set-up

A mesocosm study was conducted in the Gullmar Fjord on the west coast of Sweden (58°15.9' N, 11°28.9' E) in the framework of the German national project on 'Biological Impacts of Ocean Acidification' (BIOACID) (figure 1). Detailed information about the mesocosm design and experimental application is provided in [7]. Briefly, 10 mesocosms were deployed in the Gullmar Fjord by R/V Alkor on 29 January 2013. The 18 m long enclosure bags were filled with fjord water after the retreat of sea-ice on 7 March, well before the start of the spring phytoplankton bloom. Each mesocosm bag enclosed about 55 m<sup>3</sup> of fjord water, including the natural plankton community present at the time of closure. While five mesocosms were kept untreated as controls, the carbonate chemistry in the remaining five mesocosms was manipulated to establish elevated *p*CO<sub>2</sub> at an initial level of 1100 µatm (for details on the manipulation approach, see Riebesell *et al.* [7]). The mesocosms were sampled every second day from 8 March to 28 June, covering a period of 107 days. In total, 45 parameters were measured in all mesocosms, providing a detailed overview of the environmental conditions in the mesocosms and the development of the enclosed plankton community.

### 2.2. Cell isolation

We attempted to isolate cells from all 10 mesocosms. Viable samples were obtained from five of the high CO<sub>2</sub> mesocosms and three of the control mesocosms. Our experiment required isolating the same species of diatom from the majority of the mesocosms. *Skeletonema marinoi* was present in most mesocosms by the end of the experiment, although densities were low. Individual chains of *S. marinoi* were isolated from mesocosms and used to obtain monoclonal cultures. Samples were taken after the end of the mesocosm experiment (days 107–111) because net hauls would have been disruptive to other studies taking place at the same time. Plankton nets (mesh size 10 µm) were hauled over the whole depth of each mesocosm at 0.5 m s<sup>-1</sup>. Four hauls were done per mesocosm; total volume covered per mesocosm was 1064 l. Nets were emptied into 20 l carboys pre-filled with filtered (0.2 µm) mesocosm water. Sampling gear was sterilized between mesocosms by soaking in 80% ethanol and rinsing in Milli-Q water. The carboys were stored in the dark at 10°C for 1–4 days before isolations were conducted.

Cells were isolated on a 5 µm mesh, and collected by rinsing the mesh with sterile-filtered mesocosm water into Petri dishes.

*Skeletonema marinoi* was visually identified using an inverted light microscope (Leica DMIL). No other species of *Skeletonema* have been reported in the Gullmar Fjord (Swedish Meteorological and Hydrological Institute database SHARK/Svenskt HavsARKiv). Individual chains of cells were isolated with a 10 µl pipette and placed in a single well of a 24 well plate in 1 ml of f/8 medium [8] made from sterile-filtered water pooled from all 10 mesocosms. Growing isolates were transferred to 50 ml culture flasks containing 20 ml of media. Cultures were grown at 10°C at 100 µmol photons m<sup>-2</sup> s<sup>-1</sup> on a 12 L:12 D cycle, and 20 µl of culture was transferred into fresh media every 5 days. This was done in the Sven Lovén Centre for Marine Sciences in Kristineberg, Sweden. There was a laboratory contamination event approximately five weeks after cells were isolated, and cultures were cleaned by reisolating *S. marinoi* cells. Surviving uncontaminated (cleaned) isolates are used for all work below. For the results reported in this manuscript, the final numbers of isolates from the three control mesocosms were 5, 3 and 7 isolates. The final numbers of isolates from the high CO<sub>2</sub> mesocosms were 3, 8, 0, 9 and 11 isolates.

### 2.3. Laboratory culture conditions


Cultures were moved to the GEOMAR Helmholtz Centre for Ocean Research Kiel, Germany, for growth assays, and acclimated over 20 days to f/8 medium with artificial seawater [9] by replacing half the medium at each transfer over four transfers. In the final medium, 2% of the volume was seawater from the mesocosms, salinity was 35, and total alkalinity (TA) was adjusted to 2380 µEq l<sup>-1</sup> using sodium bicarbonate. Cultures were kept in incubators (RUMED Light Thermostat Type 1201) at 5°C with 90–100 µmol photons m<sup>-2</sup> s<sup>-1</sup> on a 12 L:12 D cycle. Temperature was changed in one step. This temperature is closer to the temperature of the Gullmar Fjord during the mesocosm experiment; it was not possible to culture at 5°C at the Sven Lovén Centre. Growth rates were measured under these conditions. Cultures were acclimated to growth conditions for four 5-day transfers prior to measurement.

### 2.4. Carbonate chemistry manipulations

Carbon dioxide concentrations were manipulated by bubbling media with 400 µatm *p*CO<sub>2</sub> or 2400 µatm *p*CO<sub>2</sub> air for 48 h, and then mixing these in appropriate proportions to make 400 or 1000 µatm *p*CO<sub>2</sub> growth media. The resulting CO<sub>2</sub> concentrations were verified as follows: CO<sub>2</sub> concentration and dissolved inorganic carbon (DIC) were calculated from TA and pH by measuring the pH of the media at 5°C (3 × 60 ml samples per flask), using CO2SYS (v. 2.1) and accounting for phosphate and silicate concentrations in the growth medium.

### 2.5. Growth rate measurements

For growth assays, three isolates were randomly chosen per mesocosm that we had uncontaminated samples from. We measured growth rates at 400 and 1000 µatm *p*CO<sub>2</sub>. Cultures were inoculated at 30 cells ml<sup>-1</sup> in a total volume of 65 ml of f/8 medium, and placed in the incubator in a random order. Cells were counted microscopically using an Utermöhl chamber [10] at 0 and 3 days. The cell division rates were calculated as  $T_d = \ln(d_2/d_1)/(\Delta t \cdot \ln 2)$ , where  $T_d$  is the doubling rate,  $d_1$  the initial cell density,  $d_2$  the final cell density and  $\Delta t$  the total time for the observations. DIC-drawdown was kept below 2% during acclimation and below 1% during growth measurements. This was verified by measuring pH and as described in the Carbonate chemistry manipulations section. Cultures were growing exponentially during acclimation and growth measurements.



**Figure 1.** (a) Single mesocosm unit, consisting of flotation frame, mesocosm bag and sediment trap. (b) Experimental set-up, consisting of 10 mesocosm units, of which five are kept at ambient  $p\text{CO}_2$  level of approximately  $400 \mu\text{atm}$  (control, numbers 1, 3, 5, 9 and 10) and five are manipulated to yield a  $p\text{CO}_2$  level of approximately  $1000 \mu\text{atm}$  projected for the end of this century in the case of unabated  $\text{CO}_2$  emissions (numbers 2, 4, 6, 7 and 8). (Online version in colour.)

## 2.6. Statistical analyses

Data were analysed as a mixed model in an R environment using the nlme package [11]. The response variable is growth rate in the laboratory. The effects included are mesocosm  $\text{CO}_2$  level, laboratory  $\text{CO}_2$  level, mesocosm identity and clone. Mesocosm and laboratory  $\text{CO}_2$  levels were modelled as fixed effects with interaction. Mesocosm identity and clone identity were random effects, with clone nested within mesocosm identity. Note that all results (evolutionary and plastic responses) are discussed using the same statistical test in order to avoid multiple tests on the same dataset. Data are available as an online data supplement.

## 3. Results

There is a direct response to selection for growth in a high  $\text{CO}_2$  environment in *S. marinoi* (effect of mesocosm  $\text{CO}_2 \times$  laboratory  $\text{CO}_2$ :  $t_{1,61} = -3.45$ ,  $p = 0.001$ ). The direct response to selection is measured by comparing the growth of the high  $\text{CO}_2$ -evolved lineages and ambient  $\text{CO}_2$ -evolved lineages when grown in high  $\text{CO}_2$  conditions in the laboratory (figure 2), and it reflects heritable differences in growth in a stable environment that are attributable to having evolved in different environments. The shorter doubling times of the high  $\text{CO}_2$ -evolved lineages indicate that *S. marinoi* from high  $\text{CO}_2$  mesocosms have evolved in response to high  $\text{CO}_2$ , and they divide about 1.3 times faster under high  $\text{CO}_2$  laboratory conditions than do lineages from control mesocosms. The doubling rate for lineages from high  $\text{CO}_2$  mesocosms is  $20.53 \pm 1.87$  h (mean  $\pm$  s.d.) at high  $\text{CO}_2$ , while the doubling rates for lineages from control mesocosms is  $24.32 \pm 3.89$  h at high  $\text{CO}_2$ .


A plastic response is a change in phenotype of a single genotype to environmental change. Here, this corresponds to the difference in growth rates of a single isolate when it is grown in ambient versus high  $\text{CO}_2$  conditions in the laboratory, and reflects the ability of lineages to respond to changes in  $\text{CO}_2$ . Overall, the plastic response to short-term changes in  $\text{CO}_2$  levels is to increase growth rates (same analysis as

above; effect of laboratory  $\text{CO}_2$  level  $t_{1,61} = -2.19$ ,  $p = 0.0318$ ). This is driven by the responses of the lineages evolved in the high  $\text{CO}_2$  mesocosms (interaction between laboratory and mesocosm  $\text{CO}_2$  levels  $t_{1,61} = -3.45$ ,  $p = 0.001$ ; figure 2). While isolates of *S. marinoi* evolved in control mesocosms do not show a plastic growth response to  $\text{CO}_2$  enrichment, lineages evolved in the high  $\text{CO}_2$  mesocosms do. This shows that the plastic response to rapid changes in  $\text{CO}_2$  has evolved in the high  $\text{CO}_2$  mesocosms. Full model output is in the electronic supplementary material, appendix.

## 4. Discussion

Experimental evolution allows researchers to watch evolution in real time, and connect evolutionary responses to environmental drivers. Here, we show that microbial evolution experiments can be carried out in enclosed natural plankton communities, where experimental design and measures of evolutionary responses are the same as in laboratory experiments. These *in situ* experiments can be directly compared with laboratory experiments to link general mechanisms to particular outcomes.

Our focal species, the marine diatom *S. marinoi*, evolved in response to growth under high  $\text{CO}_2$  conditions for over 100 days as part of an enclosed microbial community. Both the growth rate at high  $\text{CO}_2$  and the plastic response to changes in  $\text{CO}_2$  levels evolved. The direct response to selection was large, with lineages evolved at high  $\text{CO}_2$  having a 1.3 $\times$  growth advantage over lineages from the control mesocosms when both were grown at high  $\text{CO}_2$  under laboratory conditions. Since the evolutionary response to selection is to increase growth, it is likely to be adaptive, or part of a more complex phenotypic change that is, on balance, adaptive [1]. While our results show unambiguously that evolution occurred in response to high  $\text{CO}_2$ , the fitness advantage associated with it within the mesocosms cannot be reasonably extrapolated from growth rates in the laboratory. That being said, if fitness were determined entirely by



**Figure 2.** Cell division rates in hours per day for *S. marinoi* at 400 and 1000  $\mu\text{atm}$   $p\text{CO}_2$  in laboratory growth experiments.  $\text{CO}_2$  levels in top grey panels indicate the level of  $\text{CO}_2$  in the mesocosm where the lineages evolved.  $\text{CO}_2$  levels indicated on the bottom x-axis indicate  $\text{CO}_2$  level under which growth was measured in the laboratory. Points show cell division rates for individual lineages.

growth rate, this would translate into about a 33% fitness advantage. However, since growth is not the only component of fitness, this is likely to be an overestimate, especially if faster growing lineages are more likely to be grazed. Because high- and control- $\text{CO}_2$  mesocosms also differ in their communities [12] and abiotic environment [13] as a result of the  $\text{CO}_2$  manipulation, we cannot say how much of the evolutionary response to  $\text{CO}_2$  enrichment is directly driven by  $\text{CO}_2$  versus indirectly. A parallel laboratory experiment where *S. marinoi* evolved in environments that differ only in  $\text{CO}_2$  levels (e.g. [3]) would be needed to partition the evolutionary response into components attributable to direct and indirect drivers.

Our results raise the possibility that local changes in  $\text{CO}_2$  levels could drive adaptation in local populations [14]. Interestingly, the asymmetry in the responses of the diatoms from the control and high  $\text{CO}_2$  mesocosms, where the high  $\text{CO}_2$ -evolved lineages outgrow the control lineages at high  $\text{CO}_2$ , but the control lineages do not outgrow the high  $\text{CO}_2$ -evolved lineages at control levels of  $\text{CO}_2$ , has been seen in some evolution experiments using high  $\text{CO}_2$  as a driver for phytoplankton evolution, though in other cases, high  $\text{CO}_2$ -evolved lineages grew poorly or died at ambient  $\text{CO}_2$  (reviewed in [3]). Elevated  $\text{CO}_2$  may be able to drive local adaptation even if increases in growth rates are transient or absent, since marine picoplankton evolved for hundreds of generations in a high  $\text{CO}_2$  environment maintained an increase in competitive ability even when they did not show increased growth in the absence of competitors in laboratory high  $\text{CO}_2$  environments [15].

Previous studies show that plastic responses to  $\text{CO}_2$  enrichment are idiosyncratic between, and even within, diatom species [16], reporting that *Skeletonema* spp. can respond

plastically to changes in  $\text{CO}_2$  by increasing growth [17] or not [16]. However, the composition of synthetic [18] and natural [19] diatom assemblages changes in response to  $\text{CO}_2$  enrichment, indicating that shifts in relative fitness can be large enough to allow evolution in such assemblages. We find that even though the plastic response to  $\text{CO}_2$  enrichment in *S. marinoi* isolated from control mesocosms is absent, lineages evolved in high  $\text{CO}_2$  mesocosms both respond plastically to  $\text{CO}_2$  enrichment and grow faster at high  $\text{CO}_2$ . This is in line with studies in green algae showing that more plastic lineages are likely to be selected in novel environments [15]. The maximum number of generations of *S. marinoi* possible in the mesocosm experiment was approximately 100, making it unlikely that novel mutations fuelled evolution here. Because dominant mutations of very large effect could have had time to fix had they arisen early in the mesocosm experiment, we cannot rule out the possibility that novel genetic variation arose during the mesocosm experiment. However, our data suggest that it is more likely that natural selection acted predominantly on pre-existing variation, favouring more plastic genotypes in the high  $\text{CO}_2$  environment and less plastic genotypes in the control environment. Our reasoning is that the fastest-growing high  $\text{CO}_2$ -evolved lineages are within the range of the control-evolved lineages, even though the average growth rate is faster. In addition, based on the sampling effort required to do this study, populations of *S. marinoi* were relatively small in the mesocosms, meaning that the supply of novel mutations would have also been low. This, alongside the variation seen among lineages in terms of plastic responses, suggests that there is substantial within-population variation in plastic responses to changes in  $\text{CO}_2$  in this species.

Ocean change research has made the strongest progress in recent years at the level of single species or strains with respect to their plastic (short-term) responses to single environmental changes. It is, however, the evolutionary (long-term) response of natural communities to a multitude of environmental alterations that we need to understand to make reliable predictions of future changes in marine ecosystems. Providing this information by stepping up from single to multiple drivers, from single strains to communities and ecosystems, and from plastic to evolutionary responses is a major challenge. Using mesocosm studies for experimental evolution offers a way to investigate evolutionary outcomes in natural populations that is directly comparable with laboratory evolution experiments, linking

evolution in single species and community experiments. This study shows that investigating evolutionary adaptation at the community level in near-natural environmental settings is feasible and that approaches such as the one taken here will help paint a more realistic picture of the future of ocean ecosystems.

**Acknowledgements.** We thank L. Bach, T. Boxhammer, S. Febiri, M. Haunost, A. Ludwig, J. Scheinin and M. Sswat for technical assistance in the laboratory and during the mesocosm campaign.

**Funding statement.** This project was funded by the BMBF project 'Biological Impacts of Ocean Acidification' (BIOACID) and through the Leibniz Award 2012 of the German Science Foundation (DFG) to U.R. and a Royal Society UK University Research Fellowship to S.C.

## References

- Elena SF, Lenski RE. 2003 Evolution experiments with microorganisms: the dynamics and genetic bases of adaptation. *Nat. Rev. Genet.* **4**, 457–469.
- Merilä J, Hendry AP. 2014 Climate change, adaptation, and phenotypic plasticity: the problem and the evidence. *Evol. Appl.* **7**, 1–14. (doi:10.1111/eva.12137)
- Collins S, Rost B, Ryneerson TA. 2013 Evolutionary potential of marine phytoplankton under ocean acidification. *Evol. Appl.* **7**, 140–155. (doi:10.1111/eva.12120)
- Bassar RD, Lopez-Sepulcre A, Reznick DN, Travis J. 2013 Experimental evidence for density-dependent regulation and selection on trinidadian guppy life histories. *Am. Nat.* **181**, 25–38. (doi:10.1086/668590)
- Odum EP. 1984 The mesocosm. *BioScience* **34**, 558–562. (doi:10.2307/1309598)
- Riebesell U, Bellerby RGJ, Grossart HP, Thingstad F. 2008 Mesocosm CO<sub>2</sub> perturbation studies: from organism to community level. *Biogeosciences* **5**, 1157–1164. (doi:10.5194/bg-5-1157-2008)
- Riebesell U *et al.* 2012 Technical note: a mobile sea-going mesocosm system; new opportunities for ocean change research. *Biogeosci. Discuss.* **9**, 12 985–13 017. (doi:10.5194/bgd-9-12985-2012)
- Guillard RRL. 1975 *Culture of phytoplankton for feeding marine invertebrates*. (link.springer.com), pp. 29–60. Boston, MA: Springer US.
- Kester DR, Duedall IW, Connors DN, Pytkowicz RM. 1967 Preparation of artificial seawater. *Limnol. Oceanogr.* **12**, 176–179. (doi:10.4319/lo.1967.12.1.0176)
- Utermöhl H. 1958 Zur Vervollkommnung der quantitativen Phytoplankton-Methodik. *Mitt. Int. Ver. Theor. Angew. Limnol.* **9**, 1–38.
- R Core Team. 2014 *R: a language and environment for statistical computing*. Vienna, Austria: R Foundation for Statistical Computing.
- Brussaard CPD, Noordeloos AAM, Witte H, Collenteur MCJ, Schulz K, Ludwig A, Riebesell U. 2013 Arctic microbial community dynamics influenced by elevated CO<sub>2</sub> levels. *Biogeosciences* **10**, 719–731. (doi:10.5194/bg-10-719-2013)
- Schulz KG *et al.* 2013 Temporal biomass dynamics of an Arctic plankton bloom in response to increasing levels of atmospheric carbon dioxide. *Biogeosciences* **10**, 161–180. (doi:10.5194/bg-10-161-2013)
- Blanquart F, Kaltz O, Nuismer SL, Gandon S. 2013 A practical guide to measuring local adaptation. *Ecol. Lett.* **16**, 1195–1205. (doi:10.1111/ele.12150)
- Schaum CE, Collins S. 2014 Plasticity predicts evolution in a marine alga. *Proc. R. Soc. B* **281**, 20141486. (doi:10.1098/rspb.2014.1486)
- Gao K, Campbell DA. 2014 Photophysiological responses of marine diatoms to elevated CO<sub>2</sub> and decreased pH: a review. *Funct. Plant Biol.* **41**, 449–459. (doi:10.1071/FP13247)
- Gervais F, Riebesell U. 2001 Effect of phosphorus limitation on elemental composition and stable carbon isotope fractionation in a marine diatom growing under different CO<sub>2</sub> concentrations. *Limnol. Oceanogr.* **46**, 497–504. (doi:10.4319/lo.2001.46.3.0497)
- Tatters AO *et al.* 2013 Short- and long-term conditioning of a temperate marine diatom community to acidification and warming. *Phil. Trans. R. Soc. B* **368**, 20120437. (doi:10.1098/rspb.2012.0437)
- Kim J-M, Lee K, Shin K, Kang J-H, Lee H-W, Kim M, Jang P-G, Jang M-C. 2006 The effect of seawater CO<sub>2</sub> concentration on growth of a natural phytoplankton assemblage in a controlled mesocosm experiment. *Limnol. Oceanogr.* **51**, 1629–1636.