



THE UNIVERSITY *of* EDINBURGH

Edinburgh Research Explorer

Draft Genome Sequence of a Multidrug-Resistant *Acinetobacter baumannii* Strain from Chile

Citation for published version:

Opazo, A, Lopes, BS, García, P, Domínguez Yévenes, M, Lima, C, Bello-Toledo, H, González-Rocha, G & Amyes, SGB 2015, 'Draft Genome Sequence of a Multidrug-Resistant *Acinetobacter baumannii* Strain from Chile' *Genome announcements*, vol. 3, no. 4. DOI: 10.1128/genomeA.00687-15

Digital Object Identifier (DOI):

[10.1128/genomeA.00687-15](https://doi.org/10.1128/genomeA.00687-15)

Link:

[Link to publication record in Edinburgh Research Explorer](#)

Document Version:

Publisher's PDF, also known as Version of record

Published In:

Genome announcements

Publisher Rights Statement:

This is an open-access article distributed under the terms of the Creative Commons Attribution 3.0 Unported license.

General rights

Copyright for the publications made accessible via the Edinburgh Research Explorer is retained by the author(s) and / or other copyright owners and it is a condition of accessing these publications that users recognise and abide by the legal requirements associated with these rights.

Take down policy

The University of Edinburgh has made every reasonable effort to ensure that Edinburgh Research Explorer content complies with UK legislation. If you believe that the public display of this file breaches copyright please contact openaccess@ed.ac.uk providing details, and we will remove access to the work immediately and investigate your claim.



Draft Genome Sequence of a Multidrug-Resistant *Acinetobacter baumannii* Strain from Chile

Andrés Opazo,^{a,b} Bruno S. Lopes,^{a,c} Patricia García,^d Mariana Domínguez Yévenes,^b Celia Lima,^b Helia Bello-Toledo,^b Gerardo González-Rocha,^b Sebastian G. B. Amyes^a

Medical Microbiology, The Chancellor's Building, University of Edinburgh, Edinburgh, United Kingdom^a; Laboratorio de Investigación en Agentes Antibacterianos, Facultad de Ciencias Biológicas, Departamento de Microbiología, Universidad de Concepción, Concepción, Chile^b; Department of Microbiology, School of Medicine and Dentistry, University of Aberdeen, Aberdeen, United Kingdom^c; Laboratorio de Microbiología, Departamento de Laboratorios Clínicos, Escuela de Medicina, Pontificia Universidad Católica de Chile, Santiago, Chile^d

***Acinetobacter baumannii* strain Ab5 was isolated in the year 2007 in Chile, being one of the first multidrug-resistant (MDR) cases reported in the country. Here, we present the very first draft genome sequence of an MDR Chilean strain, which shows the presence of diverse resistance and acquired virulence genes.**

Received 20 May 2015 Accepted 20 May 2015 Published 2 July 2015

Citation Opazo A, Lopes BS, García P, Domínguez Yévenes M, Lima C, Bello-Toledo H, González-Rocha G, Amyes SGB. 2015. Draft genome sequence of a multidrug-resistant *Acinetobacter baumannii* strain from Chile. *Genome Announc* 3(4):e00687-15. doi:10.1128/genomeA.00687-15.

Copyright © 2015 Opazo et al. This is an open-access article distributed under the terms of the [Creative Commons Attribution 3.0 Unported license](https://creativecommons.org/licenses/by/4.0/).

Address correspondence to Andrés Opazo, andopazo@udec.cl.

Acinetobacter baumannii is a nosocomial pathogen normally classified as multidrug resistant (MDR). Alarming, the number of MDR *A. baumannii* isolates has increased in the last few years, making it a difficult microorganism to treat (1, 2), as infections caused by it are associated with high rates of mortality and long periods of hospitalization (3). The rates of *Acinetobacter* species MDR isolates in Latin America have increased lately, as resistance to imipenem in Chile increased from 0.0% during 1997 to 1999, to 47.4% in the 2008 to 2010 period (4), and 65.8% in 2014 (Antibiotic Resistant Group from Chilean Society for Infectious Diseases, personal communication).

In this work, we studied the genetic features of the *A. baumannii* MDR strain Ab5 through whole-genome sequencing (WGS), as this was one of the first sentinel cases of MDR isolates in Chile. To date, there are no whole-genome sequences of *A. baumannii* strains from Chile available in GenBank. The Ab5 strain was isolated at the Hospital of the Catholic University of Santiago in 2007 from a patient who developed a wound infection. Diverse antibiotics were tested by the disk diffusion method and by the agar dilution test, according to the Clinical and Standards Institute (CLSI) guidelines (5). Ab5 was found to be resistant to ceftazidime, imipenem, meropenem, gentamicin, cefotaxime, ciprofloxacin, cefepime, and piperacillin-tazobactam. The MICs to carbapenems were 8 µg/ml to meropenem and 16 µg/ml to imipenem.

WGS was performed utilizing the Illumina HiSeq 2000 platform. The data generated were assembled using the Velvet assembly pipeline (6), with contigs >200 bp, resulting in 107 contigs, a total length of 3,937,075 bp, and an N_{50} contig size of 173,003 bp; meanwhile, the average G+C content was 38.9%, coinciding with the data associated with the *Acinetobacter* genus.

Genome annotation was performed utilizing the Rapid Annotations using Subsystems Technology (RAST) (7), and the IGS Annotation Engine was used for structural and functional annotation (8). Manatee was used to view annotations (<http://manatee.sourceforge.net/>), while ResFinder (9), PathogenFinder (10), and

multilocus sequence typing (MLST) of total genome-sequenced bacteria (11) were utilized for complementary analyzes.

We identified 3,657 coding sequences (CDSs), 57 pseudogenes, 7 rRNAs, and 57 tRNAs. Additionally, Ab5 belonged to sequence type 162 (ST-162), which was previously detected in Brazil. We identified the *bla*_{OXA-58}, *bla*_{OXA-51}, *aph*(3')-Via, and the DNA topoisomerase type II genes that might explain its MDR phenotype. Interestingly, we found genes associated with iron acquisition, such as the *pitADC*, *piaABC*, and *piuABC* operons, which have been identified in *Streptococcus pneumoniae*. We also identified 25 pathogenesis-associated operons derived from *Mycobacterium* spp., whereas one detected operon, involved in internalin-like proteins synthesis, may have been imported from *Listeria*.

This first approach reflects the ability of *A. baumannii* to capture a wide range of genes derived from different species, making it a troublesome pathogen to treat in the nosocomial environment.

Nucleotide sequence accession numbers. This whole-genome shotgun project has been deposited at DDBJ/EMBL/GenBank under the accession no. [LANH00000000](https://www.ncbi.nlm.nih.gov/nuclink/LANH00000000). The version described in this paper is version LANH01000000.

ACKNOWLEDGMENTS

This work was supported through funds granted by the Chilean National Commission for Scientific and Technological Research (CONICYT) and by the National Fund for Scientific and Technological Development (FONDECYT) of Chile (project 3150286).

REFERENCES

- Evans BA, Hamouda A, Amyes SG. 2013. The rise of carbapenem-resistant *Acinetobacter baumannii*. *Curr Pharm Des* 19:223–238. <http://dx.doi.org/10.2174/1381612811306020223>.
- Peleg AY, Seifert H, Paterson DL. 2008. *Acinetobacter baumannii*: emergence of a successful pathogen. *Clin Microbiol Rev* 21:538–582. <http://dx.doi.org/10.1128/CMR.00058-07>.

3. Falagas ME, Bliziotis IA, Siempos II. 2006. Attributable mortality of *Acinetobacter baumannii* infections in critically ill patients: a systematic review of matched cohort and case-control studies. *Crit Care* 10:R48. <http://dx.doi.org/10.1186/cc4869>.
4. Gales AC, Castanheira M, Jones RN, Sader HS. 2012. Antimicrobial resistance among Gram-negative bacilli isolated from Latin America: results from SENTRY Antimicrobial Surveillance Program (Latin America, 2008–2010). *Diagn Microbiol Infect Dis* 73:354–360. <http://dx.doi.org/10.1016/j.diagmicrobio.2012.04.007>.
5. Clinical and Laboratory Standards Institute. 2014. Performance standards for antimicrobial susceptibility testing: 24th Informational Supplement, M100-S24. Clinical Laboratory Standards Institute, Wayne, PA.
6. Zerbino DR, Birney E. 2008. Velvet: algorithms for *de novo* short read assembly using de Bruijn graphs. *Genome Res* 18:821–829. <http://dx.doi.org/10.1101/gr.074492.107>.
7. Overbeek R, Olson R, Pusch GD, Olsen GJ, Davis JJ, Disz T, Edwards RA, Gerdes S, Parrello B, Shukla M, Vonstein V, Wattam AR, Xia F, Stevens R. 2014. The SEED and the rapid annotation of microbial genomes using subsystems technology (RAST). *Nucleic Acids Res* 42:D206–D214. <http://dx.doi.org/10.1093/nar/gkt1226>.
8. Galens K, Orvis J, Daugherty S, Creasy HH, Angiuoli S, White O, Wortman J, Mahurkar A, Giglio MG. 2011. The IGS standard operating procedure for automated prokaryotic annotation. *Stand Genomic Sci* 4:244–251. <http://dx.doi.org/10.4056/signs.1223234>.
9. Zankari E, Hasman H, Cosentino S, Vestergaard M, Rasmussen S, Lund O, Aarestrup FM, Larsen MV. 2012. Identification of acquired antimicrobial resistance genes. *J Antimicrob Chemother* 67:2640–2644. <http://dx.doi.org/10.1093/jac/dks261>.
10. Cosentino S, Voldby Larsen M, Møller Aarestrup F, Lund O. 2013. PathogenFinder—distinguishing friend from foe using bacterial whole genome sequence data. *PLoS One* 8:e77302. <http://dx.doi.org/10.1371/journal.pone.0077302>.
11. Larsen MV, Cosentino S, Rasmussen S, Friis C, Hasman H, Marvig RL, Jelsbak L, Sicheritz-Pontén T, Ussery DW, Aarestrup FM, Lund O. 2012. Multilocus sequence typing of total-genome-sequenced bacteria. *J Clin Microbiol* 50:1355–1361. <http://dx.doi.org/10.1128/JCM.06094-11>.