



# THE UNIVERSITY *of* EDINBURGH

## Edinburgh Research Explorer

### Repeated nebulisation of non-viral CFTR gene therapy in patients with cystic fibrosis

**Citation for published version:**

Alton, EFWF, Armstrong, DK, Ashby, D, Bayfield, KJ, Bilton, D, Bloomfield, EV, Boyd, AC, Brand, J, Buchan, R, Calcedo, R, Carvelli, P, Chan, M, Cheng, SH, Collie, DDS, Cunningham, S, Davidson, HE, Davies, G, Davies, JC, Davies, LA, Dewar, MH, Doherty, A, Donovan, J, Dwyer, NS, Elgmati, HI, Featherstone, RF, Gavino, J, Gea-sorli, S, Geddes, DM, Gibson, JSR, Gill, DR, Greening, AP, Griesenbach, U, Hansell, DM, Harman, K, Higgins, TE, Hodges, SL, Hyde, SC, Hyndman, L, Innes, JA, Jacob, J, Jones, N, Keogh, BF, Limberis, MP, Lloyd-evans, P, Maclean, AW, Manvell, MC, McCormick, D, MCGovern, M, Mclachlan, G, Meng, C, Montero, MA, Milligan, H, Moyce, LJ, Murray, GD, Nicholson, AG, Osadolor, T, Parra-leiton, J, Porteous, DJ, Pringle, IA, Punch, EK, Pytel, KM, Quittner, AL, Rivellini, G, Saunders, CJ, Scheule, RK, Sheard, S, Simmonds, NJ, Smith, K, Smith, SN, Soussi, N, Soussi, S, Spearing, EJ, Stevenson, BJ, Sumner-jones, SG, Turkhila, M, Ureta, RP, Waller, MD, Wasowicz, MY, Wilson, JM & Wolstenholme-hogg, P 2015, 'Repeated nebulisation of non-viral CFTR gene therapy in patients with cystic fibrosis: a randomised, double-blind, placebo-controlled, phase 2b trial' *The Lancet Respiratory Medicine*, vol. 3, no. 9, pp. 684-691. DOI: 10.1016/S2213-2600(15)00245-3

**Digital Object Identifier (DOI):**

[10.1016/S2213-2600\(15\)00245-3](https://doi.org/10.1016/S2213-2600(15)00245-3)

**Link:**

[Link to publication record in Edinburgh Research Explorer](#)

**Document Version:**

Publisher's PDF, also known as Version of record

**Published In:**

*The Lancet Respiratory Medicine*

**General rights**

Copyright for the publications made accessible via the Edinburgh Research Explorer is retained by the author(s) and / or other copyright owners and it is a condition of accessing these publications that users recognise and abide by the legal requirements associated with these rights.

**Take down policy**

The University of Edinburgh has made every reasonable effort to ensure that Edinburgh Research Explorer content complies with UK legislation. If you believe that the public display of this file breaches copyright please contact [openaccess@ed.ac.uk](mailto:openaccess@ed.ac.uk) providing details, and we will remove access to the work immediately and investigate your claim.





# Repeated nebulisation of non-viral CFTR gene therapy in patients with cystic fibrosis: a randomised, double-blind, placebo-controlled, phase 2b trial

Eric W F W Alton\*, David K Armstrong, Deborah Ashby, Katie J Bayfield, Diana Bilton, Emily V Bloomfield, A Christopher Boyd\*, June Brand, Ruairidh Buchan, Roberto Calcedo, Paula Carvelli, Mario Chan, Seng H Cheng, D David S Collie, Steve Cunningham\*, Heather E Davidson, Gwyneth Davies, Jane C Davies\*, Lee A Davies, Maria H Dewar, Ann Doherty, Jackie Donovan, Natalie S Dwyer, Hala I Elgmati, Rosanna F Featherstone, Jemyr Gavino, Sabrina Gea-Sorli, Duncan M Geddes, James S R Gibson, Deborah R Gill\*, Andrew P Greening, Uta Griesenbach\*, David M Hansell, Katharine Harman, Tracy E Higgins\*, Samantha L Hodges, Stephen C Hyde\*, Laura Hyndman, J Alastair Innes\*, Joseph Jacob, Nancy Jones, Brian F Keogh, Maria P Limberis, Paul Lloyd-Evans, Alan W Maclean, Michelle C Manvell, Dominique McCormick, Michael McGovern, Gerry McLachlan, Cuixiang Meng, M Angeles Montero, Hazel Milligan, Laura J Moyce, Gordon D Murray\*, Andrew G Nicholson, Tina Osadolor, Javier Parra-Leiton, David J Porteous\*, Ian A Pringle, Emma K Punch, Kamila M Pytel, Alexandra L Quittner, Gina Rivellini, Clare J Saunders, Ronald K Scheule, Sarah Sheard, Nicholas J Simmonds, Keith Smith, Stephen N Smith, Najwa Soussi, Samia Soussi, Emma J Spearing, Barbara J Stevenson, Stephanie G Sumner-Jones, Minna Turkhila, Rosa P Ureta, Michael D Waller, Marguerite Y Wasowicz, James M Wilson, Paul Wolstenholme-Hogg, on behalf of the UK Cystic Fibrosis Gene Therapy Consortium



## Summary

**Background** Lung delivery of plasmid DNA encoding the CFTR gene complexed with a cationic liposome is a potential treatment option for patients with cystic fibrosis. We aimed to assess the efficacy of non-viral CFTR gene therapy in patients with cystic fibrosis.

**Methods** We did this randomised, double-blind, placebo-controlled, phase 2b trial in two cystic fibrosis centres with patients recruited from 18 sites in the UK. Patients (aged  $\geq 12$  years) with a forced expiratory volume in 1 s (FEV<sub>1</sub>) of 50–90% predicted and any combination of CFTR mutations, were randomly assigned, via a computer-based randomisation system, to receive 5 mL of either nebulised pGM169/GL67A gene–liposome complex or 0.9% saline (placebo) every 28 days (plus or minus 5 days) for 1 year. Randomisation was stratified by % predicted FEV<sub>1</sub> (<70 vs  $\geq 70\%$ ), age (<18 vs  $\geq 18$  years), inclusion in the mechanistic substudy, and dosing site (London or Edinburgh). Participants and investigators were masked to treatment allocation. The primary endpoint was the relative change in % predicted FEV<sub>1</sub>. The primary analysis was per protocol. This trial is registered with ClinicalTrials.gov, number NCT01621867.

**Findings** Between June 12, 2012, and June 24, 2013, we randomly assigned 140 patients to receive placebo (n=62) or pGM169/GL67A (n=78), of whom 116 (83%) patients comprised the per-protocol population. We noted a significant, albeit modest, treatment effect in the pGM169/GL67A group versus placebo at 12 months' follow-up (3.7%, 95% CI 0.1–7.3; p=0.046). This outcome was associated with a stabilisation of lung function in the pGM169/GL67A group compared with a decline in the placebo group. We recorded no significant difference in treatment-attributable adverse events between groups.

**Interpretation** Monthly application of the pGM169/GL67A gene therapy formulation was associated with a significant, albeit modest, benefit in FEV<sub>1</sub> compared with placebo at 1 year, indicating a stabilisation of lung function in the treatment group. Further improvements in efficacy and consistency of response to the current formulation are needed before gene therapy is suitable for clinical care; however, our findings should also encourage the rapid introduction of more potent gene transfer vectors into early phase trials.

**Funding** Medical Research Council/National Institute for Health Research Efficacy and Mechanism Evaluation Programme.

**Copyright** © Alton et al. Open Access article distributed under the terms of CC BY.

## Introduction

Cystic fibrosis has been a target for gene therapy since the CFTR gene was cloned in 1989.<sup>1</sup> Lung disease is the main cause of morbidity and mortality in individuals with cystic fibrosis, with a median age at death of 29 years (95% CI 27–31).<sup>2</sup> Early expectations of a rapid breakthrough were

based on supposed ease of access to the target respiratory epithelium via inhaled aerosols. These hopes were tempered by the subsequent realisation that the airways are well defended, in keeping with their predominant function as conducting passages, rather than absorptive surfaces.

*Lancet Respir Med* 2015

Published Online  
July 3, 2015  
[http://dx.doi.org/10.1016/S2213-2600\(15\)00245-3](http://dx.doi.org/10.1016/S2213-2600(15)00245-3)

\*Senior authors

Royal Brompton and Harefield NHS Foundation Trust, London, UK (Prof D Bilton FRCP, J Donovan PhD, Prof D M Geddes FRCP, Prof D M Hansell FRCP, B F Keogh FRCA, M A Montero MD, Prof A G Nicholson FRCPATH, T Osadolor MSc, N J Simmonds FRCP, J Jacob FRCP, N Jones BPharm, S Sheard FRCP); Imperial College London, London, UK (Prof E W F W Alton FMedSci, Prof D Ashby CStat, K J Bayfield BSc, E V Bloomfield BSc, P Carvelli BN, M Chan MSc, G Davies MBChB, Prof J C Davies FRCPCH, N S Dwyer RN, R F Featherstone BSc, S Gea-Sorli PhD, Prof U Griesenbach PhD, K Harman MRCPH, T E Higgins BSc, S L Hodges BSc, M C Manvell BSc, C Meng BSc, E K Punch BSc, K M Pytel PhD, G Rivellini, C J Saunders BSc, S N Smith PhD, N Soussi BSc, S Soussi BSc, E J Spearing MSc, M Turkhila BSc, M D Waller MRCP, M Y Wasowicz PhD, R P Ureta BN, J Gavino RN); Western General Hospital, Edinburgh, UK (R Buchan MPharm, M H Dewar BSc, H I Elgmati MRCP, Prof A P Greening FRCP, E)

J A Innes FRCPE,  
M McGovern MRCP,  
H Milligan BSc; Usher Institute  
of Population Health Sciences  
and Informatics and Centre for  
Population Health Sciences  
(Prof G D Murray PhD) and The  
Roslin Institute and R(D)SVS  
(D D S Collie MRCVS,  
G McLachlan PhD) and Centre  
for Genomic and Experimental  
Medicine, Institute of Genetics  
and Molecular Medicine  
(A C Boyd PhD, J Brand BSc,  
H E Davidson, A Doherty  
BSc, J S R Gibson BSc,  
L Hyndman MSc,  
A W Maclean BSc, J Parra-Leiton,  
Prof D J Porteous PhD,  
B J Stevenson BSc), University of  
Edinburgh, Edinburgh, UK;  
Royal Hospital for Sick  
Children, Edinburgh, UK  
(D K Armstrong MB ChB,  
S Cunningham MB ChB);  
Nuffield Division of Clinical  
Laboratory Sciences, Radcliffe  
Department of Medicine,  
University of Oxford, Oxford,  
UK (L A Davies DPhil,  
D McCormick PhD, D R Gill PhD,  
S C Hyde DPhil, L J Moyce BSc,  
I A Pringle DPhil,  
S G Sumner-Jones PhD); Gene  
Therapy Program, Department  
of Pathology and Laboratory  
Medicine, University of  
Pennsylvania, Philadelphia, PA,  
USA (R Calcedo PhD,  
M P Limberis PhD,  
Prof J M Wilson MD); Genzyme,  
a Sanofi Company,  
Framingham, MA, USA  
(S H Cheng PhD, R K Scheule PhD,  
P Wolstenholme-Hogg PhD);  
NHS Blood and Transplant,  
Bristol, UK (P Lloyd-Evans PhD,  
K Smith BSc); University of  
Miami, Miami, FL, USA  
(Prof A L Quittner PhD)

Correspondence to:  
Prof Eric Alton, Department of  
Gene Therapy, National Heart  
and Lung Institute, London  
SW3 6LR, UK  
e.alton@imperial.ac.uk

For the UK Cystic Fibrosis Gene  
Therapy Consortium see <http://www.cfgenetherapy.org.uk>

For the protocol see [http://www.cfgenetherapy.org.uk/assets/fileGallery/MD\\_Clinical\\_Trial\\_Protocol.pdf](http://www.cfgenetherapy.org.uk/assets/fileGallery/MD_Clinical_Trial_Protocol.pdf)

## Research in context

### Evidence before this study

We searched PubMed between June 1, 1992, and March 1, 2015, for studies published that included the terms "non-viral, gene therapy, cystic fibrosis" or "liposome, gene therapy, cystic fibrosis".

### Added value of this study

We report the first trial of non-viral *CFTR* gene therapy for patients with cystic fibrosis that is powered to detect clinically relevant pulmonary changes. Our study has progressed this field of research from phase 1 and 2a studies showing changes in molecular surrogates of *CFTR* function, to a phase 2b setting

assessing changes in lung function in patients with a broad range of *CFTR* mutations. Additionally, our study shows that monthly repeated application of non-viral gene therapy can be safely administered to the lungs over a 1 year period.

### Implications of all the evidence

By providing the first proof of concept that non-viral gene therapy can beneficially affect lung function, follow-up studies can assess optimum dose, dosing interval, and patient stratification at trial entry. Our findings are likely to catalyse earlier translation of more efficient vectors into first-in-man trials.

Various vectors for delivery of the *CFTR* gene into respiratory epithelial cells have been assessed. Viral approaches, including adenoviruses, adeno-associated viruses, and retroviruses, have faltered because of inefficient transduction from the luminal surface and immune responses restricting the efficacy of repeated application.<sup>3</sup> As such, research from the UK Cystic Fibrosis Gene Therapy Consortium has initially focused on non-viral vectors. Formulation and delivery of plasmid DNA-liposome complexes have been refined in a large series of preclinical studies,<sup>4,5</sup> and safety,<sup>6,7</sup> molecular efficacy, and practical doses have been assessed in several phase 1 and 2a studies in patients with cystic fibrosis.<sup>1,3</sup> We did this study to assess the clinical efficacy of the non-viral *CFTR* gene-liposome complex pGM169/GL67A<sup>8</sup> after repeated delivery to the airways.

## Methods

### Study design and participants

We did this randomised, double-blind, placebo-controlled, phase 2b trial in two cystic fibrosis centres with patients recruited from 18 sites in the UK. Eligible participants had diagnosed cystic fibrosis, were aged 12 years or older, had a forced expiratory volume in 1 s (FEV<sub>1</sub>) of 50–90% predicted, and had any combination of *CFTR* mutations.

The protocol was approved by the National Research Ethics Committee and the local Research Committees at the two dosing sites and the 16 other referral centres. Each patient, or a parent, provided written informed consent, and children provided assent.

### Randomisation and masking

We randomly assigned patients (1:1), via a computer-based randomisation system, to receive nebulised pGM169/GL67A or 0.9% saline (placebo). Randomisation was stratified by % predicted FEV<sub>1</sub> (<70 vs ≥70%), age (<18 vs ≥18 years), inclusion in the mechanistic substudy, and dosing site (London or Edinburgh). Participants in the mechanistic substudy were randomly assigned (2:1) to receive nebulised pGM169/GL67A or placebo, and could participate as part of either a nasal or bronchoscopy group, or both. Participants and investigators were masked to

treatment allocation, with the randomisation code known only by pharmacy staff at the two dosing sites.

### Procedures

Patients received 5 mL of either 0.9% saline or pGM169/GL67A complex nebulised through a Trudell AeroEclipse II device (Trudell Medical International, London, ON, Canada) at 28 day intervals (plus or minus 5 days) for 12 months. Each 5 mL dose of pGM169/GL67A contained 13.3 mg of plasmid DNA and 75 mg of the GL67A lipid mixture. Routine treatments were continued throughout the study, except for DNase, which was withheld for 24 h before and after dosing. In addition to the nebulised dose, patients in the nasal group of the mechanistic substudy received 2 mL of placebo or pGM169/GL67A divided between nasal cavities via a nasal spray device at the time of each lung dose. Patients in the bronchoscopy group followed the standard protocol, but also underwent a bronchoscopy under general anaesthesia before the first dose and 28 days (plus or minus 5 days) after the final dose.

### Outcomes

The primary efficacy endpoint was the relative change in % predicted FEV<sub>1</sub>, calculated from the mean of two baseline values (at screening and before dosing on day of the first dose) to the mean of two values (2 and 4 weeks after last dose) at study completion. Secondary outcomes included additional measurements of lung function, CT scans, and Cystic Fibrosis Questionnaire-Revised (CFQ-R) scores.<sup>9</sup> Exploratory endpoints included exercise testing, activity monitoring, and sputum inflammatory markers. Mechanistic endpoints were nasal or bronchial vector-specific DNA, mRNA, and electrophysiological assessment of *CFTR* function. We did extensive safety assessments.

### Statistical analysis

The statistical analyses were prespecified in a statistical analysis plan. With use of pilot data, we estimated the standard deviation of the relative change in % predicted FEV<sub>1</sub> in the target cystic fibrosis population to be 10% over 12 months. A total sample size of 120 assessable

patients would provide 90% power to detect a 6% difference between groups in the mean change from baseline at a two-sided 5% significance level. This power calculation was conservative because covariate adjustment can be expected to increase statistical power. We did analyses in the per-protocol population (primary analysis), predefined as participants who received at least nine doses of pGM169/GL67A or placebo, and in the intention-to-treat population, who received at least one dose of pGM169/GL67A or placebo.

We compared outcomes between groups with an ANCOVA model, with inclusion of the relevant baseline value, treatment allocation, and stratification factors (baseline predicted FEV<sub>1</sub>, age, dosing site, inclusion in substudy). Results are reported as adjusted mean differences with corresponding 95% CIs. We assessed subgroup effects by including the relevant interaction term in the ANCOVA model. To allow results from different endpoints to be plotted on a common scale, the estimated treatment effects were standardised and presented as multiples of the underlying SD. No adjustment was made to the p values to allow for multiplicity because the secondary endpoints were supportive and the corresponding p values were interpreted conservatively. We assessed bronchial and nasal biomarkers with a Mann–Whitney *U* test. A two-sided p value less than 0.05 was considered statistically significant.

The trial was overseen by an independent Data Monitoring and Ethics Committee and a Trial Steering Committee. This trial is registered with ClinicalTrials.gov, number NCT01621867.


### Role of the funding source

The funder of the study had no role in study design, data collection, data analysis, data interpretation, or writing of the report. The corresponding author had full access to all the data in the study and had final responsibility for the decision to submit for publication.

### Results

Figure 1 shows the trial profile. Between June 12, 2012, and June 24, 2013, we randomly assigned 140 patients to receive placebo (n=62) or pGM169/GL67A (n=78), of whom 136 (97%) patients comprised the intention-to-treat population and 116 (83%) patients comprised the per-protocol population (figure 1). Reasons for discontinuation in the intention-to-treat population were similar between groups (appendix). Baseline characteristics were similar between the two groups (table 1). Unless indicated otherwise, all subsequent details relate to the per-protocol population.

114 (98%) patients had paired pre-treatment and post-treatment measurements of % predicted FEV<sub>1</sub>. Of the two patients (both in the placebo group) who did not have paired measurements, one patient could not do the test because of a surgery-related pneumothorax and one withdrew because of time commitments and was



**Figure 1: Trial profile**

Numbers of patients in the intention-to-treat population are unequal because of the 2:1 allocation in the mechanistic substudy. FEV<sub>1</sub>=forced expiratory volume in 1 s.

unavailable for follow-up measurements. We recorded a significant ANCOVA-adjusted treatment effect in the pGM169/GL67A group versus placebo at 12 months' follow-up (3.7%, 95% CI 0.1–7.3; p=0.046; figure 2). The relative changes within each of the individual groups were –4.0% (95% CI –6.6 to –1.4) in the placebo group and –0.4% (–2.8 to 2.1) in the pGM169/GL67A group (figure 2). Post-hoc analysis showed that 21 (18%) patients (n=6 in the placebo group and n=15 in the pGM169/GL67A group) had an improvement in % predicted FEV<sub>1</sub> of 5% or more of their individual baseline values. For comparison, the treatment effect in patients in the intention-to-treat

See Online for appendix


	Placebo group (n=54)	pGM169/GL67A group (n=62)
Age (years)	26.0 (13.0)	23.6 (10.8)
<18 years old	17 (31%)	23 (37%)
≥18 years old	37 (69%)	39 (63%)
Sex		
Female	25 (46%)	31 (50%)
Male	29 (54%)	31 (50%)
Centre distribution number		
Edinburgh	24 (44%)	22 (35%)
London	30 (56%)	40 (65%)
Height (cm)	165.0 (10.6)	163.6 (10.9)
Weight (kg)	61.6 (15.6)	61.0 (15.7)
FEV <sub>1</sub> (% predicted)	69.0 (9.9)	69.9 (11.1)
Body-mass index (kg/m <sup>2</sup> )	22.4 (4.4)	22.4 (4.5)
Mutation class		
Phe508del/Phe508del	26 (48%)	31 (50%)
Phe508del/class 1–6	22 (41%)	23 (37%)
Not Phe508del/class 1	1 (2%)	3 (5%)
Heterozygous/homozygous class 3–6	2 (4%)	2 (3%)
Phe508del/unknown class	3 (6%)	3 (5%)

Data are mean (SD) or n (%), unless otherwise indicated.

**Table 1: Baseline and demographic characteristics**

population who had spirometry measurements both before dosing and within the protocol-defined window after their final dose (n=56 in the placebo group and n=65 in the pGM169/GL67A group) was 3.6% (95% CI 0.2–7.0; p=0.039), with the 20 patients included in the intention-to-treat, but not per-protocol, analysis, receiving a mean of 3.7 doses (SD 1.9).

Figure 3 summarises changes in a range of secondary outcomes. The treatment effect was significant for FVC (p=0.031; appendix) and CT gas trapping (p=0.048), but not for other measures of lung function, imaging, and quality of life (figure 3). We assessed whether a responder subgroup could be identified; the appendix summarises the prespecified subgroups. We noted no significant differences in the primary outcome treatment effect with respect to sex, age, CFTR mutation (phe508del homozygous vs other), *Pseudomonas* colonisation, predominant smaller or larger airway disease on CT at presentation, concurrent drugs, or treatment-associated adverse events (appendix). Although some subgroups had larger treatment effects than others, these results were typically due to a greater decline in FEV<sub>1</sub> in the placebo group, rather than to any difference of effect in the pGM169/GL67A group (appendix). Stratification by baseline % predicted FEV<sub>1</sub> suggested a difference, albeit non-significant, in treatment effect between patients with more severe disease (FEV<sub>1</sub> 49.6–69.2% predicted), who had a treatment effect of 6.4% (95% CI 0.8–12.1), and those with less severe disease (69.6–89.9% predicted), who had a treatment effect of 0.2% (–4.6 to 4.9; p<sub>interaction</sub>=0.065; appendix). In patients with more severe




**Figure 2: Timecourse of the primary outcome response to either placebo or pGM169/GL67A (A) and the individual patient responses in the pGM169/GL67A (B) and placebo (C) groups**

Error bars in panel A show the standard error of the mean. Primary outcome measurements were taken at each treatment visit before administration of study drugs. Pre and post values indicate the mean of two measurements at the respective timepoints. Positive values in panels B and C show an improvement. FEV<sub>1</sub>=forced expiratory volume in 1 s.

disease, post-trial and pre-trial changes in both the placebo group (–4.9%) and the pGM169/GL67A group (1.5%) contributed to the treatment effect. Secondary outcomes showed a similar trend favouring the more severe category (appendix).

Patients in both treatment groups received a median of three (IQR one to five) courses of oral or intravenous antibiotics during the trial. Specifically, we assessed co-administered antibiotics during the critical analysis





**Figure 3: Forest plot showing secondary outcome responses to placebo or pGM169/GL67A**

Data are mean (SD) or mean (95% CI), unless otherwise indicated. The size of the circles is proportional to the number of patients represented and the error bars show 95% CIs. Values shown for FEV<sub>1</sub> are the relative change in the % predicted FEV<sub>1</sub>. To allow results from different endpoints to be plotted on a common scale, the estimated treatment effects were standardised to be presented as multiples of the underlying SD (standardised treatment effect). FEV<sub>1</sub>=forced expiratory volume in 1 s. MEF25-75=mid-expiratory flow between 25% and 75% of FVC. KCOc=diffusion capacity of the alveolar capillary membrane, corrected for haemoglobin concentrations. TLCoc=transfer factor of the lung for carbon monoxide, corrected for haemoglobin concentrations. \*Refers to scores from the Cystic Fibrosis Questionnaire-Revised.

period from dose 11 to the end of the trial. Numbers of patients receiving any additional antibiotics were 26 (48%) in the placebo group and 30 (51%) in the pGM169/GL67A group ( $\chi^2$   $p=0.774$ ). Thus, the observed FEV<sub>1</sub> treatment effect was considered to be independent of concurrent antibiotic courses.

No clinically relevant pattern of changes could be distinguished in the exploratory outcomes of activity and exercise monitoring and serum and sputum inflammatory markers (appendix). In the bronchoscopy group of the substudy, vector-specific DNA increased in 12 (86%) of 14 patients in the pGM169/GL67A group and was below the limit of quantification in all ( $n=7$ ) placebo samples ( $p=0.001$ ; figure 4A); vector-specific mRNA was below the level of sensitivity in both groups (appendix). Changes in basal post-trial and pre-trial potential difference values did not differ significantly in either group (appendix). Figure 4B shows bronchial chloride responses using the mean of all interpretable tracings

for each patient; a negative value indicates a change in the non-cystic fibrosis direction. Patients in the placebo group ( $n=7$ ) had a median change (post-trial minus pre-trial) of 3.1 mV (range 9.3 to -1.2) and those in the pGM169/GL67A group ( $n=10$ ) had a change of -1.3 mV (4.0 to -5.8;  $p=0.032$ ; figure 4B). Five (50%) of ten patients in the pGM169/GL67A group had values that were more negative than the largest response in the placebo group (figure 4). In the same analysis with only the most negative value recorded for each patient at any timepoint, patients in the placebo group had a median post-trial minus pre-trial change of 2.6 mV (range 9.3 to -1.2) and those in the pGM169/GL67A group had a change of -2.8 mV (4.0 to -16.8 mV;  $p=0.088$ ; figure 4C). Six (60%) patients in the pGM169/GL67A group had values that were more negative than the largest response in the placebo group (figure 4). The appendix shows absolute bronchial potential difference values.

In patients in the nasal group of the substudy, vector-specific DNA increased in all the 17 patients given pGM169/GL67A. Despite apparent pGM169 contamination in some samples, the change in pGM169 concentrations differed significantly between the groups (appendix); no vector-specific mRNA was quantifiable in either group. We noted no significant changes in the baseline, zero chloride, or isoprenaline responses (appendix). Four (29%) of 14 pGM169/GL67A patients had mean post-trial minus pre-trial treatment responses (ranging from  $-3.4$  mV to  $-7.0$  mV) that were more negative than the largest response in the placebo group ( $n=6$ ; appendix). The appendix shows absolute nasal potential difference values.


All patients had adverse events, with no significant difference between groups for either total events or within the nine predefined adverse event categories (table 2). One patient in the placebo group and one

patient in the pGM169/GL67A group discontinued study treatment because of adverse events (fatigue and increased respiratory symptoms and flu-like symptoms, respectively). We recorded six serious adverse events, all in the pGM169/GL67A group (appendix). Neither the Data Monitoring and Ethics Committee nor the Trial Steering Committee regarded any serious adverse event as related to study drug; however, one event was considered to be possibly related to a trial procedure (bronchoscopy). We noted no clinically relevant changes in haematology, biochemistry, conversion of anti-CFTR T cells, anti-DNA antibodies, histology, or lipid staining (appendix) and no patients died during the study.

**Discussion**

We report the first trial of non-viral based gene therapy for cystic fibrosis, powered to detect clinically relevant pulmonary changes. After monthly dosing for 1 year, we recorded evidence of a beneficial effect of gene therapy versus placebo on FEV<sub>1</sub>. No effect of sex, age, or whether patients were homozygous for the most common F508del CFTR mutation could be detected. No clinically important adverse events attributable to treatment with pGM169/GL67A were reported.

Although these findings are encouraging, they should be put into perspective. We noted a stabilisation of FEV<sub>1</sub> in the pGM169/GL67A group rather than an improvement. This stabilisation took place over a 1 year period and further work will be needed to see if this effect is maintained. The reduction in FEV<sub>1</sub> in the placebo group was within the range reported in some other prospective trials<sup>10-12</sup> and is consistent with a median survival of 29 years, but is greater than would be expected from registry data.<sup>2</sup> Three factors are likely to have influenced this difference. First, the requirement for clinical stability at trial entry meant that patients might have been at their optimum respiratory health at this stage. Second, the enthusiasm of patients to enter the trial, accompanied by a focus on self-care, might have resulted in short-term improvements in lung function during the recruitment period. Both factors are likely to lead to a subsequent decline in lung function as patients regress to their mean values. Third, we included all available data, whether from stable patients or those with exacerbations, by contrast with registry data, which focuses on measurements obtained at annual review. Stabilisation of lung disease in itself is a worthwhile aim and we would caution against the bar being set too high for novel therapeutics in cystic fibrosis populations with an unselected range of mutations. The large response to ivacaftor in patients with class III mutations takes place in the context of correctly localised CFTR protein. By contrast, much smaller improvements in lung function were shown in the ivacaftor-lumacaftor trial for the most common mutation (phe508del) in which the CFTR protein is misfolded and mislocalised.<sup>13</sup>



**Figure 4:** Assessment of DNA from bronchial brushings in the placebo ( $n=7$ ) and pGM169/GL67A ( $n=14$ ) subgroups (A) and the response of the bronchial epithelium to perfusion with a zero chloride solution containing isoprenaline  $10 \mu\text{M}$  (B, C)

Horizontal bars show median values. Each circle in panel A represents an individual patient. Each symbol in panels B and C shows the change in response from trial start to finish for the relevant treatment in an individual patient. Of the 16 participants in the bronchoscopy subgroup, 15 individuals had post-dose bronchoscopies, of whom 14 individuals generated samples for DNA and mRNA molecular analysis. The plotted value in panel B is the mean of all interpretable recordings (range 1-3), and in panel C is the most negative value obtained from all interpretable recordings, at each timepoint for that patient. A more negative value is in the non-cystic fibrosis direction. LOQ=limit of quantification, PBNQ=positive but not quantifiable.

	Placebo group (n=54)	pGM169/GL67A group (n=62)
Lower airway respiratory symptoms	7.9	9.0
Gastrointestinal symptoms	2.1	1.8
Fever or flu-like symptoms	1.1	1.4
Headache	1.2	1.1
Upper airway symptoms	2.3	3.4
Elevated liver function tests	0.3	0.4
Haematuria	0.2	0.2
Isolated raised inflammatory markers	0.8	0.7
Other	3.2	3.3
Total	19.1	21.2

Data are mean number of times the respective symptom was experienced by each patient during the trial. Values were calculated by dividing the total number of the relevant adverse event by the total number of relevant patients in that group.

**Table 2:** Adverse events

The response in our study was heterogeneous, with apparent responders and non-responders. The data suggest that an approximate doubling of treatment effect was achieved in patients with more severe disease stratified by baseline FEV<sub>1</sub>, supported by trends in other clinically relevant secondary measures. A larger trial with a stratified trial entry design, powered to assess subgroups, and that addresses the mechanisms of response heterogeneity, will be important to verify or refute these data. This differential response could relate to the dose deposited in the airways; in patients with lower baseline FEV<sub>1</sub> the relatively more obstructed smaller airways result in a larger proportion of the 5 mL dose being deposited in the larger airways. In pre-trial studies we assessed airway deposition in patients with cystic fibrosis with varying FEV<sub>1</sub> severity with technetium-99m labelled human serum albumin of similar droplet size (3–4 µm, using a different nebuliser system) to the pGM169/GL67A formulation. Bronchial airway (generations 2–8) fractional deposition was 2.9% of delivered dose (standard error of the mean [SEM] 0.2; n=33) in patients with 70–90% predicted FEV<sub>1</sub> and roughly twice as great (6.0%, SEM 1.0; n=23) in those with 50–70% predicted FEV<sub>1</sub>. An additional contributory factor to this enhanced efficacy might be the increased mitotic rate of more severely affected tissues,<sup>14</sup> which decreases the proportion of time that the nuclear membrane is intact, the membrane acting as a barrier to plasmid DNA entry to the nucleus.

We cannot rule out that the changes recorded in the present study are the result of a non-specific response to the pGM169/GL67A formulation. The placebo was 0.9% saline rather than a scrambled or CFTR-deleted plasmid-liposome complex. We selected 0.9% saline partly on the basis of pragmatic financial considerations, but mainly for ethical considerations, not wishing to expose patients with cystic fibrosis to first-in-man repeated pulmonary dosing of an untested product that might direct the expression of an immunologically active peptide or novel non-coding RNA molecule with deleterious biological functions. Furthermore, we wanted to compare progression on therapy with the natural history of the disease. In terms of alternative explanations for the effects we noted, we know of no evidence that monthly nebulisation of 0.9% saline is deleterious to lung function, nor that liposome alone produces physiological improvements in either patients without,<sup>15</sup> or those with<sup>16</sup> cystic fibrosis. Delivery of non-CFTR encoding plasmid DNAs to the human airways has not been associated with a gain in CFTR chloride-channel function, nor improvement in any cystic fibrosis-related assay,<sup>17,18</sup> and plasmid DNA is generally associated with pro-inflammatory, rather than non-specific, beneficial effects.<sup>19</sup> We did not identify any pathophysiological changes in the airways, such as inflammation or remodelling, nor any changes in bacterial species that might otherwise explain the outcomes. Nevertheless, we cannot exclude that

DNA-liposome complexes augment host defences, stimulate mucus clearance, or enhance bacterial killing to an extent undetectable on semi-quantitative routine culture.

Results showing more robust changes in molecular CFTR surrogates would have been reassuring. Despite extensive optimisation of quantitative realtime-PCR assays, the pGM169-derived mRNA assay has poor sensitivity and is adversely affected by the inclusion of high levels of total RNA or modest concentrations of pGM169 plasmid DNA. In ovine studies we have shown that a 20 mL nebulised dose of pGM169/GL67A, four times that used in the present trial, is the lower threshold for reproducible detection of mRNA with this assay in airway tissue samples (unpublished).<sup>6</sup> Thus, our inability to detect pGM169-derived mRNA after delivery of 5 mL of pGM169/GL67A to the human airways, although disappointing, was not surprising. In human tissues, we have noted the low sensitivity of assays assessing vector-specific mRNA from human samples in vivo,<sup>16,20,21</sup> and have noted the greater sensitivity of detection of electrophysiological changes, consistent with findings in this study.<sup>17,18,22</sup>

The ratio of area sampled to area dosed is small. Although we recorded significant chloride secretory changes in the bronchial, but not the nasal, epithelium, we caution against placing undue weight on either observation. The size of the groups in the mechanistic substudy was limited by both the practicality of the procedures and the acceptability to patients of the additional invasive tests, leading to low statistical power for these measures. We would instead conclude that modest variable changes can be shown with currently available assays that remain insufficiently sensitive to detect changes in low levels of CFTR function when assessed in vivo in humans; further optimisation in these or other assays is needed.

Although we are encouraged by the first demonstration of a significant beneficial effect in lung function compared with placebo associated with gene therapy in patients with cystic fibrosis, the mean difference was modest, only recorded in some individuals, and at the lower end of the range of results seen in clinical trials which result in changes in patient-related care.<sup>23,24</sup> We did not formally assess infective exacerbations in view of the fairly small patient numbers in our study, but use of antibiotic courses as a surrogate identified no obvious treatment advantage. The treatment effect is consistent with a clinically meaningful benefit from the perspective of the European Medicine Agency;<sup>25</sup> however, further improvements in efficacy and consistency of response to the current formulation, or its combination with CFTR potentiators, are needed before gene therapy is suitable for clinical practice. Furthermore, our findings should encourage the rapid introduction of more potent gene transfer vectors into early phase trials, now that much of the groundwork has been established.



The data reported here provide the first proof of concept that repeated administration of non-viral CFTR gene therapy can safely change clinically relevant parameters, providing another step along the path of translational cystic fibrosis gene therapy.

#### Contributors

EWFWA, ACB, SC, JCD, DMG, DRG, APG, UG, SCH, TEH, JAI, GDM, and DJP conceived, designed, and analysed the overall study. DA and GDM designed and coordinated data collection and statistical analysis. DKA, KJB, DB, PC, GD, NSD, HIE, RFF, JG, JSRG, DMH, KH, SLH, JJ, BFK, MM, EKP, ALQ, CJS, SSh, NJS, NS, EJS, SNS, RPU, and MDW assessed patient outcomes and undertook and analysed individual in-vivo assays. EVB, MHD, and SSo coordinated and undertook the administration of the trial. RB, NJ, PL-E, GR, and KS oversaw receipt, preparation, and dispensing of study drug. JB, RC, MC, HED, AD, JD, SG-S, LH, MPL, AWM, MCM, DM, CM, MAM, HM, LJM, AGN, TO, JP-L, IAP, KMP, BJS, SGS-J, MT, MYW, and JMW designed, undertook, and analysed in-vitro assays. SHC, RKS, and PW-H coordinated the production of lipid 67A. DDSC, LAD, and GM designed, undertook, and analysed studies of study drug delivery.

#### Declaration of interests

ACB, AD, APG, AGN, AWM, BFK, BJS, CM, CJS, DKA, DA, DB, DM, DMH, DMG, DDSC, DJP, DRG, EKP, EJS, EVB, EWFWA, GD, GM, GDM, GR, HED, HM, HIE, IAP, JAI, JB, JCD, JD, JG, JSRG, JJ, JP-L, JMW, KJB, KH, KMP, KS, LAD, LH, LJM, MC, MCM, MHD, MM, MT, MYW, MAM, MDW, MPL, NSD, NJ, NJS, NS, PC, PL-E, PW-H, RB, RC, RFF, RPU, SC, SCH, SG-S, SLH, SSh, SSo, SGS-J, SNS, TEH, TO, and UG report grants from the National Institute for Health Research, the Cystic Fibrosis Trust, Just Gene Therapy, Medicor Foundation, and other support from Genzyme, a Sanofi company, related to the submitted work during the conduct of the study. ALQ, JCD, JMW, MCM, NJS, RKS, SC, and SHC report fees, grants, honoraria, and patents outside of the submitted work. ACB, DJP, DRG, EWFWA, JAI, JCD, LAD, SCH, and UG report patents related to the submitted work.

#### Acknowledgments

We thank the patients, carers, and families who gave so much to this trial; those who so generously supported the programme of work that led to this trial by giving to the Cystic Fibrosis Trust; the Medicor Foundation and the Cystic Fibrosis Trust for part-funding the cost of the plasmid DNA and lipid; the National Institute for Health Research (NIHR) Clinical Research Network and Just Gene Therapy for additional funding; Sandra Griffiths, Nayan Das, Amanda Bravery, Juan Gonzales-Maffe, and Ginny Picot (Imperial Clinical Trials Unit); Jermaine Wright, Fabricio Ghiraldi, Stephen Man, Judith Foy, and Viba Teli (Royal Brompton Hospital Pharmacy) and Peter Brown (Royal Marsden Pharmacy); Leticia Brown, Michelle Watson-Dotchin, and Michelle Pugh for administrative support; the Efficacy and Mechanism Evaluation Programme team, including Lucy Knight and Dani Preedy, who guided us throughout the trial; Tony Hickson and Gursharan Randhawa for help with the intellectual property and commercialisation aspects; the Data Monitoring and Ethics Committee (Colin Wallace, Caroline Elston, George Dickson, Julie Morris) and the Trial Steering Committee (Ashley Woodcock, Nikki Samsa, Brandon Wainwright, Pierre Lehn, Janet Allen) for their wise and helpful advice; and the teams at the referring sites who gave so generously of their time: Jeremy Hull (Oxford Churchill); Nick Bell and Kathryn Bateman (Bristol); Judy Ryan (Papworth); Maya Desai, Michelle Taberner, and Edward Nash (Birmingham); Paul Aurora and Ammani Prasad (Great Ormond Street); Judith Duguid, Alexandra Highton, and Timothy Ho (Frimley); Siobhan Carr and Catherine Lambert (Royal London); Thomas Daniels and Victoria Brown (Southampton); Daniel Peckham and Amy Drifill (Leeds); Felicity Perrin and Mike Waller (King's College London); Gordon MacGregor, Steve Bicknell, Ewan Ross, and Anne Devenny (Glasgow); Carol Dryden (Wishaw); Richard Brooker, Graham Devereux, Owen Dempsey (Aberdeen); Helen Rodgers and Jonathon McCormick (Dundee); and Steven Bourke and David Spencer (Dundee). This study was funded by the Efficacy and Mechanism Evaluation (EME) Programme, a Medical Research Council and NIHR

partnership. The research was supported by the NIHR Respiratory Biomedical Research Unit at the Royal Brompton and Harefield NHS Foundation Trust, and the NIHR Imperial Biomedical Research Centre at Imperial College Healthcare NHS Trust and Imperial College London. The UK Cystic Fibrosis Gene Therapy Consortium would not have come together, and this trial would not have happened, without the vision and leadership of Rosie Barnes during her tenure as Chief Executive of the Cystic Fibrosis Trust.

#### References

- Griesenbach U, Alton EW. Moving forward: cystic fibrosis gene therapy. *Hum Mol Genet* 2013; **22**: R52–58.
- Cystic Fibrosis Trust. UK Cystic Fibrosis Registry annual data report 2013. 2014. <http://www.cysticfibrosis.org.uk/media/598466/annual-data-report-2013-jul14.pdf> (accessed April 26, 2015).
- Gill DR, Hyde SC. Delivery of genes into the CF airway. *Thorax* 2014; **69**: 962–64.
- Davies LA, Nunez-Alonso GA, McLachlan G, Hyde SC, Gill DR. Aerosol delivery of DNA/liposomes to the lung for cystic fibrosis gene therapy. *Hum Gene Ther Clin Dev* 2014; **25**: 97–107.
- McLachlan G, Davidson H, Holder E, et al. Pre-clinical evaluation of three non-viral gene transfer agents for cystic fibrosis after aerosol delivery to the ovine lung. *Gene Ther* 2011; **18**: 996–1005.
- Alton EW, Baker A, Baker E, et al. The safety profile of a cationic lipid-mediated cystic fibrosis gene transfer agent following repeated monthly aerosol administration to sheep. *Biomaterials* 2013; **34**: 10267–77.
- Alton EW, Boyd AC, Cheng SH, et al. Toxicology study assessing efficacy and safety of repeated administration of lipid/DNA complexes to mouse lung. *Gene Ther* 2014; **21**: 89–95.
- Hyde SC, Pringle IA, Abdullah S, et al. CpG-free plasmids confer reduced inflammation and sustained pulmonary gene expression. *Nat Biotechnol* 2008; **26**: 549–51.
- Quittner AL, Sawicki GS, McMullen A, et al. Psychometric evaluation of the Cystic Fibrosis Questionnaire-Revised in a national sample. *Qual Life Res* 2012; **21**: 1267–78.
- Kerem E, Konstan MW, De Boeck K, et al. Ataluren for the treatment of nonsense-mutation cystic fibrosis: a randomised, double-blind, placebo-controlled phase 3 trial. *Lancet Respir Med* 2014; **2**: 539–47.
- Ramsey BW, Pepe MS, Quan JM, et al. Intermittent administration of inhaled tobramycin in patients with cystic fibrosis. Cystic Fibrosis Inhaled Tobramycin Study Group. *N Engl J Med* 1999; **340**: 23–30.
- Saiman L, Marshall BC, Mayer-Hamblett N, et al. Azithromycin in patients with cystic fibrosis chronically infected with *Pseudomonas aeruginosa*: a randomized controlled trial. *JAMA* 2003; **290**: 1749–56.
- Boyle MP, Bell SC, Konstan MW, et al. A CFTR corrector (lumacaftor) and a CFTR potentiator (ivacaftor) for treatment of patients with cystic fibrosis who have a phe508del CFTR mutation: a phase 2 randomised controlled trial. *Lancet Respir Med* 2014; **2**: 527–38.
- Leigh MW, Kylander JE, Yankaskas JR, Boucher RC. Cell proliferation in bronchial epithelium and submucosal glands of cystic fibrosis patients. *Am J Respir Cell Mol Biol* 1995; **12**: 605–12.
- Chadwick SL, Kingston HD, Stern M, et al. Safety of a single aerosol administration of escalating doses of the cationic lipid GL-67/DOPE/DMPE-PEG5000 formulation to the lungs of normal volunteers. *Gene Ther* 1997; **4**: 937–42.
- Alton EW, Stern M, Farley R, et al. Cationic lipid-mediated CFTR gene transfer to the lungs and nose of patients with cystic fibrosis: a double-blind placebo-controlled trial. *Lancet* 1999; **353**: 947–54.
- Gill DR, Southern KW, Mofford KA, et al. A placebo-controlled study of liposome-mediated gene transfer to the nasal epithelium of patients with cystic fibrosis. *Gene Ther* 1997; **4**: 199–209.
- Zabner J, Cheng SH, Meeker D, et al. Comparison of DNA-lipid complexes and DNA alone for gene transfer to cystic fibrosis airway epithelia in vivo. *J Clin Invest* 1997; **100**: 1529–37.
- Bazzani RP, Cai Y, Hebel HL, Hyde SC, Gill DR. The significance of plasmid DNA preparations contaminated with bacterial genomic DNA on inflammatory responses following delivery of lipoplexes to the murine lung. *Biomaterials* 2011; **32**: 9854–65.
- Hyde SC, Southern KW, Gileadi U, et al. Repeat administration of DNA/liposomes to the nasal epithelium of patients with cystic fibrosis. *Gene Ther* 2000; **7**: 1156–65.

- 
- 21 Rose AC, Goddard CA, Colledge WH, Cheng SH, Gill DR, Hyde SC. Optimisation of real-time quantitative RT-PCR for the evaluation of non-viral mediated gene transfer to the airways. *Gene Ther* 2002; **9**: 1312–20.
- 22 Konstan MW, Davis PB, Wagener JS, et al. Compacted DNA nanoparticles administered to the nasal mucosa of cystic fibrosis subjects are safe and demonstrate partial to complete cystic fibrosis transmembrane regulator reconstitution. *Hum Gene Ther* 2004; **15**: 1255–69.
- 23 Jones AP, Wallis C. Dornase alfa for cystic fibrosis. *Cochrane Database Syst Rev* 2010; **3**: CD001127.
- 24 Wilms EB, Touw DJ, Heijerman HG, van der Ent CK. Azithromycin maintenance therapy in patients with cystic fibrosis: a dose advice based on a review of pharmacokinetics, efficacy, and side effects. *Pediatr Pulmonol* 2012; **47**: 658–65.
- 25 European Medicines Agency. Report of the workshop on endpoints for cystic fibrosis clinical trials 2012. [https://www.ecfs.eu/files/webfm/webfiles/File/documents/EMA\\_Workshop2012.pdf](https://www.ecfs.eu/files/webfm/webfiles/File/documents/EMA_Workshop2012.pdf). (accessed April 26, 2015).