



THE UNIVERSITY *of* EDINBURGH

Edinburgh Research Explorer

Convergence of tamed Euler schemes for a class of stochastic evolution equations

Citation for published version:

Gyongy, I, Sabanis, S & Siska, D 2016, 'Convergence of tamed Euler schemes for a class of stochastic evolution equations' *Stochastic Partial Differential Equations: Analysis and Computations*, vol. 4, no. 2, pp. 225-245. DOI: 10.1007/s40072-015-0057-7

Digital Object Identifier (DOI):

[10.1007/s40072-015-0057-7](https://doi.org/10.1007/s40072-015-0057-7)

Link:

[Link to publication record in Edinburgh Research Explorer](#)

Document Version:

Peer reviewed version

Published In:

Stochastic Partial Differential Equations: Analysis and Computations

General rights

Copyright for the publications made accessible via the Edinburgh Research Explorer is retained by the author(s) and / or other copyright owners and it is a condition of accessing these publications that users recognise and abide by the legal requirements associated with these rights.

Take down policy

The University of Edinburgh has made every reasonable effort to ensure that Edinburgh Research Explorer content complies with UK legislation. If you believe that the public display of this file breaches copyright please contact openaccess@ed.ac.uk providing details, and we will remove access to the work immediately and investigate your claim.



CONVERGENCE OF TAMED EULER SCHEMES FOR A CLASS OF STOCHASTIC EVOLUTION EQUATIONS

ISTVÁN GYÖNGY, SOTIRIOS SABANIS, DAVID ŠIŠKA

ABSTRACT. We prove stability and convergence of a full discretization for a class of stochastic evolution equations with super-linearly growing operators appearing in the drift term. This is done by using the recently developed tamed Euler method, which employs a fully explicit time stepping, coupled with a Galerkin scheme for the spatial discretization.

1. INTRODUCTION

In this paper we investigate the convergence of full discretizations, explicit in time, of stochastic evolution equations

$$du(t) = Au(t)dt + Bu(t)dW(t), \quad t \in [0, T] \quad (1.1)$$

with the drift term governed by a super-linearly growing operator. When the operator appearing in the drift term grows at most linearly then the classical explicit Euler scheme applied to stochastic evolution equations is convergent (when coupled appropriately with the spatial discretization), see, for example, Gyöngy and Millet [6]. If the operator appearing in the drift term grows faster than linearly then one would, in general, not expect the explicit Euler scheme to be convergent (this is the case even in the setting of fully deterministic evolution equations). Instead, one would typically consider the implicit Euler scheme which is convergent in this situation (see, for example, Gyöngy and Millet [6]). The price one pays is the increased computational effort required at each time step of the numerical scheme.

Hutzenthaler, Jentzen and Kloeden [9] have observed that the absolute moments of explicit Euler approximations for stochastic differential equations with super-linearly growing coefficients may diverge to infinity at finite time. This led to development of “tamed” Euler schemes for stochastic differential equations. This was pioneered in Hutzenthaler, Jentzen and Kloeden [10] and, using different techniques, in Sabanis [16]. A taming-like technique in the form of truncation has been proposed by Roberts and Tweedie [14] in the context of Markov chain Monte Carlo methods. Further work on explicit numerical approximations of stochastic differential equations with super-linearly growing coefficients can be found in Tretyakov and Zhang [19], Hutzenthaler and Jentzen [12], Sabanis [17] as well Dareiotis, Kumar and Sabanis [18].

Moreover Hutzenthaler, Jentzen and Kloeden [11] have demonstrated that to apply multilevel Monte Carlo methods (see Heinrich [7], [8] and Giles [5]) to stochastic differential equations with super-linearly growing coefficients one must “tame” the explicit Euler scheme. In this paper we use the idea of “taming” to devise a new type of a convergent explicit scheme for a class of stochastic evolution equations with super-linearly growing operators in the drift term.

The article is organised as follows. In Section 2 we introduce the numerical scheme, give the precise assumptions and state the main result in Theorem 2.3. Essential a priori estimates are proved in Section 3. In Section 4 we first use the a priori estimates and a compactness argument to extract weakly convergent subsequences and limits of the approximation. The remaining step is to identify the weak limit of the approximation of the nonlinear term with the nonlinear term in the equation. This is done using a monotonicity argument in Section 4 where Theorem 2.3 is finally proved. In Section 5 we provide examples of stochastic partial differential equations where the numerical scheme can be applied.

2. MAIN RESULTS

Let $T > 0$. Let $(\Omega, \mathcal{F}, \mathbb{P})$ be a probability space and let $(\mathcal{F}_t)_{t \in [0, T]}$ be a filtration such that \mathcal{F}_0 contains all the \mathbb{P} -null sets of \mathcal{F} .

Let $K > 0$ and $p \in [2, \infty)$ be given constants. Let $p^* := p/(p-1)$. For a reflexive, separable Banach space $(X, \|\cdot\|_X)$ let X^* and $\|\cdot\|_{X^*}$ denote its dual space and the norm on the dual space respectively. For $f \in X^*$ and $v \in X$ we use $\langle f, v \rangle$ to denote the duality pairing. By $L^p(0, T; X)$ we denote the Lebesgue–Bochner space of equivalence classes of measurable functions with values in X that satisfy

$$\|x\|_{L^p(0, T; X)} := \left(\int_0^T \|x(t)\|_X^p dt \right)^{1/p} < \infty.$$

By $L^p(\Omega; X)$ we denote the Lebesgue–Bochner space of random variables with values in X and such that the norm

$$\|x\|_{L^p(\Omega; X)} := (\mathbb{E}(\|x\|_X^p))^{1/p}$$

is finite. Finally by $\mathcal{L}^p(X)$ we denote the Lebesgue–Bochner space of $dt \times \mathbb{P}$ -equivalence classes of $(\mathcal{F}_t)_{t \in [0, T]}$ -adapted and X -valued stochastic process that satisfy

$$\|x\|_{\mathcal{L}^p(X)} := \left(\mathbb{E} \int_0^T \|x(t)\|_X^p dt \right)^{1/p} < \infty.$$

We say that an operator $C : X \times \Omega \rightarrow X^*$ is measurable with respect to some $\mathcal{G} \subseteq \mathcal{F}$ if for any $v, w \in X$ the real-valued random variable $\langle Cv, w \rangle$ is \mathcal{G} -measurable.

We assume that, with respect to $(\mathcal{F}_t)_{t \in [0, T]}$, $(W_t)_{t \in [0, T]}$ is a cylindrical Q -Wiener process with $Q = I$ on a separable Hilbert space $(U, (\cdot, \cdot)_U, |\cdot|_U)$. We assume that there are $(V_1, \|\cdot\|_{V_1})$ and $(V_2, \|\cdot\|_{V_2})$, separable and reflexive Banach spaces that are densely and continuously embedded in H , where $(H, (\cdot, \cdot), |\cdot|)$ is a Hilbert space identified with its dual. We thus have two Gelfand triples

$$V_i \hookrightarrow H \hookrightarrow V_i^*, \quad i \in \{1, 2\}.$$

Let A_i with $i \in \{1, 2\}$ be operators defined on $V_i \times \Omega$ with values in V_i^* . Let B_i with $i \in \{1, 2\}$ be operators defined on $V_i \times \Omega$ with values in $L_2(U, H)$, where $L_2(U, H)$ is the space of Hilbert–Schmidt operators from U to H .

Let $V := V_1 \cap V_2$ and let the norm in V be given by $\|\cdot\| := \|\cdot\|_{V_1} + \|\cdot\|_{V_2}$. Assume that V is separable and dense in both V_1 and V_2 . Using Gajewski, Gröger and Zacharias [4, Kapitel I, Satz 5.13] one observes that the dual V^* of V can be identified with

$$V_1^* + V_2^* := \{f = f_1 + f_2 : f_1 \in V_1^*, f_2 \in V_2^*\}$$

and that for all $f \in V^*$

$$\|f\|_{V^*} = \inf\{\max(\|f_1\|_{V_1^*}, \|f_2\|_{V_2^*}) : f = f_1 + f_2, f_1 \in V_1^*, f_2 \in V_2^*\}.$$

We consider stochastic evolution equations of the form

$$du(t) = [A_1u(t) + A_2u(t)]dt + [B_1u(t) + B_2u(t)]dW(t), \quad t \in [0, T], \quad (2.1)$$

where $u(0) = u_0$ with u_0 a given H -valued and \mathcal{F}_0 -measurable random variable. Let $A := A_1 + A_2$ and $B := B_1 + B_2$. The operator A is defined on $V \times \Omega$ with values in V^* and the operator B is defined on $V \times \Omega$ with values in $L_2(U, H)$. Then we can write (2.1) as (1.1).

We impose the following assumptions on the operators.

Assumption 1. Let $A_i : V_i \times \Omega \rightarrow V_i^*$ be \mathcal{F}_0 -measurable operators for $i \in \{1, 2\}$. Let $B_i : V_i \times \Omega \rightarrow L_2(U, H)$ be such that for any $v \in V_i$, $u \in U$ and $h \in H$ the real-valued random variable $((B_i v)u, h)$ is \mathcal{F}_0 -measurable for $i \in \{1, 2\}$. Moreover assume that the following conditions hold.

Monotonicity:

$$2\langle Av - Aw, v - w \rangle + \|Bv - Bw\|_{L_2(U, H)}^2 \leq K|v - w|^2 \quad \text{for all } v, w \in V.$$

Coercivity: there is $\mu > 0$ such that

$$2\langle A_1v, v \rangle + \|B_1v\|_{L_2(U, H)}^2 \leq -\mu\|v\|_{V_1}^2 + K(1 + |v|^2) \quad \text{for all } v \in V_1.$$

and

$$2\langle A_2v, v \rangle + \|B_2v\|_{L_2(U, H)}^2 \leq K(1 + |v|^2) \quad \text{for all } v \in V_2.$$

Growth:

$$\|A_1v\|_{V_1^*}^2 \leq K(1 + \|v\|_{V_1}^2) \quad \text{for all } v \in V_1$$

and

$$\|A_2v\|_{V_2^*}^p \leq K(1 + \|v\|_{V_2}^p) \quad \text{for all } v \in V_2.$$

as well as

$$\|Bv\|_{L_2(U, H)}^2 \leq K(1 + |v|^2) \quad \text{for all } v \in H.$$

Hemicontinuity: for any v, w and z in V

$$\langle A(v + \epsilon w), z \rangle \rightarrow \langle Av, z \rangle \quad \text{as } \epsilon \rightarrow 0.$$

We now define what is meant by solution of (1.1).

Definition 2.1 (Solution). Let u_0 be an \mathcal{F}_0 -measurable H -valued random variable. We say that a continuous, H -valued and $(\mathcal{F}_t)_{t \in [0, T]}$ -adapted process u is a solution to (1.1) if u is $dt \times \mathbb{P}$ almost everywhere V -valued, if $u \in \mathcal{L}^2(V_1) \cap \mathcal{L}^p(V_2)$ and if for every $t \in [0, T]$ and every $v \in V$, almost surely,

$$(u(t), v) = (u_0, v) + \int_0^t \langle Au(s), v \rangle ds + \int_0^t (v, Bu(s))dW(s).$$

To the best knowledge of the authors, existence and uniqueness has not been proved for this class of stochastic evolution equations. Pardoux [13] considers the situation where the stochastic evolution equation is governed by a sum of monotone, coercive and hemicontinuous operators satisfying certain growth condition. However the operator A_2 in our case only satisfies a type of ‘‘degenerate’’ coercivity condition. Hence the existence theorem from Pardoux [13] does not apply. We prove that a solution to (1.1) must be unique in Theorem 2.2 and we prove existence of the solution in Theorem 2.3.

Theorem 2.2 (Uniqueness). *The solution of (1.1), specified by Definition 2.1, is unique, provided that the Growth and Monotonicity conditions in Assumption 1 are satisfied.*

We prove Theorem 2.2 in Section 4. Let us now describe the discretization scheme for the stochastic evolution equation (2.1). For the space discretization let $(V_m)_{m \in \mathbb{N}}$ be a Galerkin scheme for V . To be precise we assume that $V_m \subseteq V$ are finite dimensional spaces with the dimension of V_m equal to m . We further assume that $V_m \subseteq V_{m+1}$ for all $m \in \mathbb{N}$ and that

$$\lim_{m \rightarrow \infty} \inf \{ \|v - \varphi\| : \varphi \in V_m \} = 0 \quad \forall v \in V.$$

(this is referred to as the limited completeness of the Galerkin scheme). We will need the following projection operators.

Assumption 2. For any $m \in \mathbb{N}$ let $\Pi_m : V^* \rightarrow V_m$ satisfy the following:

- (1) For any $v \in V_m$, $\Pi_m v = v$.
- (2) If $f \in V^*$ and $v \in V$ then $\langle f, \Pi_m v \rangle = \langle v, \Pi_m f \rangle$.
- (3) If $g, h \in H$ then $(\Pi_m g, h) = (\Pi_m h, g)$ and $|\Pi_m h| \leq |h|$.
- (4) There is a constant, depending on m and denoted by $\mathfrak{c}(m)$, such that

$$|\Pi_m f|^2 \leq \mathfrak{c}(m) \|f\|_{V^*}^2 \quad \text{for all } f \in V^*.$$

In applications this assumption is easily satisfied. In particular if $\{\varphi_j \in V : j = 1, 2, \dots\}$ is an orthonormal basis in H then taking $V_m := \text{span}\{\varphi_1, \dots, \varphi_m\}$ is a Galerkin scheme for V . Taking $\Pi_m f := \sum_{j=1}^m \langle f, \varphi_j \rangle \varphi_j$ satisfies the first three conditions in Assumption 2. Moreover, the following holds

$$|\Pi_m f|^2 = \left| \sum_{j=1}^m \langle f, \varphi_j \rangle \varphi_j \right|^2 = \sum_{j=1}^m \langle f, \varphi_j \rangle^2 \leq \|f\|_{V^*}^2 \sum_{j=1}^m \|\varphi_j\|_V^2 = \mathfrak{c}(m) \|f\|_{V^*}^2,$$

where $\mathfrak{c}(m) := \sum_{j=1}^m \|\varphi_j\|_V^2$. Thus the fourth condition in Assumption 2 is also satisfied. Let $\{\chi_i\}_{i \in \mathbb{N}}$ be an orthonormal basis of U . Fix $k \in \mathbb{N}$ and define

$$W_k(t) := \sum_{j=1}^k (W(t), \chi_j)_U \chi_j.$$

For the time discretization take $n \in \mathbb{N}$, let $\tau_n := T/n$ and define the grid points on an equidistant grid as $t_i^n := \tau_n i$, $i = 0, 1, \dots, n$. Further consider some sequence $((n_\ell, m_\ell, k_\ell))_{\ell \in \mathbb{N}}$ such that n_ℓ, m_ℓ and k_ℓ all go to infinity as $\ell \rightarrow \infty$.

Let c denote a generic positive constant that may depend on T , on the constants arising in the continuous embeddings $V_i \hookrightarrow H \hookrightarrow V_i^*$, $i = 1, 2$ and on the constants arising in Assumptions 1 and 3 but that is always independent of the discretization parameters m , k and n . Define $\kappa_{n_\ell}(t) = t_i^{n_\ell}$ if $t \in [t_i^{n_\ell}, t_{i+1}^{n_\ell})$ for $i = 0, \dots, n_\ell - 1$ and $\kappa_{n_\ell}(T) = T$. Fix some $\ell \in \mathbb{N}$ (and hence m_ℓ, n_ℓ and k_ℓ). Let $u_\ell(0)$ be a V_{m_ℓ} valued \mathcal{F}_0 -measurable approximation of u_0 . For example we can take $u_\ell(0) := \Pi_{m_\ell} u_0$ but other approximations are possible. For $t > 0$ we define a process u_ℓ by

$$\begin{aligned} u_\ell(t) &= u_\ell(0) + \int_0^t \Pi_{m_\ell} [A_1 u_\ell(\kappa_{n_\ell}(s)) + A_{2,\ell} u_\ell(\kappa_{n_\ell}(s))] ds \\ &\quad + \int_0^t \Pi_{m_\ell} B u_\ell(\kappa_{n_\ell}(s)) dW_{k_\ell}(s), \end{aligned} \tag{2.2}$$

where we use the ‘‘tamed’’ operator $A_{2,\ell}$ defined by

$$A_{2,\ell} v := \frac{1}{1 + n_\ell^{-1/2} |\Pi_{m_\ell} A_2 v|} A_2 v \tag{2.3}$$

for any $v \in V_2$. We will use the following notation: $\bar{u}_\ell(t) := u_\ell(\kappa_{n_\ell}(t))$ and $a_\ell(v) := \Pi_{m_\ell}[A_1 v + A_{2,\ell} v]$. Then (2.2) is equivalent to

$$u_\ell(t) = u_\ell(0) + \int_0^t a_\ell(\bar{u}_\ell(s)) ds + \int_0^t \Pi_{m_\ell} B \bar{u}_\ell(s) dW_{k_\ell}(s). \quad (2.4)$$

In particular at the time-grid points we have, for $i = 0, 1, \dots, n_\ell - 1$,

$$u_\ell(t_{i+1}^{n_\ell}) = u_\ell(t_i^{n_\ell}) + a_\ell(u_\ell(t_i^{n_\ell})) \tau_{n_\ell} + \Pi_{m_\ell} B u_\ell(t_i^{n_\ell}) \Delta W_{k_\ell}(t_{i+1}),$$

where $\Delta W_{k_\ell}(t_{i+1}^{n_\ell}) := W_{k_\ell}(t_{i+1}^{n_\ell}) - W_{k_\ell}(t_i^{n_\ell})$. This in turn is equivalent to

$$\frac{u_\ell(t_{i+1}^{n_\ell}) - u_\ell(t_i^{n_\ell})}{\tau_{n_\ell}} = a_\ell(u_\ell(t_i^{n_\ell})) + \Pi_{m_\ell} B u_\ell(t_i^{n_\ell}) \frac{\Delta W_{k_\ell}(t_{i+1}^{n_\ell})}{\tau_{n_\ell}}.$$

We list below the properties which are satisfied by the tamed operator $A_{2,\ell}$. These are consequences of its structure and the assumed properties of A_2 . For brevity let, for any $v \in V_2$,

$$T_\ell(v) := \frac{1}{1 + n_\ell^{-1/2} |\Pi_{m_\ell} A_2 v|}. \quad (2.5)$$

Then for any $v \in V_2$,

$$|\Pi_{m_\ell} A_{2,\ell} v| = T_\ell(v) |\Pi_{m_\ell} A_2 v| \leq n_\ell^{1/2} \quad (2.6)$$

and also, using the Growth assumption on A_2 ,

$$\|A_{2,\ell} v\|_{V_2^*}^{p^*} = T_\ell(v)^{p^*} \|A_2 v\|_{V_2^*}^{p^*} \leq K(1 + \|v\|_{V_2}^p). \quad (2.7)$$

Furthermore, using the Coercivity assumption on A_2 , we note that for all $v \in V_{m_\ell}$ we have

$$2\langle A_{2,\ell} v, v \rangle = 2T_\ell(v) \langle A_2 v, v \rangle \leq K(1 + |v|^2). \quad (2.8)$$

Thus the weaker coercivity assumption that has been made about A_2 is retained. Consider, for a moment, that A_2 satisfies the ‘‘usual’’ coercivity condition

$$2\langle A_2 v, v \rangle + \|B_2 v\|_{L_2(U,H)}^2 \leq -\mu \|v\|_{V_2}^p + K(1 + |v|^2) \quad \text{for all } v \in V_2.$$

We see that in this case the best coercivity we can get from this for $A_{2,\ell}$ is again only (2.8). Hence to obtain the necessary a priori estimates we will need an interpolation inequality between V_2 and V_1 with H .

Assumption 3. There are constants $\lambda \in [0, 2/p)$ and $\Lambda > 0$ such that for any $v \in V$

$$\|v\|_{V_2} \leq \Lambda \|v\|_{V_1}^\lambda |v|^{1-\lambda}.$$

Note that in order to overcome the difficulty with coercivity it would suffice to have Assumption 3 satisfied with $\lambda \in [0, 2/p]$. However monotonicity of A_2 is not preserved by taming. To overcome this we will need to show that $A_{2,\ell} \bar{u}_\ell - A_2 \bar{u}_\ell \rightarrow 0$ in $\mathcal{L}^{p^*}(V_2^*)$. To achieve this we use the fact that $\lambda \in [0, 2/p)$ in Lemma 4.3 and the following observation: Assumption 3 implies that there is $\eta > 0$ such that

$$\|v\|_{V_2}^{p(1+\eta)} \leq c \|v\|_{V_1}^2 |v|^\rho,$$

where $\rho := (1 - \lambda)p(1 + \eta)$. From this it follows that if $v \in L^2(\Omega; L^2(0, T; V_1))$ and $v \in L^{2\rho}(\Omega; L^\infty(0, T; H))$ then

$$\mathbb{E} \int_0^T \|v(s)\|_{V_2}^{p(1+\eta)} ds \leq c \left[\mathbb{E} \sup_{s \leq t} |v(s)|^{2\rho} + \mathbb{E} \left(\int_0^T \|v(s)\|_{V_1}^2 ds \right)^2 \right].$$

Thus we see that Assumption 3 allows us to control the approximate solution in the $L^{p(1+\eta)}((0, T) \times \Omega; V_2)$ norm, provided that we can control the approximate solution in the norms of $L^2(\Omega; L^2(0, T; V_1))$ and $L^{2\rho}(\Omega; L^\infty(0, T; H))$.

Let us take $q_0 := \max(4, 2\rho)$. Now we can state the main result of this paper.

Theorem 2.3. *Let Assumptions 1, 2 and 3 be satisfied. Let $u_0 \in L^{q_0}(\Omega; H)$ and let $u_\ell(0) \rightarrow u_0$ in $L^{q_0}(\Omega; H)$. Assume that $\frac{c(m_\ell)}{n_\ell} \rightarrow 0$ as $\ell \rightarrow \infty$. Then there exists a unique solution u to (1.1) and $\bar{u}_\ell \rightarrow u$ in $\mathcal{L}^2(V_1)$ and in $\mathcal{L}^p(V_2)$ and $u_\ell(T) \rightarrow u(T)$ in $L^2(\Omega; H)$ as $\ell \rightarrow \infty$.*

In Section 5 we provide examples of stochastic partial differential equations where Theorem 2.3 can be applied. We also compute $c(m)$ in case of the spectral Galerkin method to make the implications of the space-time coupling constraint more explicit. The crucial point is that the requirement is no more onerous than in the case of equations with operators growing at most linearly.

3. A PRIORI ESTIMATES

We start with an important observation that allows us to use standard results on bounds of stochastic integrals driven by finite dimensional Wiener processes.

Remark 3.1. Recall that $(\chi_j)_{j \in \mathbb{N}}$ is an orthonormal basis in U . Moreover recall that $\bar{u}_\ell(t) := u_\ell(\kappa_{n_\ell}(t))$ and that $a_\ell(v) := \Pi_{m_\ell}[A_1 v + A_{2,\ell} v]$. For each $j \in \mathbb{N}$ a Wiener processes \mathcal{W}_j is obtained by taking $\mathcal{W}_j(t) := (W(t), \chi_j)_U$. If $i \neq j$ then \mathcal{W}_i and \mathcal{W}_j are independent. Furthermore (2.4) is equivalent to

$$u_\ell(t) = u_\ell(0) + \int_0^t a_\ell(\bar{u}_\ell(s)) ds + \sum_{j=1}^{k_\ell} \int_0^t \Pi_{m_\ell} B \bar{u}_\ell(s) \chi_j d\mathcal{W}_j(s).$$

Fix $\ell \in \mathbb{N}$ (and thus k_ℓ, m_ℓ and n_ℓ are also fixed). Then using the Growth assumptions on A_1 and B , Assumption 2 as well as (2.6) one observes that $|a_\ell(v)|^2 \leq 2c(m)K(1 + |v|^2) + 2n_\ell$ and $|\Pi_{m_\ell} B v \chi_j|^2 \leq K(1 + |v|^2)$. Hence one knows that, for $q \geq 1$,

$$\mathbb{E} \sup_{0 \leq t \leq T} |u_\ell(t)|^q < \infty,$$

provided that $\mathbb{E}|u_\ell(0)|^q < \infty$. Clearly, at this point, one cannot claim that this bound is independent of ℓ .

One applies Itô's formula to (2.4) to obtain

$$\begin{aligned} |u_\ell(t)|^2 &= |u_\ell(0)|^2 + \int_0^t \left[2\langle a_\ell(\bar{u}_\ell(s)), u_\ell(s) \rangle + \sum_{j=1}^{k_\ell} |\Pi_{m_\ell} B \bar{u}_\ell(s) \chi_j|^2 \right] ds \\ &\quad + \int_0^t 2\langle B \bar{u}_\ell(s), u_\ell(s) \rangle dW_{k_\ell}(s), \end{aligned}$$

which can be rewritten as

$$\begin{aligned} |u_\ell(t)|^2 &= |u_\ell(0)|^2 + \int_0^t \left[2\langle a_\ell(\bar{u}_\ell(s)), \bar{u}_\ell(s) \rangle + \sum_{j=1}^{k_\ell} |\Pi_{m_\ell} B \bar{u}_\ell(s) \chi_j|^2 \right] ds \\ &\quad + 2 \int_0^t \langle a_\ell(\bar{u}_\ell(s)), u_\ell(s) - \bar{u}_\ell(s) \rangle ds \\ &\quad + \int_0^t 2\langle B \bar{u}_\ell(s), u_\ell(s) \rangle dW_{k_\ell}(s), \end{aligned} \tag{3.1}$$

in order to apply the coercivity assumption so as to obtain the a priori estimates for the discretized equation.

First we concentrate on the term that arises from the ‘‘correction’’ that one has to make to use the coercivity assumption due to the use of an explicit scheme.

Lemma 3.2. *Let the Growth condition in Assumption 1 be satisfied. Let Assumption 2 hold. Let $q \geq 1$ be given. Then*

$$\begin{aligned} & \mathbb{E} \left(\frac{1}{\tau_{n_\ell}} \int_0^t |u_\ell(s) - \bar{u}_\ell(s)|^2 ds \right)^q \\ & \leq c_{T,q} \left(1 + (\mathfrak{c}(m)\tau_{n_\ell})^q \mathbb{E} \left(\int_0^t \|\bar{u}_\ell(s)\|_{V_1}^2 ds \right)^q + \mathbb{E} \int_0^t |\bar{u}_\ell(s)|^{2q} ds \right) \end{aligned} \quad (3.2)$$

and

$$\begin{aligned} & \mathbb{E} \left(\int_0^t |a_\ell(\bar{u}_\ell(s))| |u_\ell(s) - \bar{u}_\ell(s)| ds \right)^q \\ & \leq c_{T,q} \left(1 + (\mathfrak{c}(m)\tau_{n_\ell})^q \mathbb{E} \left(\int_0^t \|\bar{u}_\ell(s)\|_{V_1}^2 ds \right)^q + \mathbb{E} \int_0^t |\bar{u}_\ell(s)|^{2q} ds \right). \end{aligned} \quad (3.3)$$

Proof. From (2.2) it is clear that

$$\begin{aligned} I_{1,\ell}(t) & := \mathbb{E} \left(\frac{1}{\tau_{n_\ell}} \int_0^t |u_\ell(s) - \bar{u}_\ell(s)|^2 ds \right)^q \\ & = \mathbb{E} \left(\int_0^t \frac{1}{\tau_{n_\ell}} \left| \int_{\kappa_{n_\ell}(s)}^s a_\ell(\bar{u}_\ell(r)) dr + \int_{\kappa_{n_\ell}(s)}^s \Pi_{m_\ell} B \bar{u}_\ell(r) dW_{k_\ell}(r) \right|^2 ds \right)^q \\ & \leq 2^q \mathbb{E} \left(\frac{1}{\tau_{n_\ell}} \int_0^t \left| \int_{\kappa_{n_\ell}(s)}^s a_\ell(\bar{u}_\ell(r)) dr \right|^2 + \left| \int_{\kappa_{n_\ell}(s)}^s \Pi_{m_\ell} B \bar{u}_\ell(r) dW_{k_\ell}(r) \right|^2 ds \right)^q. \end{aligned}$$

Applying Hölder's inequality yields

$$\begin{aligned} I_{1,\ell}(t) & \leq 2^q \mathbb{E} \left(\frac{1}{\tau_{n_\ell}} \int_0^t \left[(s - \kappa_{n_\ell}(s)) \int_{\kappa_{n_\ell}(s)}^s |a_\ell(\bar{u}_\ell(r))|^2 dr \right. \right. \\ & \quad \left. \left. + \left| \int_{\kappa_{n_\ell}(s)}^s \Pi_{m_\ell} B \bar{u}_\ell(r) dW_{k_\ell}(r) \right|^2 \right] ds \right)^q \\ & \leq c_q \mathbb{E} \left(\frac{1}{\tau_{n_\ell}} \int_0^t \tau_{n_\ell}^2 |a_\ell(\bar{u}_\ell(s))|^2 ds \right)^q \\ & \quad + c_q \mathbb{E} \left(\frac{1}{\tau_{n_\ell}} \int_0^t \left| \int_{\kappa_{n_\ell}(s)}^s \Pi_{m_\ell} B \bar{u}_\ell(r) dW_{k_\ell}(r) \right|^2 ds \right)^q. \end{aligned}$$

Using Assumption 2 and (2.6) one obtains

$$\begin{aligned} I_{1,\ell}(t) & \leq c_q \mathbb{E} \left(\int_0^t \tau_{n_\ell} [2\mathfrak{c}(m) \|A_1 \bar{u}_\ell(s)\|_{V_1^*}^2 + 2n_\ell] ds \right)^q \\ & \quad + c_q \mathbb{E} \left(\int_0^t \frac{1}{\tau_{n_\ell}} \left| \int_{\kappa_{n_\ell}(s)}^s \Pi_{m_\ell} B \bar{u}_\ell(r) dW_{k_\ell}(r) \right|^2 ds \right)^q := I_{2,\ell}(t) + I_{3,\ell}(t). \end{aligned} \quad (3.4)$$

The Growth assumption on A_1 implies that

$$I_{2,\ell}(t) \leq c_{T,q} \left(1 + (\mathfrak{c}(m)\tau_{n_\ell})^q \mathbb{E} \left(\int_0^t \|\bar{u}_\ell(s)\|_{V_1}^2 ds \right)^q \right).$$

Using Hölder's inequality leads to

$$I_{3,\ell}(t) \leq c_{T,q} \mathbb{E} \int_0^t \frac{1}{\tau_{n_\ell}^q} \left| \int_{\kappa_{n_\ell}(s)}^s \Pi_{m_\ell} B \bar{u}_\ell(r) dW_{k_\ell}(r) \right|^{2q} ds.$$

Due to Remark 3.1 and the Growth assumption on B one observes that

$$\begin{aligned} I_{3,\ell}(t) &\leq c_{T,q} \int_0^t \frac{1}{\tau_{n_\ell}^q} \mathbb{E} \left| \int_{\kappa_{n_\ell}(s)}^s \|B\bar{u}_\ell(r)\|_{L_2(U,H)}^2 dr \right|^q ds \\ &\leq c_{T,q} \mathbb{E} \int_0^t \frac{1}{\tau_{n_\ell}^q} (s - \kappa_{n_\ell}(s))^q \|B\bar{u}_\ell(s)\|_{L_2(U,H)}^{2q} ds \\ &\leq c_{T,q} \left(1 + \mathbb{E} \int_0^t |\bar{u}_\ell(s)|^{2q} ds \right). \end{aligned}$$

This implies (3.2). Moreover Assumption 2 and (2.6) imply that

$$\begin{aligned} I_\ell(t) &:= \mathbb{E} \left(\int_0^t |a_\ell(\bar{u}_\ell(s))| |u_\ell(s) - \bar{u}_\ell(s)| ds \right)^q \\ &\leq 2^q \mathbb{E} \left(\int_0^t \tau_{n_\ell} |a_\ell(\bar{u}_\ell(s))|^2 + \frac{1}{\tau_{n_\ell}} |u_\ell(s) - \bar{u}_\ell(s)|^2 ds \right)^q \\ &\leq c_q \mathbb{E} \left(\int_0^t [\mathbf{c}(m)\tau_{n_\ell} \|A_1 \bar{u}_\ell(s)\|_{V_1^*}^2 + \tau_{n_\ell} n_\ell] ds \right)^q \\ &\quad + c_q \mathbb{E} \left(\frac{1}{\tau_{n_\ell}} \int_0^t |u_\ell(s) - \bar{u}_\ell(s)|^2 ds \right)^q. \end{aligned}$$

Applying the Growth assumption on A_1 yields

$$\begin{aligned} I_\ell(t) &\leq c_{T,q} \left(1 + (\mathbf{c}(m)\tau_{n_\ell})^q \mathbb{E} \left(\int_0^t \|\bar{u}_\ell(s)\|_{V_1}^2 ds \right)^q \right) \\ &\quad + c_q \mathbb{E} \left(\frac{1}{\tau_{n_\ell}} \int_0^t |u_\ell(s) - \bar{u}_\ell(s)|^2 ds \right)^q. \end{aligned} \tag{3.5}$$

Using (3.2) in (3.5) concludes the proof. \square

Theorem 3.3 (A priori estimate). *Let the Coercivity and Growth conditions in Assumption 1 hold. Let Assumption 2 be satisfied. Let $q \geq 1$ be given and assume that $\mathbb{E}|u_\ell(0)|^{2q} < c$ and that $u_\ell(0)$ is \mathcal{F}_0 -measurable. There is $\epsilon \in (0, \infty)$ such that for all $\ell \in \{\ell' \in \mathbb{N} : \mathbf{c}(m_{\ell'})\tau_{n_{\ell'}} < \epsilon\}$ we have, for any $t \in [0, T]$,*

$$\mathbb{E} \sup_{s \in [0,t]} |u_\ell(s)|^{2q} + \mu^q \mathbb{E} \left(\int_0^t \|\bar{u}_\ell(s)\|_{V_1}^2 ds \right)^q \leq c(1 + \mathbb{E}|u_\ell(0)|^{2q}).$$

Proof. Applying the Coercivity assumption in (3.1), raising to power $q \geq 1$, taking the supremum over $s \leq t$ and taking the expectation yields

$$\begin{aligned} &\mathbb{E} \sup_{s \leq t} |u_\ell(s)|^{2q} + \mu^q \mathbb{E} \left(\int_0^t \|\bar{u}_\ell(s)\|_{V_1}^2 ds \right)^q \leq c_{T,q} \left[1 + \mathbb{E}|u_\ell(0)|^{2q} \right. \\ &\quad + \mathbb{E} \left(\int_0^t |\bar{u}_\ell(s)|^2 ds \right)^q + \mathbb{E} \left(\int_0^t |a_\ell(\bar{u}_\ell(s))| |u_\ell(s) - \bar{u}_\ell(s)| ds \right)^q \\ &\quad \left. + \mathbb{E} \sup_{s \leq t} \left| \int_0^s (u_\ell(s), B\bar{u}_\ell(s)) dW_{k_\ell}(s) \right|^q \right]. \end{aligned} \tag{3.6}$$

Using Lemma 3.2 in (3.6) results in

$$\begin{aligned} & \mathbb{E} \sup_{s \leq t} |u_\ell(s)|^{2q} + \frac{\mu^q}{2} \mathbb{E} \left(\int_0^t \|\bar{u}_\ell(s)\|_{V_1}^2 ds \right)^q \\ & \leq c_{T,q} \left[1 + \mathbb{E}|u_\ell(0)|^{2q} + \mathbb{E} \left(\int_0^t |\bar{u}_\ell(s)|^2 ds \right)^q \right. \\ & \quad \left. + \mathbb{E} \int_0^t |\bar{u}_\ell(s)|^{2q} ds + \mathbb{E} \sup_{s \leq t} \left(\int_0^s (u_\ell(r), B\bar{u}_\ell(r)) dW_{k_\ell}(r) \right)^q \right]. \end{aligned} \quad (3.7)$$

Using Burkholder–Davis–Gundy inequality one obtains

$$\begin{aligned} I_\ell & := c_{T,q} \mathbb{E} \sup_{s \leq t} \left| \int_0^s (u_\ell(r), B\bar{u}_\ell(r)) dW_{k_\ell}(r) \right|^q \\ & \leq c_{T,q} \mathbb{E} \left| \int_0^t |u_\ell(s)|^2 \|B\bar{u}_\ell(s)\|_{L_2(U,H)}^2 ds \right|^{q/2} \\ & \leq c_{T,q} \mathbb{E} \left[\sup_{s \leq t} |u_\ell(s)|^q \left(\int_0^t \|B\bar{u}_\ell(s)\|_{L_2(U,H)}^2 ds \right)^{q/2} \right]. \end{aligned}$$

Young's inequality and the Growth assumption on B imply that

$$\begin{aligned} I_\ell & \leq \frac{1}{2} \mathbb{E} \sup_{s \leq t} |u_\ell(s)|^{2q} + c \mathbb{E} \left(\int_0^t \|B\bar{u}_\ell(s)\|_{L_2(U,H)}^2 ds \right)^q \\ & \leq \frac{1}{2} \mathbb{E} \sup_{s \leq t} |u_\ell(s)|^{2q} + c \left(1 + \int_0^t \mathbb{E} \sup_{r \leq s} |u_\ell(r)|^{2q} ds \right). \end{aligned}$$

Applying this in (3.7) leads to

$$\begin{aligned} \frac{1}{2} \mathbb{E} \sup_{s \leq t} |u_\ell(s)|^{2q} + \frac{\mu^q}{2} \mathbb{E} \left(\int_0^t \|\bar{u}_\ell(s)\|_{V_1}^2 ds \right)^q & \leq c \left[1 + \mathbb{E}|u_\ell(0)|^{2q} \right. \\ & \quad \left. + \int_0^t \mathbb{E} \sup_{r \leq s} |u_\ell(r)|^{2q} ds \right]. \end{aligned}$$

Application of Gronwall's lemma yields

$$\mathbb{E} \sup_{s \in [0,t]} |u_\ell(s)|^{2q} + \mu^q \mathbb{E} \left(\int_0^t \|\bar{u}_\ell(s)\|_{V_1}^2 ds \right)^q \leq c (1 + \mathbb{E}|u_\ell(0)|^{2q}). \quad (3.8)$$

□

Now we use Theorem 3.3 and Assumption 3 to obtain the remaining required estimates.

Corollary 3.4 (Remaining a priori estimates). *Let the Growth and Coercivity conditions in Assumption 1 be satisfied. Let Assumptions 2 and 3 hold. Let $u_\ell(0)$ be bounded in $L^{q_0}(\Omega; H)$, uniformly with respect to ℓ . There is $\epsilon \in (0, \infty)$ such that for all $\ell \in \{\ell' \in \mathbb{N} : \mathfrak{c}(m'_\ell \tau_{n_{\ell'}}) < \epsilon\}$ we have*

$$\mathbb{E} \int_0^T \|A_1 \bar{u}_\ell\|_{V_1^*}^2 ds \leq c, \quad \mathbb{E} \int_0^T \|B\bar{u}_\ell\|_{L_2(U,H)}^2 ds \leq c. \quad (3.9)$$

Furthermore

$$\mathbb{E} \int_0^T |u_\ell(s) - \bar{u}_\ell(s)|^2 ds \leq c \tau_{n_\ell}. \quad (3.10)$$

Finally, for some $\eta > 0$,

$$\mathbb{E} \int_0^T \|\bar{u}_\ell\|_{V_2}^{p(1+\eta)} ds \leq c \quad (3.11)$$

and

$$\mathbb{E} \int_0^T \|A_2 \bar{u}_\ell\|_{V_2^*}^{p^*(1+\eta)} ds \leq c, \quad \mathbb{E} \int_0^T \|A_{2,\ell} \bar{u}_\ell\|_{V_2^*}^{p^*(1+\eta)} ds \leq c. \quad (3.12)$$

Proof. Inequality (3.9) follows directly from the Growth assumptions on A_1 and B and from Theorem 3.3 with $q = 1$. Using (3.2), together with Theorem 3.3 with $q = 1$, yields (3.10). Since $u_\ell(0)$ is assumed to be bounded in $L^{q_0}(\Omega; H)$, uniformly in ℓ , one can conclude, using Theorem 3.3, that

$$\mathbb{E} \sup_{s \in [0,t]} |u_\ell(s)|^{2\rho} \leq c \quad \text{and} \quad \mathbb{E} \left(\int_0^t \|\bar{u}_\ell(s)\|_{V_1}^2 ds \right)^2 \leq c.$$

This, together with Assumption 3, yields (3.11). Finally, (3.11), the assumption on the growth of A_2 and (2.7) lead to (3.12). \square

4. CONVERGENCE

Having obtained the required a priori estimates we can use compactness arguments to extract weakly convergent subsequences of the approximation.

Lemma 4.1. *Let the Growth and Coercivity conditions in Assumption 1 hold. Let Assumptions 2 and 3 be satisfied. Let $u_\ell(0) \rightarrow u_0$ in $L^{q_0}(\Omega; H)$. Let $\frac{c(m_\ell)}{n_\ell} \rightarrow 0$ as $\ell \rightarrow \infty$. Then there is a subsequence of the sequence ℓ , which we denote ℓ' , and $u \in \mathcal{L}^2(V_1) \cap \mathcal{L}^p(V_2)$ such that, as $\ell' \rightarrow \infty$,*

$$\bar{u}_{\ell'} \rightharpoonup u \quad \text{in } \mathcal{L}^2(V_1) \quad \text{and in } \mathcal{L}^p(V_2).$$

Furthermore there are $a_1^\infty \in \mathcal{L}^2(V_1^)$, $a_2^\infty \in \mathcal{L}^{p^*}(V_2^*)$ and $b^\infty \in \mathcal{L}^2(L_2(U, H))$ such that, as $\ell' \rightarrow \infty$,*

$$A_1 \bar{u}_{\ell'} \rightharpoonup a_1^\infty \quad \text{in } \mathcal{L}^2(V_1^*), \quad A_{2,\ell'} \bar{u}_{\ell'} \rightharpoonup a_2^\infty \quad \text{in } \mathcal{L}^{p^*}(V_2^*)$$

and

$$B \bar{u}_{\ell'} \rightharpoonup b^\infty \quad \text{in } \mathcal{L}^2(L_2(U, H)).$$

Finally, there is $\xi \in L^{q_0}(\Omega; H)$ such that $\bar{u}_{\ell'}(T) = u_{\ell'}(T) \rightharpoonup \xi$ in $L^{q_0}(\Omega; H)$.

Proof. The sequence (\bar{u}_ℓ) is bounded in $\mathcal{L}^2(V_1)$ due to Theorem 3.3 and in $\mathcal{L}^p(V_2)$ due to Corollary 3.4. The sequences $(A_1 \bar{u}_\ell)$, $(A_{2,\ell} \bar{u}_\ell)$ and $(B \bar{u}_\ell)$ are bounded in $\mathcal{L}^2(V_1^*)$, $\mathcal{L}^{p^*}(V_2^*)$ and in $\mathcal{L}^2(L_2(U, H))$ respectively, due to Corollary 3.4. Finally the sequence $(u_\ell(T))$ is bounded in $L^{q_0}(\Omega; H)$ due to Theorem 3.3.

Since it is assumed that V_1 , V_2 are reflexive, it follows that $\mathcal{L}^2(V_1)$ and $\mathcal{L}^p(V_2)$ are reflexive. A bounded sequence in a reflexive Banach space must have a weakly convergent subsequence (see e.g. Brézis [2, Theorem 3.18]). Applying this to the sequences in question concludes the proof of the lemma. \square

Let $a^\infty := a_1^\infty + a_2^\infty$. Then $a^\infty \in \mathcal{L}^{p^*}(V^*)$. Due to Lemma 4.1 $a_\ell(\bar{u}_\ell) \rightharpoonup a^\infty$ in $\mathcal{L}^{p^*}(V^*)$ as $\ell' \rightarrow \infty$, provided that $\frac{c(m_\ell)}{n_\ell} \rightarrow 0$. The following lemma provides the equation satisfied by the weak limits of the approximations.

Lemma 4.2. *Let the Growth and Coercivity conditions in Assumption 1 hold. Let Assumptions 2 and 3 be satisfied. Let $u_\ell(0) \rightarrow u_0$ in $L^{q_0}(\Omega; H)$. Let $\frac{c(m_\ell)}{n_\ell} \rightarrow 0$ as $\ell \rightarrow \infty$. Then there is an H -valued adapted continuous process \tilde{u} on $[0, T]$ such that $u = \tilde{u} dt \times \mathbb{P}$ -almost everywhere on $(0, T) \times \Omega$. Furthermore, for almost every $(t, \omega) \in (0, T) \times \Omega$,*

$$\tilde{u}(t) = u_0 + \int_0^t a^\infty(s) ds + \int_0^t b^\infty(s) dW(s) \quad (4.1)$$

and almost surely

$$\tilde{u}(T) = u_0 + \int_0^T a^\infty(s)ds + \int_0^T b^\infty(s)dW(s). \quad (4.2)$$

In the rest of this paper we will write u instead of \tilde{u} for notational simplicity.

Proof. Fix $M \in \mathbb{N}$. Let φ be a V_M -valued adapted stochastic process such that $|\varphi(t)| < M$ for all $t \in [0, T]$ and $\omega \in \Omega$. For $g \in U$ let $\tilde{\Pi}_m g := \sum_{j=1}^m (\chi_j, g)U\chi_j$ and note that for any $v \in V$ one has $Bv\tilde{\Pi}_m \in L_2(U, H)$. From (2.2) one observes that

$$\begin{aligned} (u_{\ell'}(t), \varphi(t)) &= (u_{\ell'}(0), \varphi(t)) + \left\langle \int_0^t a_{\ell'}(\bar{u}_{\ell'}(s))ds, \varphi(t) \right\rangle \\ &\quad + \left(\int_0^t B\bar{u}_{\ell'}(s)\tilde{\Pi}_{m_{\ell'}}dW(s), \varphi(t) \right). \end{aligned} \quad (4.3)$$

Let $G : \mathcal{L}^{p^*}(V^*) \rightarrow \mathcal{L}^{p^*}(V^*)$ be given by $(Gv)(t) := \int_0^t v(s)ds$. Moreover, let $H : \mathcal{L}^2(L_2(U, H)) \rightarrow \mathcal{L}^2(H)$ be given by $(Hv)(t) := \int_0^t v(s)dW(s)$. Integrating (4.3) from 0 to T and taking the expectation yields

$$\begin{aligned} \mathbb{E} \int_0^T (u_{\ell'}(t), \varphi(t))dt &= \mathbb{E} \int_0^T (u_{\ell'}(0), \varphi(t))dt \\ &\quad + \mathbb{E} \int_0^T \langle (Ga_{\ell'}(\bar{u}_{\ell'}))(t), \varphi(t) \rangle dt + \mathbb{E} \int_0^T ((HB\bar{u}_{\ell'}\tilde{\Pi}_{m_{\ell'}})(t), \varphi(t)) dt. \end{aligned}$$

The operator G is linear and bounded and as such it is weakly-weakly continuous. This operator H is clearly linear. Furthermore, due to Itô's isometry,

$$\begin{aligned} \|Hv\|_{\mathcal{L}^2(H)}^2 &= \mathbb{E} \int_0^T |(Hv)(s)|^2 ds \\ &= \int_0^T \mathbb{E} \left| \int_0^t v(s)dW(s) \right|^2 dt = \mathbb{E} \int_0^T \int_0^t |v(s)|^2 ds dt \leq T \|v\|_{\mathcal{L}^2(L_2(U, H))}^2. \end{aligned}$$

Thus the operator H is also bounded. It follows that H is also weakly-weakly continuous. Therefore, taking the limit as $\ell' \rightarrow \infty$ and using Lemma 4.1, one obtains

$$\begin{aligned} \mathbb{E} \int_0^T (u(t), \varphi(t))dt &= \mathbb{E} \int_0^T (u_0, \varphi(t))dt \\ &\quad + \mathbb{E} \int_0^T \langle (Ga^\infty)(t), \varphi(t) \rangle dt + \mathbb{E} \int_0^T ((Hb^\infty)(t), \varphi(t)) dt. \end{aligned}$$

This holds for any φ as specified at the beginning of the proof. By letting $M \rightarrow \infty$ and using the limited completeness of the Galerkin scheme it follows that this also holds for any $\varphi \in \mathcal{L}^p(V)$. Thus

$$u(t) = u_0 + \int_0^t a^\infty(s)ds + \int_0^t b^\infty(s)dW(s) \quad (4.4)$$

holds for almost all $(t, \omega) \in (0, T) \times \Omega$.

Let φ be a V_M -valued and \mathcal{F}_T -measurable random variable such that $\mathbb{E}\|\varphi\|_V^2 < \infty$. Setting $t = T$ in (4.3) and taking the expectation yields

$$\begin{aligned} \mathbb{E}(u_{\ell'}(T), \varphi) &= \mathbb{E}(u_{\ell'}(0), \varphi) \\ &\quad + \mathbb{E}\langle (Ga_{\ell'}(\bar{u}_{\ell'}))(T), \varphi \rangle + \mathbb{E}((HB\bar{u}_{\ell'}\tilde{\Pi}_{m_{\ell'}})(T), \varphi). \end{aligned}$$

Let $\ell' \rightarrow \infty$. The weak-weak continuity of the operators G and H , together with Lemma 4.1, implies that

$$\mathbb{E}(\xi, \varphi) = \mathbb{E}(u_0, \varphi) + \mathbb{E}\langle (Ga^\infty)(T), \varphi \rangle + \mathbb{E}\langle (Hb^\infty)(T), \varphi \rangle. \quad (4.5)$$

Letting $M \rightarrow \infty$ and again using the limited completeness of the Galerkin scheme shows that the above equality holds for any \mathcal{F}_T -measurable $\varphi \in L^2(\Omega; V)$. If one now applies Itô's formula to (4.4) then one obtains an adapted process \tilde{u} with paths in $C([0, T]; H)$ that is equal to u almost surely. Furthermore, for any $\varphi \in L^2(\Omega; V)$ and due to continuity of \tilde{u} ,

$$\begin{aligned} \mathbb{E}(\xi - \tilde{u}(T), \varphi) &= \lim_{t \rightarrow T} \mathbb{E}(\xi - \tilde{u}(t), \varphi) \\ &= \lim_{t \rightarrow T} \mathbb{E} \left\langle \int_t^T a^\infty(s) ds + \int_t^T b^\infty(s) dW(s), \varphi \right\rangle = 0. \end{aligned}$$

Thus $\xi = \tilde{u}(T)$. This together with (4.5) implies (4.2). \square

All that remains to be done to prove Theorem 2.3 is to identify a^∞ with Au and b^∞ with Bu and to show strong convergence of $u_\ell(T)$ to $u(T)$. To that end we would like to use monotonicity of A . In order to overcome the difficulty arising from the fact that the tamed operator $A_{2,\ell}$ does not preserve the monotonicity property of A_2 we need the following lemma.

Lemma 4.3. *Let the Growth and Coercivity conditions in Assumption 1 hold. Let Assumptions 2 and 3 be satisfied. Let $u_\ell(0) \rightarrow u_0$ in $L^{q_0}(\Omega; H)$. Let $\frac{c(m_\ell)}{n_\ell} \rightarrow 0$ as $\ell \rightarrow \infty$. Then*

$$\mathbb{E} \int_0^T \|A_2 \bar{u}_\ell(s) - A_{2,\ell} \bar{u}_\ell(s)\|_{V_2^*}^{p^*} ds \rightarrow 0 \quad \text{as } \ell \rightarrow \infty.$$

Proof. Consider some $M > 0$. Recall that T_ℓ is given by (2.5). Then

$$\begin{aligned} I_\ell &:= \mathbb{E} \int_0^T \|A_{2,\ell} \bar{u}_\ell(s) - A_2 \bar{u}_\ell(s)\|_{V_2^*}^{p^*} ds \\ &= \mathbb{E} \int_0^T (1 - T_\ell(\bar{u}_\ell(s)))^{p^*} \|A_2 \bar{u}_\ell(s)\|_{V_2^*}^{p^*} \mathbf{1}_{\{\|A_2 \bar{u}_\ell(s)\|_{V_2^*} \leq M\}} ds \\ &\quad + \mathbb{E} \int_0^T (1 - T_\ell(\bar{u}_\ell(s)))^{p^*} \|A_2 \bar{u}_\ell(s)\|_{V_2^*}^{p^*} \mathbf{1}_{\{\|A_2 \bar{u}_\ell(s)\|_{V_2^*} > M\}} ds \\ &=: I_{1,\ell,M} + I_{2,\ell,M}. \end{aligned}$$

It is observed that

$$\begin{aligned} I_{1,\ell,M} &\leq \mathbb{E} \int_0^T \frac{n_\ell^{-1/2} |\Pi_{m_\ell} A_2 \bar{u}_\ell(s)|}{1 + n_\ell^{-1/2} |\Pi_{m_\ell} A_2 \bar{u}_\ell(s)|} \|A_2 \bar{u}_\ell(s)\|_{V_2^*}^{p^*} \mathbf{1}_{\{\|A_2 \bar{u}_\ell(s)\|_{V_2^*} \leq M\}} ds \\ &\leq \mathbb{E} \int_0^T \frac{\tau_{n_\ell}^{1/2} T^{-1/2} c(m)^{1/2} M}{1 + n_\ell^{-1/2} |\Pi_{m_\ell} A_2 \bar{u}_\ell(s)|} M^{p^*} ds \leq (c(m) \tau_{n_\ell})^{1/2} T^{1/2} M^{1+p^*}. \end{aligned}$$

Recall that due to Corollary 3.4 one knows that

$$\mathbb{E} \int_0^T \|A_2 \bar{u}_\ell(s)\|_{V_2^*}^{p^*(1+\eta)} ds < c$$

with c independent of ℓ . Thus the sequence $(\|A_2 \bar{u}_\ell\|_{V_2^*}^{p^*})_{\ell \in \mathbb{N}}$ is uniformly integrable on $(0, T) \times \Omega$ with respect to $dt \times P$. Hence for any $\epsilon > 0$ there exists M such that $I_{2,\ell,M} < \epsilon/2$ for all ℓ . Finally, since $\frac{c(m_\ell)}{n_\ell} \rightarrow 0$ as $\ell \rightarrow \infty$, one can choose ℓ large such that $I_{1,\ell,M} < \epsilon/2$. \square

We now prove Theorem 2.2. This is needed to later show that the whole sequence of approximations converges rather than just a subsequence.

Proof of Theorem 2.2. Assume that u_1 and u_2 are two distinct solutions to (1.1) such that $u_1(0) = u_2(0) = u_0$. One would now like to apply Itô's formula for the square of the norm from Pardoux [13, Chapitre 2, Theoreme 5.2]. To that end one immediately observes that $u_1 - u_2 \in \mathcal{L}^2(V_1) \cap \mathcal{L}^p(V_2)$ and that $u_1(0) - u_2(0) = 0 \in L^2(\Omega; H)$. Moreover

$$\|Au_1 - Au_2\|_{\mathcal{L}^2(V_1^*) + \mathcal{L}^{p^*}(V_2^*)} = \|A_1u_1 - A_1u_2\|_{\mathcal{L}^2(V_1^*)} + \|A_2u_1 - A_2u_2\|_{\mathcal{L}^{p^*}(V_2^*)}.$$

Using the Growth assumption on A_1 one observes that

$$\begin{aligned} \mathbb{E} \int_0^T \|A_1u_1(s) - A_1u_2(s)\|_{V_1^*}^2 ds &\leq \mathbb{E} \int_0^T 2[\|A_1u_1(s)\|_{V_1^*}^2 + \|A_1u_2(s)\|_{V_1^*}^2] ds \\ &\leq K \mathbb{E} \int_0^T 2[(1 + \|u_1(s)\|_{V_1}^2) + (1 + \|u_2(s)\|_{V_1}^2)] ds \\ &\leq c(1 + \|u_1\|_{\mathcal{L}^2(V_1)}^2 + \|u_2\|_{\mathcal{L}^2(V_1)}^2) < \infty. \end{aligned}$$

Also, using the Growth assumption on A_2 one obtains

$$\mathbb{E} \int_0^T \|A_2u_1(s) - A_2u_2(s)\|_{V_2^*}^{p^*} ds \leq c(1 + \|u_1\|_{\mathcal{L}^p(V_2)}^p + \|u_2\|_{\mathcal{L}^p(V_2)}^p) < \infty.$$

Thus $Au_1 - Au_2 \in \mathcal{L}^2(V_1^*) \cup \mathcal{L}^{p^*}(V_2^*)$. Finally, using the Growth assumption on B one deduces that $Bu_1 - Bu_2 \in \mathcal{L}^2(L_2(U, H))$. Hence the aforementioned Itô's formula for the square of the norm can be applied, yielding

$$\begin{aligned} e^{-Kt}|u_1(t) - u_2(t)|^2 &= -K \int_0^t e^{-Ks}|u_1(s) - u_2(s)|^2 ds \\ &\quad + \int_0^t e^{-Ks} [2\langle Au_1(s) - Au_2(s), u_1(s) - u_2(s) \rangle \\ &\quad + \|Bu_1(s) - Bu_2(s)\|_{L_2(U, H)}^2] ds + M(t), \end{aligned}$$

where

$$M(t) := \int_0^t e^{-Ks} (u_1(s) - u_2(s), (Bu_1(s) - Bu_2(s)) dW(s)).$$

One then observes, due to the monotonicity of $A : V \times \Omega \rightarrow V^*$, that

$$e^{-Kt}|u_1(t) - u_2(t)|^2 \leq M(t)$$

and hence M is non-negative. It is also a real-valued continuous local martingale, and thus a supermartingale. Furthermore it starts from 0 and thus, almost surely, $M(t) = 0$ for all $t \in [0, T]$. One thus concludes that $u_1(t) = u_2(t)$ for all $t \in [0, T]$ almost surely. \square

Finally we can prove Theorem 2.3.

Proof of Theorem 2.3. Recall that $a_\ell(v) := \Pi_{m_\ell}[A_1v + A_{2,\ell}v]$. Applying Itô's formula to the scheme (2.2) and taking expectations yields

$$\begin{aligned} e^{-KT} \mathbb{E}|u_{\ell'}(T)|^2 &= \mathbb{E}|u_{\ell'}(0)|^2 - K \mathbb{E} \int_0^T e^{-Ks}|u_{\ell'}(s)|^2 ds \\ &\quad + \mathbb{E} \int_0^T e^{-Ks} \left(2\langle a_{\ell'}(\bar{u}_{\ell'}(s)), \bar{u}_{\ell'}(s) \rangle + \|\Pi_{m_{\ell'}} B \bar{u}_{\ell'}(s)\|_{L_2(U, H)}^2 \right) ds \\ &\quad + \mathbb{E} \int_0^T 2e^{-Ks} \langle a_{\ell'}(\bar{u}_{\ell'}(s)), u_{\ell'}(s) - \bar{u}_{\ell'}(s) \rangle ds. \end{aligned}$$

Let

$$I_{1,\ell'} := \mathbb{E} \int_0^T \langle a_{\ell'}(\bar{u}_{\ell'}(s)), u_{\ell'}(s) - \bar{u}_{\ell'}(s) \rangle ds.$$

Using Hölder's inequality results in

$$I_{1,\ell'} \leq \left(\mathbb{E} \int_0^T |a_{\ell'}(\bar{u}_{\ell'}(s))|^2 ds \right)^{1/2} \left(\mathbb{E} \int_0^T |u_{\ell'}(s) - \bar{u}_{\ell'}(s)|^2 ds \right)^{1/2}.$$

Using Assumption 2 and Corollary 3.4 yields

$$I_{1,\ell'} \leq c(\mathbf{c}(m_{\ell'})\tau_{n_{\ell'}})^{1/2}.$$

Thus,

$$\begin{aligned} e^{-Ks} \mathbb{E}|u_{\ell'}(T)|^2 &\leq \mathbb{E}|u_{\ell'}(0)|^2 - K \mathbb{E} \int_0^T e^{-Ks} |u_{\ell'}(s)|^2 ds \\ &+ \mathbb{E} \int_0^T e^{-Ks} (2\langle a_{\ell'}(\bar{u}_{\ell'}(s)), \bar{u}_{\ell'}(s) \rangle + \|B\bar{u}_{\ell'}(s)\|_{L_2(U,H)}^2) ds + c(\mathbf{c}(m_{\ell'})\tau_{n_{\ell'}})^{1/2} \end{aligned}$$

and one may proceed with a monotonicity argument. Let $w \in \mathcal{L}^p(V)$. Then

$$\begin{aligned} e^{-KT} \mathbb{E}|u_{\ell'}(T)|^2 &\leq \mathbb{E}|u_{\ell'}(0)|^2 - K \mathbb{E} \int_0^T e^{-Ks} |u_{\ell'}(s)|^2 ds \\ &+ \mathbb{E} \int_0^T e^{-Ks} [2\langle a_{\ell'}(\bar{u}_{\ell'}(s)) - a_{\ell'}(w(s)), \bar{u}_{\ell'}(s) - w(s) \rangle \\ &\quad + 2\langle a_{\ell'}(w(s)), \bar{u}_{\ell'}(s) - w(s) \rangle + 2\langle a_{\ell'}(\bar{u}_{\ell'}(s)), w(s) \rangle \\ &\quad + 2(Bw(s), B\bar{u}_{\ell'}(s))_{L_2(U,H) \times L_2(U,H)} - \|Bw(s)\|_{L_2(U,H)}^2 \\ &\quad + \|B\bar{u}_{\ell'}(s) - Bw(s)\|_{L_2(U,H)}^2] ds + c(\mathbf{c}(m_{\ell'})\tau_{n_{\ell'}})^{1/2}. \end{aligned}$$

Using the Monotonicity assumption on A one obtains

$$\begin{aligned} e^{-KT} \mathbb{E}|u_{\ell'}(T)|^2 &\leq \mathbb{E}|u_{\ell'}(0)|^2 \\ &+ K \mathbb{E} \int_0^T e^{-Ks} (|\bar{u}_{\ell'}(s)|^2 - |u_{\ell'}(s)|^2 - 2(\bar{u}_{\ell'}(s), w(s)) + |w(s)|^2) ds \\ &+ \mathbb{E} \int_0^T e^{-Ks} \langle A_{2,\ell'} \bar{u}_{\ell'}(s) - A_2 \bar{u}_{\ell'}(s), \bar{u}_{\ell'}(s) - w(s) \rangle ds \\ &+ \mathbb{E} \int_0^T e^{-Ks} \langle A_2 w(s) - A_{2,\ell'} w(s), \bar{u}_{\ell'}(s) - w(s) \rangle ds \tag{4.6} \\ &+ \mathbb{E} \int_0^T e^{-Ks} [2\langle a_{\ell'}(w(s)), \bar{u}_{\ell'}(s) - w(s) \rangle \\ &\quad + 2\langle a_{\ell'}(\bar{u}_{\ell'}(s)), w(s) \rangle \\ &\quad + 2(Bw(s), B\bar{u}_{\ell'}(s))_{L_2(U,H) \times L_2(U,H)} - \|Bw(s)\|_{L_2(U,H)}^2] ds \\ &+ c(\mathbf{c}(m_{\ell'})\tau_{n_{\ell'}})^{1/2}. \end{aligned}$$

Taking limit inferior as $\ell' \rightarrow \infty$, using the weak lower-semi-continuity of the norm, Lemma 4.1, Lemma 4.3 and Corollary 3.4, one observes that

$$\begin{aligned} e^{-KT} \mathbb{E}|u(T)|^2 &\leq \mathbb{E}|u_0|^2 + K \mathbb{E} \int_0^T e^{-Ks} [-2(u(s), w(s)) + |w(s)|^2] ds \\ &+ \mathbb{E} \int_0^T e^{-Ks} [2\langle Aw(s), u(s) - w(s) \rangle + 2\langle a^\infty(s), w(s) \rangle \\ &\quad + 2(Bw(s), b^\infty(s))_{L_2(U,H) \times L_2(U,H)} - \|Bw(s)\|_{L_2(U,H)}^2] ds. \end{aligned} \tag{4.7}$$

Applying Itô's formula to (4.1) and taking expectations yields

$$\begin{aligned} e^{-KT} \mathbb{E}|u(T)|^2 &= \mathbb{E}|u(0)|^2 - K \mathbb{E} \int_0^T e^{-Ks} |u(s)|^2 ds \\ &\quad + \mathbb{E} \int_0^T e^{-Ks} [2\langle a^\infty(s), u(s) \rangle + \|b^\infty(s)\|_{L_2(U,H)}^2] ds \end{aligned} \quad (4.8)$$

Subtracting this from (4.7) one arrives at

$$\begin{aligned} 0 \leq \mathbb{E} \int_0^T e^{-Ks} [K|u(s) - w(s)|^2 + 2\langle Aw(s), u(s) - w(s) \rangle \\ + 2\langle a^\infty(s), w(s) - u(s) \rangle - \|Bu(s) - b^\infty(s)\|_{L_2(U,H)}^2] ds. \end{aligned} \quad (4.9)$$

Note that so far w was arbitrary. It will now be used to identify the nonlinear terms. First, one takes $w = u$ and observes that,

$$0 \leq -\mathbb{E} \int_0^T e^{-Ks} \|Bu(s) - b^\infty(s)\|_{L_2(U,H)}^2 ds \leq 0$$

which implies $b^\infty = Bu$. Next, one sets $w = u + \epsilon z$ with $\epsilon > 0$ and $z \in \mathcal{L}^p(V)$ in (4.9). Dividing by $\epsilon > 0$ leads to

$$0 \leq \mathbb{E} \int_0^T e^{-Ks} [K\epsilon|z|^2 + 2\langle A(u(s) + \epsilon z(s)), -z(s) \rangle + 2\langle a^\infty(s), z(s) \rangle] ds.$$

Using hemicontinuity of A while letting $\epsilon \rightarrow 0$ results in

$$\mathbb{E} \int_0^T \langle Au(s), z(s) \rangle ds \leq \mathbb{E} \int_0^T \langle a^\infty(s), z(s) \rangle ds.$$

This holds for an arbitrary $z \in \mathcal{L}^p(V)$ and hence, in particular, for $-z$. Thus one obtains that $a^\infty = Au$.

Due to Theorem 2.2, the solution u to (1.1) is unique. Thus the whole sequences of approximations converges rather than just the subsequence denoted by ℓ' .

Finally, in order to show that $u_\ell(T) \rightarrow u(T)$ in $L^2(\Omega; H)$, one uses (4.6) and (4.7) with $w = u$ together with $a^\infty = Au$ and $b^\infty = Bu$. Consequently, the weak-lower-semi-continuity of the norm and Lemma 4.1 lead to

$$\begin{aligned} e^{-KT} \mathbb{E}|u(T)|^2 &\leq \liminf_{\ell \rightarrow \infty} e^{-KT} \mathbb{E}|u_\ell(T)|^2 \leq \mathbb{E}|u_0|^2 - K \mathbb{E} \int_0^T e^{-Ks} |u(s)|^2 ds \\ &\quad + \mathbb{E} \int_0^T e^{-Ks} [2\langle Au(s), u(s) \rangle + \|Bu(s)\|_{L_2(U,H)}^2] ds. \end{aligned}$$

Thus, due to (4.8),

$$0 \leq \liminf_{\ell \rightarrow \infty} \mathbb{E}|u_\ell(T)|^2 - \mathbb{E}|u(T)|^2 \leq 0.$$

From Lemma 4.1 and Lemma 4.2, one already knows that $u_\ell(T) \rightharpoonup u(T)$ in $L^2(\Omega; H)$. This is a uniformly convex space (as it is a Hilbert space). Thus one concludes that $u_\ell(T) \rightarrow u(T)$ in $L^2(\Omega; H)$. For this see, e.g., Brézis [2, Proposition 3.32]. \square

5. EXAMPLES

In this section we give examples of three equations which fit into our framework. In all three examples the interpolation inequality is a consequence of the Gagliardo–Nirenberg inequality (see, for example, [15, Theorem 1.24]). The first example is the equation:

$$du = [\nabla a(\nabla u) - |u|^{p-2}u] dt + u dW \quad \text{on } \mathcal{D} \times (0, T)$$

with $u = 0$ on the boundary of the domain \mathcal{D} and $u(\cdot, 0) = u_0$ given. Here $a : \mathbb{R}^d \rightarrow \mathbb{R}^d$ can be nonlinear but it is assumed to be continuous, monotone and growing at most linearly. If we take $a_i(z) = z_i$ then $\nabla a(\nabla u) = \Delta u$ and this equation is the stochastic Ginzburg–Landau equation. An example of a nonlinear function is $a_i(z) = \frac{2 + \exp(-z_i)}{1 + \exp(-z_i)}$. Moreover \mathcal{D} is a bounded Lipschitz domain in \mathbb{R}^d , $d = 1, 2, 3$ and $p \in [2, 6)$ if $d = 1$, $p \in [2, 4)$ if $d = 2$ and $p \in [2, 10/3)$ if $d = 3$. In our framework $H = L^2(\mathcal{D})$, $V_1 = H_0^1(\mathcal{D})$ and $V_2 = L^p(\mathcal{D})$ (using the standard notation for Lebesgue and Sobolev spaces).

The second is the stochastic Swift–Hohenberg equation:

$$du = [(\gamma^2 - (1 + \Delta)^2)u - |u|^{p-2}u]dt + dW \quad \text{on } \mathcal{D} \times (0, T)$$

with appropriate boundary and initial conditions. The domain \mathcal{D} is assumed to be a bounded Lipschitz domain in \mathbb{R}^2 . With Dirichlet boundary conditions we would take $V_1 = H_0^2(\mathcal{D})$ and $V_2 = L^p(\mathcal{D})$ with $p \in [2, 6)$.

The third example is the spatially extended stochastic FitzHugh–Nagumo system for signal propagation in nerve cells (originally stated by FitzHugh [3] as a system of ordinary differential equations, see Bonaccorsi and Mastrogiacomo [1] for mathematical analysis of the spatially extended stochastic version):

$$\begin{aligned} du &= (\Delta u + u - u^3 - v)dt + dW \\ dv &= c_1(u - c_2v + c_3)dt \end{aligned} \quad \text{on } (0, 1) \times (0, T),$$

together with appropriate initial data for u and v as well as homogeneous Neumann boundary conditions for u only. In this situation $V_1 = H^1((0, 1)) \times L^2((0, 1))$ while $V_2 = L^4((0, 1)) \times L^2((0, 1))$.

We now provide estimates on the constant $\mathfrak{c}(m)$ in the particular case when $\mathcal{D} = (0, \pi)^2 \subset \mathbb{R}^2$ and we use a spectral Galerkin method to construct the spaces V_m . To that end define

$$\varphi_{n_1 n_2}(x_1, x_2) := \frac{2}{\pi} \sin(n_1 x_1) \sin(n_2 x_2).$$

Let $V_m = \text{span}\{\varphi_{n_1 n_2} : n_1 = 1, \dots, m, n_2 = 1, \dots, m\}$. Then

$$\Pi_m f := \sum_{n_1=1, n_2=1}^m \langle f, \varphi_{n_1 n_2} \rangle \varphi_{n_1 n_2}$$

satisfies Assumption 2. Moreover we can calculate

$$\begin{aligned} \mathfrak{c}(m) &= \sum_{n_1=1, n_2=1}^m \left(\|\varphi_{n_1 n_2}\|_{L^2(\mathcal{D})}^2 + \|\nabla \varphi_{n_1 n_2}\|_{L^2(\mathcal{D}; \mathbb{R}^2)}^2 + \|\varphi_{n_1 n_2}\|_{L^p(\mathcal{D})}^{2/p} \right) \\ &= m^2 (1 + 2 + c_p), \end{aligned}$$

where c_p depends only on p . Hence, in order to apply Theorem 2.3, we need a sequence (m_ℓ, n_ℓ, k_ℓ) such that $\frac{m_\ell^2}{n_\ell} \rightarrow 0$ as $\ell \rightarrow \infty$. This means that we need to choose $n_\ell = \lfloor m_\ell^{2+\delta} \rfloor$ for some $\delta > 0$.

We also note that if $\mathcal{D} = (0, \pi)^d$ then an analogous construction of V_m would lead to the conclusion that we need $n_\ell = \lfloor m_\ell^{d+\delta} \rfloor$ for some $\delta > 0$. Crucially we see that the space-time coupling requirement is no more onerous than in the case of equations with operators growing at most linearly.

ACKNOWLEDGEMENT

The authors would like to thank the referees for their comments on the paper.

REFERENCES

- [1] S. Bonaccorsi and E. Mastrogiacomo. Analysis of the Stochastic FitzHugh–Nagumo System. *Infin. Dimens. Anal. Quantum Probab. Relate. Top.*, 11(03):427–446, 2008.
- [2] H. Brézis. *Functional analysis, Sobolev spaces and partial differential equations*. Springer, New York, 2010.
- [3] R. FitzHugh. Impulses and Physiological States in Theoretical Models of Nerve Membrane. *Biophys. J.*, 1:445–466, 1961.
- [4] H. Gajewski, K. Gröger, and K. Zacharias. *Nichtlineare Operatorgleichungen und Operatordifferentialgleichungen*. Akademie-Verlag, Berlin, 1974.
- [5] M.B. Giles. Multilevel Monte Carlo path simulation. *Oper. Res.*, 56:607–617, 2008.
- [6] I. Gyöngy and A. Millet. On discretization schemes for stochastic evolution equations. *Potential Anal.*, 23(2):99–134, 2005.
- [7] S. Heinrich. Monte Carlo complexity of global solution of integral equations. *J. Complexity*, 14(2):151–175, 1998.
- [8] S. Heinrich. Multilevel Monte Carlo methods. In *Large-Scale Scientific Computing*, vol. 2179, pp. 58–67 of *Lect. Notes Comput. Sci.*, 2001.
- [9] M. Hutzenthaler and A. Jentzen and P.E. Kloeden. Strong and weak divergence in finite time of Euler’s method for stochastic differential equations with non-globally Lipschitz continuous coefficients. *Proc. R. Soc. A* 467:1563–1576, 2011.
- [10] M. Hutzenthaler and A. Jentzen and P.E. Kloeden. Strong convergence of an explicit numerical method for SDEs with non-globally Lipschitz continuous coefficients. *Ann. Appl. Probab.*, 22:1611–1641, 2012.
- [11] M. Hutzenthaler and A. Jentzen and P.E. Kloeden. Divergence of the multilevel Monte Carlo Euler method for nonlinear stochastic differential equations. *Ann. Appl. Probab.*, 23(5):1913–1966, 2013.
- [12] M. Hutzenthaler and A. Jentzen. Numerical Approximations of stochastic differential equations with non-globally Lipschitz coefficients. *Mem. Amer. Math. Soc.* 4:1–112, 2015.
- [13] É. Pardoux. Sur des équations aux dérivées partielles stochastiques monotones. *C. R. Acad. Sci. Paris Sér. A-B*, 275:A101–A103, 1972.
- [14] G. O. Roberts and R. L. Tweedie. Exponential convergence of Langevin distributions and their discrete approximations. *Bernoulli*, 2(4):341–363.
- [15] T. Roubíček. *Nonlinear partial differential equations with applications*. Birkhäuser, Basel, 2005.
- [16] S. Sabanis. A note on tamed Euler approximations. *Electron. Commun. Probab.*, 18(47):1–10, 2013.
- [17] S. Sabanis. Euler approximations with varying coefficients: the case of superlinearly growing diffusion coefficients. *arXiv:1308.1796 [math.PR]*, 2015.
- [18] K. Dareiotis, C. Kumar and S. Sabanis. On Tamed Euler Approximations of SDEs Driven by Lévy Noise with Applications to Delay Equations. *arXiv:1403.0498 [math.PR]*, 2015.
- [19] M. V. Tretyakov and Z. Zhang. A fundamental mean-square convergence theorem for SDEs with locally Lipschitz coefficients and its applications. *SIAM J. Numer. Anal.*, 51(6):3135–3162, 2013.