



THE UNIVERSITY *of* EDINBURGH

Edinburgh Research Explorer

Large-scale screening and characterization of enteroviruses and kobuviruses infecting pigs in Vietnam

Citation for published version:

Nguyen, DV, Anh, PH, Cuong, NV, Hoa, NT, Carrique-Mas, J, Hien, VB, Sharp, C, Rabaa, M, Berto, A, Campbell, J, Baker, S, Farrar, J, Woolhouse, ME, Bryant, JE & Simmonds, P 2016, 'Large-scale screening and characterization of enteroviruses and kobuviruses infecting pigs in Vietnam' *Journal of General Virology*, vol. 97, no. 2, pp. 378-388. DOI: 10.1099/jgv.0.000366

Digital Object Identifier (DOI):

[10.1099/jgv.0.000366](https://doi.org/10.1099/jgv.0.000366)

Link:

[Link to publication record in Edinburgh Research Explorer](#)

Document Version:

Publisher's PDF, also known as Version of record

Published In:

Journal of General Virology

General rights

Copyright for the publications made accessible via the Edinburgh Research Explorer is retained by the author(s) and / or other copyright owners and it is a condition of accessing these publications that users recognise and abide by the legal requirements associated with these rights.

Take down policy

The University of Edinburgh has made every reasonable effort to ensure that Edinburgh Research Explorer content complies with UK legislation. If you believe that the public display of this file breaches copyright please contact openaccess@ed.ac.uk providing details, and we will remove access to the work immediately and investigate your claim.



Large-scale screening and characterization of enteroviruses and kobuviruses infecting pigs in Vietnam

Nguyen Van Dung,¹ Pham Hong Anh,² Nguyen Van Cuong,² Ngo Thi Hoa,^{2,5} Juan Carrique-Mas,² Vo Be Hien,³ C. Sharp,¹ M. Rabaa,² A. Berto,² James Campbell,² Stephen Baker,^{2,5,6} Jeremy Farrar,² Mark E. Woolhouse,⁴ Juliet E. Bryant^{2,5} and Peter Simmonds^{1,4}

Correspondence

Peter Simmonds

peter.simmonds@ed.ac.uk

¹Infection and Immunity Division, Roslin Institute, University of Edinburgh, Easter Bush, Edinburgh EH25 9RG, UK

²Oxford University Clinical Research Unit, 764 Vo Van Kiet, W.1, District 5, Ho Chi Minh City, Vietnam

³Subdepartment of Animal Health, Dong Thap Province, Vietnam

⁴Centre for Immunity, Infection and Evolution, University of Edinburgh, Ashworth Laboratories, Kings Buildings, West Mains Road, Edinburgh EH9 3JT, UK

⁵Nuffield Department of Medicine, Oxford University, Oxford OX3 7BN, UK

⁶London School of Hygiene and Tropical Medicine, Keppel Street, Bloomsbury, London WC1E 7HT, UK

A recent survey of pigs in Dong Thap province, Vietnam identified a high frequency of enterovirus species G (EV-G) infection (144/198; 72.7 %). Amongst these was a plethora of EV-G types (EV-G1, EV-G6 and four new types EV-G8–EV-G11). To better characterize the genetic diversity of EV-G and investigate the possible existence of further circulating types, we performed a larger-scale study on 484 pig and 45 farm-bred boar faecal samples collected in 2012 and 2014, respectively. All samples from the previous and current studies were also screened for kobuviruses. The overall EV infection frequency remained extremely high (395/484; 81.6 %), but with comparable detection rates and viral loads between healthy and diarrhoeic pigs; this contrasted with less frequent detection of EV-G in boars (4/45; 8.9 %). EV was most frequently detected in pigs ≤ 14 weeks old (~ 95 %) and declined in older pigs. Infections with EV-G1 and EV-G6 were most frequent, whilst less commonly detected types included EV-G3, EV-G4 and EV-G8–EV-G11, and five new types (EV-G12–EV-G16). In contrast, kobuvirus infection frequency was significantly higher in diarrhoeic pigs (40.9 versus 27.6 %; $P=0.01$). Kobuviruses also showed contrasting epizootologies and age associations; a higher prevalence was found in boars (42 %) compared with domestic pigs (29 %), with the highest infection frequency amongst pigs >52 weeks old. Although genetically diverse, all kobuviruses identified belonged to the species *Aichivirus C*. In summary, this study confirms infection with EV-G was endemic in Vietnamese domestic pigs and exhibits high genetic diversity and extensive inter-type recombination.

Received 11 October 2015

Accepted 8 December 2015

INTRODUCTION

Porcine enteroviruses (PEVs) are members of the family *Picornaviridae* and originally consisted of 13 types

The GenBank/EMBL/DDBJ accession numbers for the sequences obtained in this study are KT265880–KT266194.

One supplementary figure and three supplementary tables are available with the online Supplementary Material.

(PEV-1–PEV-13). Further studies on their genetic relatedness led to the reassignment of these viruses into three separate picornavirus genera: *Enterovirus*, *Sapelovirus* and *Teschovirus* (Kaku *et al.*, 2001; Knowles, 2006; Krumbholz *et al.*, 2002; Zell *et al.*, 2001). Of these, PEV-9 and PEV-10 were classified as EV-G1 and EV-G2 serotypes in enterovirus G, with the subsequently characterized EV-G types EV-G3–EV-G6 reported in single publications from Hungary and South Korea (Boros *et al.*, 2011, 2012a, b;

Moon *et al.*, 2012). EV-G7 was subsequently discovered from domestic sheep by A. Boros and N. J. Knowles (unpublished). More recently, our molecular characterization of EVs in a small number of PCR-positive faecal samples from domestic pigs identified a further four types (EV-G8–EV-11) within this species. The presence of such a large amount of genetic diversity within such a modest number of samples suggests that the full range of EV-G types in this virus species has yet to be fully characterized (Van Dung *et al.*, 2014). In contrast to teschoviruses, infections with EV-G have not been linked to any enteric or other disease presentations (Knowles, 2006) except for skin lesions (Knowles, 1988) and flaccid paralysis in an experimental infection of pigs in China (Yang *et al.*, 2013). Viruses in this species have been reported from both healthy and diarrhoeic pigs with no statistically significant difference in infection rates between the two groups (Van Dung *et al.*, 2014).

Porcine kobuvirus (PKV) was first identified in 2008 in Hungary (Reuter *et al.*, 2008). Whilst screening 15 faecal samples from pigs for calicivirus (norovirus and sapovirus) using reverse transcription-PCR, Reuter *et al.* (2008) detected, in addition to a band specific for sapovirus, a non-specific, strong band of ~1.1 kb in all tested samples visible on agarose gel. The nucleotide sequence of the non-specific PCR product showed most similarity to the 3C/3D region of bovine and human aichiviruses. Upon analysis of the obtained complete nucleotide sequence, the new PKV was assigned as a member of a new species (*Aichivirus C*) in the genus *Kobuvirus*, which also includes the species *Aichivirus A* and *Aichivirus B*, originally detected in humans and oxen, respectively. The species *Aichivirus C* consists of a single type, PKV-1. The existence of PKV-1 in pigs has been well documented in different countries worldwide (Amimo *et al.*, 2014; Barry *et al.*, 2011; Chen *et al.*, 2013; Di Bartolo *et al.*, 2015; Fan *et al.*, 2013; Khamrin *et al.*, 2009, 2010; Okitsu *et al.*, 2012; Park *et al.*, 2010; Reuter *et al.*, 2013; Sisay *et al.*, 2013); however, in Vietnam, PKV-1 has only been reported from river water (Inaba *et al.*, 2014).

We have performed this larger-scale screening study to better understand the prevalence, genetic diversity and

disease association of EVs and kobuviruses infecting pigs and farm-bred boars (hereafter referred to as 'boars') in Vietnam.

RESULTS

Detection of EVs

EV RNA was detected in all but one (103/104; 99 %) of the farms containing domestic pigs and in three from six (50 %) farms housing boars. In total, 395 of 484 (81.6 %) samples from domestic pigs and four of 55 (8.9 %) boar samples were PCR-positive (Table 1). Pigs ≤14 weeks old were the most frequently infected with faecal detection frequencies of nearly 95 %. Lower detection frequencies were observed in older pigs with the lowest rate (50.5 %) in pigs >1 year old. Combining screening data with that of our previous samples, we found no significant difference in the frequency of detection of EV infection between the two groups in all age ranges (Table 1). The lack of any direct causative role of EV-G in enteric disease in pigs was further demonstrated by our observation that pigs with diarrhoea showed slightly lower EV-G RNA viral loads than found in healthy pigs ($P=0.008$, Mann–Whitney test; Fig. 1).

Detection of kobuviruses

The overall detection frequencies of kobuviruses in domestic pigs and boars were 29.3 (200/682) and 42.2 % (19/45), respectively. In contrast to EV-G, kobuviruses were significantly associated with diarrhoea in pigs, with an infection frequency of 40.9 % (36/88) in diarrhoeic pigs and an infection frequency of 27.6 % (164/594) in asymptomatic pigs ($P=0.01$; χ^2 -test). This positive association was, however, most marked in pigs >52 weeks old in which the highest prevalence (67.9 %) was recorded. For pigs ≤23 weeks old, infection frequencies were highest in sucklers and generally decreased with age.

In order to elucidate whether PKV viral loads were associated with the clinical status of pigs, a real-time PCR

Table 1. Detection of EVs and PKVs in pigs

Pigs	Age (weeks)	No. of pigs tested*	EV prevalence (%)*	P-value†	PKV prevalence (%)*	P-value†	EV/PKV co-infection (%)*	P-value†
Suckler	<7	11/78	90.9/93.6	0.56	63.6/59	1	63.6/53.8	0.75
Weaner	7–14	32/147	93.8/94.6	1	18.8/25.9	0.4	18.8/24.5	0.49
Grower	15–23	8/117	75/82.9	0.63	25/16.2	0.62	25/13.7	0.32
Gilts	24–52	0/68	–/70.6	–	–/26.5	–	–/17.6	–
Matured	>52	28/106	39.3/48.1	0.4	67.9/25.5	2.7×10^{-5}	28.6/13.2	0.08
Total	–	79/516‡	72.2/79.1	0.17	43/28.7	0.01	29.1/23.3	0.26

*Data are presented in pairs of diarrhoeic/non-diarrhoeic samples.

†Fisher's exact test or χ^2 -test.

‡Only samples from the previous and current studies with available data (age and clinical status) are included.




Fig. 1. Box plot of EV RNA viral load of EV-G screen-positive sample suspensions from healthy and diarrhoeic pigs. Plots represent median and interquartile range (with outliers). The Mann–Whitney test was applied to compare medians between groups.

protocol was developed. The assay had a detection limit of 100 PKV RNA copies per reaction and an amplification efficiency of 89 % (data not shown). It showed good specificity (no false positives in 20 samples negative by nested PCR) and sensitivity (97/100 samples positive by nested PCR were detected by real-time PCR). The median \log_{10} viral load (7.7 copies ml^{-1}) of PKV in 41 matured pig samples was significantly higher than that (6.4 copies ml^{-1}) in the group of 56 younger pig samples ($P=0.0003$, Mann–Whitney test; Fig. 2a). However, there was no significant difference in the viral loads (median \log_{10} viral loads of 7.99 and 7.56 copies ml^{-1} , respectively) between matured pigs with and without diarrhoea ($P=0.22$ by Mann–Whitney test; Fig. 2b).

The differences in the co-infection rates were not significant in any age categories between the two groups of diarrhoeic and non-diarrhoeic pigs ($P>0.05$). Co-infection with EV and PKV was most common in the youngest pigs (Table 1).

Genetic characterization of porcine EVs

A range of EV variants in a subsample of 177 positive samples representing all sampling districts, C_t value

ranges, ages and clinical status of infected pigs, and four boar positive samples were typed by sequencing of the VP1 region. All VP1 sequences obtained from 130 samples (including four from boars) clustered with the species G EVs (Fig. 3). Phylogenetic and sequence distance analysis of this region showed the most commonly identified previously described type was EV-G1 ($n=57$ sequences, three boar-derived), followed by EV-G6 ($n=29$), EV-G11 ($n=9$), EV-G4, EV-G8 and EV-G9 ($n=6$ of each type), EV-G3 ($n=5$) and EV-G10 ($n=2$). The other 10 VP1 nucleotide sequences were substantially divergent ($>25\%$) from other EV-G types and have been assigned as new types EV-G12–EV-G16 by the International Committee on Taxonomy of Viruses Picornavirus Study Group (Fig. 3).

Analysis of data shows that young pigs were more frequently infected with EV-G1 and EV-G6 compared with other EV-G types, with median ages of 2 and 1.75 months, respectively. Meanwhile, other types were generally found in older pigs (median age 4 months) (Table S1, available in the online Supplementary Material). These differences were statistically significant ($P<0.0001$ and $P=0.0001$ by Mann–Whitney test; Fig. 4).

Sequence and phylogenetic analysis of PKVs

In total, 184 VP1 sequences were obtained from 173 positive samples, including 15 from boars (Fig. 5). Co-infection with two kobuvirus variants was detected in 11 samples by sequencing of VP1. All the VP1 sequences in the current study clustered within aichivirus C with nucleotide and amino acid sequence distances ranging from 0 to 22.3 % (mean 14.3 %) and 0 to 17.6 % (mean 8.5 %), respectively. In comparison with VP1 sequences of aichivirus C available in GenBank, the novel sequences produced here showed 9.1–23.6 and 0.45–17.5 % nucleotide and amino acid distances, respectively. In the resulting phylogeny, sequences from Vietnam were segregated into six groups (groups 1–6) with sequences from boars representing five distinct lineages, whilst those from domestic pigs clustering in three large groups. Groups 1–3 appeared to be most closely related to sequences from China (GenBank accession numbers KJ452348, KF539763 and KF157926), whilst groups 4–6 were most closely related to sequences from the USA (GenBank accession number JX987506), Hungary (GenBank accession number GQ249161.1) and Thailand (GenBank accession number AB624490), respectively. To test whether specific groups were associated with diarrhoea, association index values were calculated using genetic groups 1, 3 and 6 for group assignments and diarrhoea/non-diarrhoea as the associated variable. The calculated association index values of 0.45, 0.85 and 0.66 for the three groups, respectively, provided no evidence for a difference in clinical outcomes between groups.




Fig. 2. PKV RNA viral load of pig sample suspensions in (a) different age groups and (b) healthy or diarrhoeic matured pigs. Bars show median values. The Mann–Whitney test was applied to compare medians between groups.

Recombination of new types EV-G8 and EV-9

To perform a recombination analysis, complete coding region sequences of two new types were obtained by overlapping PCR amplifications. Phylogenetic relationships between EV-G8 and EV-G9 and other EV-G types for which complete genome sequences were available were inconsistent in several different regions of the genome (5' UTR, VP4/VP2, VP1 and 3D), indicative of recombination (Fig. 6). Most obviously, whilst sequences from the 5' UTR and VP4/VP2 and VP1 regions of the two novel EV-G types (EV-G8 and EV-G9) clustered separately from all other known EV-G types, sequences from the 3D regions of these novel viruses clustered together and formed a well-supported monophyletic clade (Fig. 6). To identify sites where recombination was likely to have occurred, a nucleotide distance scan (Fig. 7a) and grouping scan (Fig. 7b) were performed using EV-G9 as the reference sequence. EV-G9 was equally divergent from EV-G8 in P1 as it was from other types, but divergence values between EV-G8 and EV-G9 plunged at the boundary of P1/P2 to values equal to or lower than the mean pairwise distance of sequences within EV-G9. The analysis localized the breakpoint to the middle of the 2A region.

DISCUSSION

This is the first large-scale study of the prevalence and genetic diversity of EVs and kobuviruses in pigs and boars,

providing substantial new information on their high infection frequencies and co-circulation of multiple types and lineages of both viruses in domestic pigs and boars in Vietnam.

The overall detection frequency of EVs in domestic pigs (81.6 %) was notably higher than those reported in Spain (0 %), Italy (7.5 %), China (8.3 %) and the Czech Republic (50.2 %) (Buitrago *et al.*, 2010; Prodělalová, 2012; Sozzi *et al.*, 2010; Yang *et al.*, 2013). In contrast, detection of EV RNA in boars in this study was much lower than the 50 and 69.4 % prevalence reported in Hungary (Boros *et al.*, 2012a) and the Czech Republic (Prodělalová, 2012), respectively. High rates of EV infection in young pigs (detection rate >90 %) suggested that these viruses maintain endemic transmission in this population. Infection rates declined with the age of the pigs, to 50.5 % in matured pigs, consistent with a limited period of active infection by and faecal excretion of circulating EV-G types, followed by clearance and likely immunity to reinfection. The greater likelihood of encountering the most actively circulating EV-G types (EV-G1 and EV-G6) early in life may account for their increased detection in younger pigs (Fig. 4). As a result of the more limited circulation of other EV-G types, first exposure may be delayed and therefore account for the increased age range of pigs in which these types are detected.

Even though domestic pigs and boars are typically reared separately, we found that EV-G1 infected both animal




Fig. 3. Phylogenetic comparisons of VP1 sequences (nt 2469–3317) from study samples and available sequences of other EV-G variants from GenBank. Maximum-likelihood phylogenetic trees were reconstructed using 1000 bootstrap resamples to demonstrate the robustness of groupings; bootstrap support values of $\geq 70\%$ are shown. The tree was rooted using bovine enterovirus (BEV; GenBank accession number AF123433) as outgroup. The tree was drawn to scale; bar, estimated number of substitutions per site.

populations. Furthermore, variants were phylogenetically interspersed, indicating transmission events between the two populations. How this occurs is currently unclear as the boars were sampled in farms without domestic pigs and vice versa.

Our study provided no evidence that infections with EV-G caused diarrhoea, with comparable detection frequencies in pigs with and without symptoms (Table 1), consistent with a previous study (Prodělalová, 2012). To date, only EV-G1 virus has been specifically associated with disease in pigs, based on observations from the experimental infection of pigs in China in which two of 12 specific pathogen-free 2-week-old pigs exhibited flaccid paralysis of the hind

limbs with EV RNA detected in plasma, spinal cord and brain. In that experiment, the sample supernatant (not EV isolate) was filtered through $0.22\ \mu\text{m}$ microfilters before inoculation (Yang *et al.*, 2013; Zhang *et al.*, 2012). The sample was negative for a number of pathogens in pigs (e.g. porcine teschoviruses, porcine circovirus type 2, porcine reproductive and respiratory syndrome virus, classical swine fever virus, porcine epidemic diarrhoea virus, porcine rotavirus, hepatitis E virus). However, findings from our study indicate that this must be a rare outcome of infection given the absence of any evident neurological disease in pig farms where EV-G1 (and other EV types) circulate extensively.




Fig. 4. Box plot of EV types by pig ages. Note the y -axis is displayed in a \log_{10} format. Plots represent median and interquartile range (with outliers). The Mann–Whitney test was applied to compare medians between groups.

Genetic diversity and recombination in EV-G

Findings of four new types (EV-G8–EV-G11) from only 20 positive samples in our previous study suggested that the genetic diversity of EV-G had not been characterized completely (Van Dung *et al.*, 2014). This conclusion was supported by the detection of five more novel types from 130 positive samples in the current study. There is no doubt that more new EV-G types will continue to be detected from pigs from different countries in the future. To further characterize the genetic diversity of EV-G and the potential occurrence of recombination between types, almost complete genome sequences of EV-G8 and EV-G9 were obtained and compared with those of other EV-G types. Collectively, EV-Gs show generic characteristics, such as patterns of divergence, codon usage and non-synonymous to synonymous substitutions that occur in other picornaviruses prone to extensive recombination, including EV-A–EV-C (Van Dung *et al.*, 2014; Oberste *et al.*, 2004; Simmonds, 2006). Consistent with these predictions, recombination in the EV-G genome was detected, with breakpoints located at the border of the 5' UTR and VP4 of the EV-G5 prototype sequence (Boros *et al.*, 2012b). Our previous study additionally identified other

recombination occurring in the HM131607-G1 genome and other EV sequences from domestic pigs in Vietnam (Van Dung *et al.*, 2014). In the current study, sequences of EV-G8 and EV-G9 variants showed significant phylogenetic incongruence between 3D (in which EV-G8 and EV-G9 sequences grouped together) and other regions (5' UTR, VP4/VP2 and VP1, where EV-G8 clustered separately from EV-G9 sequences). This inconsistency in tree topology is strongly suggestive of recombination. Fragments 3 (nt 2395–3390) and 4 (nt 2679–5419) of the EV-G9 variants, both amplified by a single PCR (Fig. S1), had a long identical overlapping region that was distinct from the corresponding region of EV-G8. This helped rule out the possibility of hybrid viruses formed by assembly error from co-infection of individual pigs with two or more EV variants. When comparing the EV-G8 and EV-G9 variants, a nucleotide sequence divergence scan (Fig. 7a) showed considerable divergence in P1; from the middle of 2B to the end of the coding region, however, divergence declined substantially to levels even lower than that observed within EV-G9 sequences alone. The grouping scan (Fig. 7b), which quantifies degrees of grouping with pre-defined groups (types), similarly identified a recombination breakpoint in the middle of the 2A region.

Infection frequencies, diversity and disease associations of PKV

In accordance with previous studies (Chen *et al.*, 2013; Park *et al.*, 2010), our study showed an association between PKV detection and the presence of diarrhoea in matured pigs (Table 1), indicating its potential aetiological role in gastrointestinal disease, at least in older pigs. In marked contrast to EV infections, however, the highest detection rates were observed in diarrhoeic pigs >52 weeks old (matured pigs). This contrasts with reports from other countries where pigs <6 months old, particularly sucklers, were most susceptible to PKV infection (Chen *et al.*, 2013; Park *et al.*, 2010). Furthermore, we found viral loads of PKV in matured pigs to be significantly higher than that in the younger pigs.

This observation of increasing detection frequencies with age may simply reflect the possibility that PKV circulates less extensively than other enteric viruses, such as EV-G, and that infections are skewed towards older age groups with greater cumulative exposure. Another hypothesis to explain increasing detection frequencies and greater viral loads in older animals is that PKV infections are naturally persistent, and pigs may sequentially acquire long-term infections with one or more PKV variants over time. Although frequencies of persistence in PKV and other kobuviruses are poorly documented, genomes of all members of this genus possess structured RNA genomes (Davis *et al.*, 2008; Simmonds *et al.*, 2004) – a generic property associated with persistence in several other picornavirus genera and amongst other families of positive-stranded RNA viruses. High rates of detection in older pigs and




Fig. 6. Phylogenetic comparisons of sequences from the (a) 5' UTR (nt 107–809), (b) VP4/VP2 (nt 810–1752), (c) VP1 (nt 2469–3317) and (d) 3D (nt 5934–7316) regions from the study samples and available sequences of other EV-G variants from GenBank. The scale bar shows number of substitutions per site (evolutionary distance).

sequence distances from each other in VP1 (Oberste *et al.*, 1999), whilst different rhinovirus serotypes in the same genus indeed show divergences of ≥ 12 –13 % (McIntyre *et al.*, 2013). As PKV VP1 sequence distances lie between these two thresholds, prediction of their serological properties using this type of comparative approach is problematic. However, the detection of different PKV variants in the same samples is more consistent with them being antigenically distinct and lacking cross-protection, comparable to what is observed with different serotypes of poliovirus and foot-and-mouth disease virus (Alexandersen & Donaldson, 2002; Alexandersen *et al.*, 2002).

CONCLUSIONS

This large-scale study showed high prevalence and genetic diversity of EV-G and PKV in pigs and boars in Vietnam. These are clearly widely distributed, with endemic infections circulating in both domestic pigs and boars, and demonstrate the ease with which enteric viruses likely transmitted by faecal/oral routes can become established in and transmit within farmed animal populations. Whilst the pathogenicity of neither virus is clearly established in either this or previous studies, further genetic and epizootiological characterization of these viruses will contribute to our understanding of their transmission

patterns and the farming practices that affect their spread. Future studies may also provide new insights into the factors that underlie the transmission and dissemination of more specifically pathogenic viruses, such as rotaviruses, that present a much greater challenge to animal welfare and productivity.

METHODS

Sampling. Faecal samples were collected from 104 farms in four districts of Dong Thap province between February and May 2012. At each farm, 10 freshly voided samples from randomly selected healthy pigs and up to four samples from pigs with diarrhoea were collected. From this collection, a subsample of 198 samples, including 70 diarrhoeic samples, were screened in a previous study (Van Dung *et al.*, 2014). In the current study, we extended the screening of EVs to the 18 remaining diarrhoeic samples and 466 samples representing three to five healthy pigs per farm. In addition, 45 faecal samples collected in April 2014 from healthy farm-bred boars that were housed in typical livestock pens on six farms in Dak Lak province were also included. All 727 samples from pigs and boars were also screened for kobuviruses.

Screening and typing of EVs. RNA was extracted from faecal samples using a MagNA Pure 96 Viral NA small volume kit (Roche) and an automated extractor (Roche) as described previously (Van Dung *et al.*, 2014). Reverse transcription using SuperScript III reverse transcriptase (Invitrogen, UK) was performed according to the manufacturer's instruction. EVs were screened by a real-time PCR




Fig. 7. (a) Nucleotide sequence divergence within EV-G9, and between EV-G9 sequences and other EV-G types. Y-axis values show mean pairwise distances between 300-base windows across the genome, incrementing by 30 bases between fragments. (b) Grouping scan analysis of the sequence 734087-G9. Sequence fragments of 300 bases incrementing by 30 bases, 100 bootstrap replicates, were compared with sequences from other types. To show gene boundaries, the P2 and P3 regions are shaded grey and a scale representation of gene positions in the EV polyprotein is shown above the graphs.

protocol (Beld *et al.*, 2004). Typing of EVs from 177 positive samples, including 10 from diarrhoeic pigs, representing ranges of geography, C_t values and ages of infected pigs, was performed by amplification and sequencing of the VP1 region as described previously (Van Dung *et al.*, 2014). VP1 products of the nested PCR were sequenced on both strands using BigDye Terminator version 3.1 (Applied Biosystems) and primers for second-round PCR.

Screening and characterization of kobuviruses. Synthesized cDNA was screened for kobuviruses by nested PCR using primers targeting the 3D region of all known kobuviruses (Table S2). The

conditions for amplification (using GoTaq DNA polymerase; Promega) were 95 °C for 5 min, and 30 cycles of denaturation (94 °C, 30 s), annealing (54 °C, 30 s) and elongation (72 °C, 45 s). Kobuviruses in positive samples were characterized by amplification and sequencing of the VP1 region with similar amplification conditions for EV VP1 as described previously (Van Dung *et al.*, 2014) but using an annealing temperature of 58 °C in both rounds.

PKV viral load measurement. Using PKV sequences available from GenBank, primers and a probe targeting the 5' UTR region were designed to develop a real-time PCR protocol for measurement of

PKV viral load. The 200 bp target region in the 5' UTR amplified from a positive sample was cloned into pGEM-T Easy Vector (Promega). The linearized recombinant plasmid was used as template for transcription using a MEGAscript T7 transcription kit (Ambion) as previously described (McLeish *et al.*, 2012). RNA transcripts were used to assess the detection limit and efficiency of the assay before application on a subsample of 100 positive samples. These samples were selected to represent healthy and diarrhoeic pigs of all age groups, with the focus on the group with disease association (matured pigs). The specificity of the assay was checked with 20 randomly selected negative pig samples that were negative for PKV, but positive for EV. Each reaction of 20 µl total volume contained primers (PKV_332s, PKV_533a) and probe (PKV_505a_probe) at final concentrations of 0.4 and 0.2 µM, respectively, cDNA (5 µl), and 10 µl of SensiFAST Probe Hi-ROX kit (Bioline) Master Mix. After initial denaturation at 95 °C (5 min), amplification was performed in 45 cycles of 95 °C for 10 s and 60 °C for 35 s (acquisition step). Primers and probe used in this study for kobuviruses are listed in Table S2.

Complete genome sequencing of EVs. Primers were designed using VP1 sequences available from this study and reference sequences from GenBank to obtain nearly complete genomes of viruses representing EV-G8 and EV-G9 (Fig. S1, Table S3). Nested PCR amplifying regions from ~700 bp to 2.5 kb in length were performed. For each reaction, 6 µl extracted RNA was used for cDNA synthesis in combination with first-round PCR using a Superscript III One-Step RT-PCR System with Platinum *Taq* High Fidelity (Invitrogen). Then, 1 µl of the first-round reaction was used as template for the second-round PCR with amplification (using GoTaq DNA polymerase; Promega) conditions as follows: 95 °C for 5 min, and 30 cycles of denaturation (94 °C, 30 s), annealing (50 °C, 30 s) and elongation (72 °C, 60 s) in a thermal cycler. A SequalPrep long PCR kit (Life Technology) was used according to the manufacturer's instruction instead of GoTaq DNA polymerase to amplify fragments > 2 kb.

Sequence analysis. Sequences were imported into SSE (Simmonds, 2012) for alignment and calculation of sequence divergence values from reference sequences of known EV types. Maximum-likelihood phylogenetic trees were reconstructed as described in our previous study (Van Dung *et al.*, 2014) using the MEGA 5.2 software package (Tamura *et al.*, 2011) with 1000 bootstrap resamples and best-fitting models (Tamura–Nei, gamma distribution for EV VP4/VP2 and VP1, Tamura three-parameter model, gamma distribution for EV 5' UTR and 3D, general time-reversible gamma distribution with invariant sites for kobuvirus VP1). Nucleotide distance scans and grouping scans (Simmonds & Midgley, 2005) implemented in SSE were applied to identify recombination breakpoints. Analysis of phylogenetic groupings was performed by determining association index values (Wang *et al.*, 2001) using the program in SSE with 100 bootstrap replicates.

Statistical analyses. Fisher exact test, χ^2 -test and Mann–Whitney test incorporated in Minitab 17 were used to compare data between groups of interest when appropriate as specified.

ACKNOWLEDGEMENTS

We would like to thank the Subdepartment of Animal Health of Dong Thap and all the farmers who participated in the survey for their support. Work has been funded by the Vietnam Initiative on Zoonotic Infections (VIZIONS), part of the Wellcome Trust Major Overseas Programme (UK).

REFERENCES

- Alexandersen, S. & Donaldson, A. I. (2002). Further studies to quantify the dose of natural aerosols of foot-and-mouth disease virus for pigs. *Epidemiol Infect* **128**, 313–323.
- Alexandersen, S., Zhang, Z. & Donaldson, A. I. (2002). Aspects of the persistence of foot-and-mouth disease virus in animals - the carrier problem. *Microbes Infect* **4**, 1099–1110.
- Amimo, J. O., Okoth, E., Junga, J. O., Ogara, W. O., Njahira, M. N., Wang, Q., Vlasova, A. N., Saif, L. J. & Djikeng, A. (2014). Molecular detection and genetic characterization of kobuviruses and astroviruses in asymptomatic local pigs in East Africa. *Arch Virol* **159**, 1313–1319.
- Barry, A. F., Ribeiro, J., Alfieri, A. F., van der Poel, W. H. M. & Alfieri, A. A. (2011). First detection of kobuvirus in farm animals in Brazil and the Netherlands. *Infect Genet Evol* **11**, 1811–1814.
- Beld, M., Minnaar, R., Weel, J., Sol, C., Damen, M., van der Avoort, H., Wertheim-van Dillen, P., van Breda, A. & Boom, R. (2004). Highly sensitive assay for detection of enterovirus in clinical specimens by reverse transcription-PCR with an armored RNA internal control. *J Clin Microbiol* **42**, 3059–3064.
- Boros, Á., Pankovics, P. & Reuter, G. (2011). Characterization of a novel porcine enterovirus in domestic pig in Hungary. *Infect Genet Evol* **11**, 1096–1102.
- Boros, A., Nemes, C., Pankovics, P., Biró, H., Kapusinszky, B., Delwart, E. & Reuter, G. (2012a). Characterization of a novel porcine enterovirus in wild boars in Hungary. *Arch Virol* **157**, 981–986.
- Boros, A., Pankovics, P., Knowles, N. J. & Reuter, G. (2012b). Natural interspecies recombinant bovine/porcine enterovirus in sheep. *J Gen Virol* **93**, 1941–1951.
- Buitrago, D., Cano-Gómez, C., Agüero, M., Fernandez-Pacheco, P., Gómez-Tejedor, C. & Jiménez-Clavero, M. Á. (2010). A survey of porcine picornaviruses and adenoviruses in fecal samples in Spain. *J Vet Diagn Invest* **22**, 763–766.
- Chen, L., Zhu, L., Zhou, Y. C., Xu, Z. W., Guo, W. Z. & Yang, W. Y. (2013). Molecular and phylogenetic analysis of the porcine kobuvirus VP1 region using infected pigs from Sichuan Province, China. *Virol J* **10**, 281.
- Davis, M., Sagan, S. M., Pezacki, J. P., Evans, D. J. & Simmonds, P. (2008). Bioinformatic and physical characterizations of genome-scale ordered RNA structure in mammalian RNA viruses. *J Virol* **82**, 11824–11836.
- Di Bartolo, I., Angeloni, G., Tofani, S., Monini, M. & Ruggeri, F. M. (2015). Infection of farmed pigs with porcine kobuviruses in Italy. *Arch Virol* **160**, 1533–1536.
- Fan, S., Sun, H., Ying, Y., Gao, X., Wang, Z., Yu, Y., Li, Y., Wang, T., Yu, Z. & other authors (2013). Identification and characterization of porcine kobuvirus variant isolated from suckling piglet in Gansu province, China. *Viruses* **5**, 2548–2560.
- Frémont, M., Coomans, D., Massart, S. & De Meirleir, K. (2013). High-throughput 16S rRNA gene sequencing reveals alterations of intestinal microbiota in myalgic encephalomyelitis/chronic fatigue syndrome patients. *Anaerobe* **22**, 50–56.
- Inaba, M., Katayama, H., Tran, N. V. T. & Furuami, H. (2014). Detection of genus *Kobuvirus* for evaluation as virus indicator for fecal contamination source track from Nhue River in Hanoi, Vietnam. In *Southeast Asian Water Environment* **5**, pp. 61–66. Edited by H. Yamamoto, H. Furumai, H. Katayama, C. Chiemchaisri, U. Puetpaiboon, C. Visvanathan & H. Satoh. London: IWA Publishing.
- Kaku, Y., Sarai, A. & Murakami, Y. (2001). Genetic reclassification of porcine enteroviruses. *J Gen Virol* **82**, 417–424.

- Khamrin, P., Maneekarn, N., Kongkaew, A., Kongkaew, S., Okitsu, S. & Ushijima, H. (2009).** Porcine kobuvirus in piglets, Thailand. *Emerg Infect Dis* **15**, 2075–2076.
- Khamrin, P., Maneekarn, N., Hidaka, S., Kishikawa, S., Ushijima, K., Okitsu, S. & Ushijima, H. (2010).** Molecular detection of kobuvirus sequences in stool samples collected from healthy pigs in Japan. *Infect Genet Evol* **10**, 950–954.
- Knowles, N. J. (1988).** The association of group III porcine enteroviruses with epithelial tissue. *Vet Rec* **122**, 441–442.
- Knowles, N. J. (2006).** Porcine enteric picornaviruses. In *Diseases of Swine*, pp. 337–345. Edited by B. E. Straw, J. J. Zimmermann, S. D. D’Allaine & D. J. Taylor. Ames, IA: Blackwell.
- Krumbholz, A., Dauber, M., Henke, A., Birch-Hirschfeld, E., Knowles, N. J., Stelzner, A. & Zell, R. (2002).** Sequencing of porcine enterovirus groups II and III reveals unique features of both virus groups. *J Virol* **76**, 5813–5821.
- McIntyre, C. L., Knowles, N. J. & Simmonds, P. (2013).** Proposals for the classification of human rhinovirus species A, B and C into genotypically assigned types. *J Gen Virol* **94**, 1791–1806.
- McLeish, N. J., Witteveldt, J., Clasper, L., McIntyre, C., McWilliam Leitch, E. C., Hardie, A., Bennett, S., Gunson, R., Carman, W. F. & other authors (2012).** Development and assay of RNA transcripts of enterovirus species A to D, rhinovirus species a to C, and human parechovirus: assessment of assay sensitivity and specificity of real-time screening and typing methods. *J Clin Microbiol* **50**, 2910–2917.
- Moon, H. J., Song, D., Seon, B. H., Kim, H. K., Park, S. J., An, D. J., Kim, J. M., Kang, B. K. & Park, B. K. (2012).** Complete genome analysis of porcine enterovirus B isolated in Korea. *J Virol* **86**, 10250.
- Oberste, M. S., Maher, K., Kilpatrick, D. R. & Pallansch, M. A. (1999).** Molecular evolution of the human enteroviruses: correlation of serotype with VP1 sequence and application to picornavirus classification. *J Virol* **73**, 1941–1948.
- Oberste, M. S., Peñaranda, S. & Pallansch, M. A. (2004).** RNA recombination plays a major role in genomic change during circulation of coxsackie B viruses. *J Virol* **78**, 2948–2955.
- Okitsu, S., Khamrin, P., Thongprachum, A., Hidaka, S., Kongkaew, S., Kongkaew, A., Maneekarn, N., Mizuguchi, M., Hayakawa, S. & Ushijima, H. (2012).** Sequence analysis of porcine kobuvirus VP1 region detected in pigs in Japan and Thailand. *Virus Genes* **44**, 253–257.
- Park, S.-J., Kim, H.-K., Moon, H.-J., Song, D.-S., Rho, S.-M., Han, J.-Y., Nguyen, V.-G. & Park, B.-K. (2010).** Molecular detection of porcine kobuviruses in pigs in Korea and their association with diarrhea. *Arch Virol* **155**, 1803–1811.
- Prodělalová, J. (2012).** The survey of porcine teschoviruses, sapeloviruses and enteroviruses B infecting domestic pigs and wild boars in the Czech Republic between 2005 and 2011. *Infect Genet Evol* **12**, 1447–1451.
- Reuter, G., Boldizsár, A., Kiss, I. & Pankovics, P. (2008).** Candidate new species of Kobuvirus in porcine hosts. *Emerg Infect Dis* **14**, 1968–1970.
- Reuter, G., Nemes, C., Boros, A., Kapusinszky, B., Delwart, E. & Pankovics, P. (2013).** Porcine kobuvirus in wild boars (*Sus scrofa*). *Arch Virol* **158**, 281–282.
- Simmonds, P. (2006).** Recombination and selection in the evolution of picornaviruses and other mammalian positive-stranded RNA viruses. *J Virol* **80**, 11124–11140.
- Simmonds, P. (2012).** SSE: a nucleotide and amino acid sequence analysis platform. *BMC Res Notes* **5**, 50.
- Simmonds, P. & Midgley, S. (2005).** Recombination in the genesis and evolution of hepatitis B virus genotypes. *J Virol* **79**, 15467–15476.
- Simmonds, P., Tuplin, A. & Evans, D. J. (2004).** Detection of genome-scale ordered RNA structure (GORS) in genomes of positive-stranded RNA viruses: Implications for virus evolution and host persistence. *RNA* **10**, 1337–1351.
- Sisay, Z., Wang, Q., Oka, T. & Saif, L. (2013).** Prevalence and molecular characterization of porcine enteric caliciviruses and first detection of porcine kobuviruses in US swine. *Arch Virol* **158**, 1583–1588.
- Sozzi, E., Barbieri, I., Lavazza, A., Lelli, D., Moreno, A., Canelli, E., Bugnetti, M. & Cordioli, P. (2010).** Molecular characterization and phylogenetic analysis of VP1 of porcine enteric picornaviruses isolates in Italy. *Transbound Emerg Dis* **57**, 434–442.
- Tamura, K., Peterson, D., Peterson, N., Stecher, G., Nei, M. & Kumar, S. (2011).** MEGA5: molecular evolutionary genetics analysis using maximum likelihood, evolutionary distance, and maximum parsimony methods. *Mol Biol Evol* **28**, 2731–2739.
- Van Dung, N., Anh, P. H., Van Cuong, N., Hoa, N. T., Carrique-Mas, J., Hien, V. B., Campbell, J., Baker, S., Farrar, J. & other authors (2014).** Prevalence, genetic diversity and recombination of species G enteroviruses infecting pigs in Vietnam. *J Gen Virol* **95**, 549–556.
- Wang, T. H., Donaldson, Y. K., Brettell, R. P., Bell, J. E. & Simmonds, P. (2001).** Identification of shared populations of human immunodeficiency virus type 1 infecting microglia and tissue macrophages outside the central nervous system. *J Virol* **75**, 11686–11699.
- Yang, S., Wang, Y., Shen, Q., Zhang, W. & Hua, X. (2013).** Prevalence of porcine enterovirus 9 in pigs in middle and eastern China. *Virology* **457**, 99.
- Zell, R., Dauber, M., Krumbholz, A., Henke, A., Birch-Hirschfeld, E., Stelzner, A., Prager, D. & Wurm, R. (2001).** Porcine teschoviruses comprise at least eleven distinct serotypes: molecular and evolutionary aspects. *J Virol* **75**, 1620–1631.
- Zhang, W., Yang, S., Shen, Q., Ren, L., Shan, T., Wei, J., Cui, L. & Hua, X. (2012).** Complete genome sequence of a novel porcine enterovirus strain in China. *J Virol* **86**, 7008–7009.